


METHYL BROMIDE

**RISK CHARACTERIZATION DOCUMENT
FOR INHALATION EXPOSURE**

DRAFT



 California Environmental Protection Agency
Department of Pesticide Regulation

October 1999

DRAFT RCD 99-02

**CALIFORNIA ENVIRONMENTAL PROTECTION AGENCY
DEPARTMENT OF PESTICIDE REGULATION**

**Paul E. Helliker
Director**

For additional copies of this report please contact:

Department of Pesticide Regulation
Pesticide Registration Branch
830 K Street
Sacramento, California 95814-3510

(916) 445-4400

METHYL BROMIDE

**RISK CHARACTERIZATION DOCUMENT
FOR INHALATION EXPOSURE**

DRAFT

Medical Toxicology, Worker Health and Safety, and
Environmental Monitoring and Pest Management Branches
Department of Pesticide Regulation
California Environmental Protection Agency

October 15, 1999

CONTRIBUTORS AND ACKNOWLEDGMENTS

Principle Author: Lori O. Lim, Ph.D., D.A.B.T.
Staff Toxicologist (Specialist)
Medical Toxicology Branch

Toxicology Data Review: Stephen J. Rinkus, Ph.D.
Staff Toxicologist (Specialist)
Medical Toxicology Branch

Reviewers: Joyce F. Gee, Ph.D.
Peter Leung, Ph.D., D.A.B.T., and
Keith Pfeifer, Ph.D., D.A.B.T.
Senior Toxicologists, Medical Toxicology Branch

Jay Schreider, Ph.D.
Primary State Toxicologist, Medical Toxicology Branch

Gary Patterson, Ph.D.
Supervising Toxicologist, Medical Toxicology Branch

Additional Contributors:

Worker and Residential Exposures -

Thomas Thongsinthusak, Ph.D.
Staff Toxicologist (Specialist)
Worker Health and Safety Branch

Air Monitoring and Buffer Zones -

Pam Wofford, M.S.
Associate Environmental Research Scientist, and

Randy Segawa
Senior Environmental Research Scientist
Environmental Monitoring and Pest Management Branch

The Department of Pesticide Regulation acknowledges the review of this document by the Office of Environmental Health Hazard Assessment, California Environmental Protection Agency; the U.S. Environmental Protection Agency; and the National Academy of Sciences. A preliminary draft of the Toxicology Profile and Hazard Identification sections was reviewed by the U.S. Environmental Protection Agency and Dr. Gerald Last, Professor at the University of California at Davis.

TABLE OF CONTENTS

	Pages
List of Tables	iii
I. TECHNICAL SUMMARY	1
II. INTRODUCTION	15
III. TOXICOLOGY PROFILE	30
A. Pharmacokinetics	30
B. Acute Toxicity	34
C. Subchronic Toxicity	46
D. Chronic Toxicity and Oncogenicity	55
E. Genotoxicity	66
F. Reproductive Toxicity	71
G. Developmental Toxicity	74
H. Neurotoxicity	78
IV. RISK ASSESSMENT	82
A. Hazard Identification	82
B. Exposure Assessment	93
C. Risk Characterization	107
V. RISK APPRAISAL	119
A. Introduction	119
B. Hazard Identification	119
C. Exposure Assessment	122
D. Risk Characterization	123
E. Issues Related to the Food Quality Protection Act	126
VI. CONCLUSIONS	129
VII. REFERENCES	130
VIII. APPENDICES	156
A. Preliminary Risk Assessment for Methyl Bromide	157
B. Document for the Proposition 65 Developmental and Reproductive Toxicant Identification Committee	167
C. Methyl Bromide Residues in Fumigated Commodities	197
D. Toxicology Summary for Methyl Bromide	204
E. Chloropicrin Use and Toxicology	234
F. Estimation of Exposure of Persons to Methyl Bromide During And/Or After Agricultural and Non-Agricultural Uses	238
G. Calculations	353
H. Methyl Bromide Air Monitoring Studies and Buffer Zones	356
I. Description of Computer Modeling Procedures for Methyl Bromide ..	365
J. Evaluation of Potential Subchronic Exposure to Methyl Bromide Using Pesticide Use Report Data	376
K. Evaluation of Charcoal Tube and SUMMA Canister Recoveries for Methyl Bromide Air Sampling	382
L. Comments from the Office of Environmental Health Hazard Assessment and Response to Comments	445

LIST OF TABLES

	pages
1. The acute lethal toxicity of technical methyl bromide	35
2. Alteration of catecholamine levels and tyrosine hydroxylase activity in the rat brain after acute inhalation exposure to methyl bromide	39
3. The neurotoxicity of methyl bromide in dogs after acute exposure	42
4. The no-observed-effect levels (NOELs) and lowest-observed-effect levels (LOELs) for acute effects of methyl bromide from selected studies	45
5. The neurotoxicity of methyl bromide in dogs after subchronic exposure	52
6. The no-observed-effect levels (NOELs) and lowest-observed-effect levels (LOELs) for subchronic effects of methyl bromide	53
7. The effects of methyl bromide in rats after chronic inhalation exposure	57
8. The incidences of histological lesions in mice after chronic inhalation exposure to methyl bromide	61
9. The incidences of histological lesions in survivors of chronic inhalation exposure to methyl bromide	62
10. Hematological findings in dogs exposed to methyl bromide fumigated feed	64
11. The no-observed-effect levels (NOELs) and lowest-observed-effect levels (LOELs) of methyl bromide from chronic oral and inhalation studies	65
12. The genotoxicity of methyl bromide	69
13. Body weight changes in rats after inhalation exposure to methyl bromide in a 2-generation reproductive toxicity study	73
14. The incidence of fetal effects in rabbits after inhalation exposure to methyl bromide during gestation	76
15. The critical no-observed-effects levels (NOELs) for the risk characterization of inhalation exposures to methyl bromide	92
16. Estimates of occupational exposures to methyl bromide in preplant soil injection	96
17. Estimates of occupational exposures to methyl bromide in commodity fumigation facilities	99
18. Ambient and area sampling air concentrations of methyl bromide in or near commodity fumigation facilities	102
19. Residential exposures to methyl bromide in fumigated homes or from living near commodity fumigation facilities	105
20. Summary of occupational and residential exposures to methyl bromide	106
21. Margins of exposure for occupational exposures to methyl bromide in preplant soil injection	109
22. Margins of exposure for occupational exposures to methyl bromide in commodity fumigation facilities	112
23. Margins of exposure for exposure to ambient and area sampling air concentrations of methyl bromide in or near commodity fumigation facilities	114
24. Margins of exposure for residential exposures to methyl bromide in fumigated homes or from living near commodity fumigation facilities	117
25. Summary of margins of exposure for occupational and residential exposures to methyl bromide	118
26. Comparison of neurotoxicity in humans and dogs	124

I. TECHNICAL SUMMARY

I.A. INTRODUCTION

In 1992, the Department of Pesticide Regulation (DPR) conducted a Preliminary Risk Assessment on methyl bromide to address acute inhalation exposures of residents reentering fumigated home. The risk assessment concluded that the reentry level of 5 ppm for residents posed a health hazard and an emergency regulation was promulgated to decrease the exposure. Subsequently, permit conditions were developed to reduce the acute exposure of workers and the residents living near fumigated fields and fumigation chambers.

This Risk Characterization Document reevaluates the acute exposure assessment conducted in 1992 and considers also subchronic and chronic exposures to methyl bromide by inhalation from all uses in California. While all toxicology studies are reviewed, the potential risk from exposure to methyl bromide residues in food will be addressed in another document.

I.A.1. Chemical Identification

Methyl bromide is a gaseous fumigant that kills insects, rodents, nematodes, weeds, and organisms that cause plant diseases. Since methyl bromide is released into the air during and after its use, there is a potential for exposure by the workers as well the general population living near the use sites. Methyl bromide is a restricted use pesticide for structural, soil, and commodity fumigations. In 1999, 51 products containing methyl bromide were registered in California. From 1993-1997, 14-17 million pounds were used each year in California.

The primary route of human exposure to methyl bromide is inhalation. Exposure may occur from accidental spills, drift, leakage, or residual levels of methyl bromide released after treatment. Signs and symptoms of inhalation exposure depend on the concentration and exposure duration. Early symptoms of acute exposure to lethal concentrations include: malaise, headache, visual disturbances, nausea, and vomiting. Later symptoms include delirium, convulsions, and respiratory failure or cardiovascular collapse leading to death. Nonlethal exposures result in neurological effects similar to the early symptoms for fatal exposure. These symptoms may persist after exposure, depending on the severity of the effects. Exposure of skin to methyl bromide results in vesication and swelling of the skin. The general population may also be exposed to methyl bromide-treated foods.

Since methyl bromide is acutely toxic, chloropicrin has been added to some methyl bromide formulations as a warning agent. However, there may not be a correlation between methyl bromide concentration in the air and the extent of the irritation induced by chloropicrin due to the differences in physical and chemical properties between these compounds. Also, chloropicrin itself is acutely toxic. With its increased use as a replacement for methyl bromide, there are increased concerns regarding the health effects from chloropicrin exposure.

I.A.2. Regulatory History

The Federal agencies have established regulatory levels for the uses of methyl bromide. For food uses, the U.S. Environmental Protection Agency (U.S. EPA) established tolerances

based on inorganic bromide with the assumption that methyl bromide is completely degraded. The U.S. EPA oral chronic reference dose (RfD) is 0.0014 mg/kg/day. In the drinking water, the one-day, ten-day, and longer-term health advisories are 0.1 mg/L for children. The longer-term health advisory is 0.5 mg/L for an adult. The lifetime health advisory is 0.01 mg/L.

For methyl bromide in the air, the U.S. EPA inhalation reference concentration (RfC) is 5×10^{-3} mg/m³. The Agency for Toxic Substances and Diseases Registry minimum risk levels (MRLs) are 50 ppb, 50 ppb, and 5 ppb for acute, intermediate, and chronic exposure scenarios, respectively. For occupational exposure, the federal Occupation Safety and Health Administration permissible exposure limit (PEL) is 20 ppm while California established a lower limit of 5 ppm and a ceiling of 20 ppm. The reentry level is 1 ppm for structural fumigation within the wall voids. Methyl bromide is a Class I ozone depleter and its use is regulated by the U.S. Clean Air Act and the United Nations Montreal Protocol.

In California, the use of methyl bromide is regulated by permit conditions. To limit the acute exposures of workers and residents, the minimum buffer zones are 30 and 100 feet for worker and residents, respectively. The distances of the buffer zones vary depending on the use pattern. The need for the permit conditions was based on the Preliminary Risk Assessment conducted in 1992 (Appendix A). In 1993, methyl bromide, as a structural fumigant, was administratively listed as a developmental toxicant by the Office of Environmental Health Hazard Assessment of the California Environmental Protection Agency under Proposition 65 via the provision for listing due to the federal label warning requirement. However, the Proposition 65 Developmental and Reproductive Toxicity Identification Committee decided not to expand the listing to all uses of methyl bromide because results from laboratory animals did not “clearly” show that methyl bromide was a developmental toxicant.

I.A.3. Environmental Fate

Methyl bromide is degraded in the environment. The rate of hydrolysis was enhanced by elevated temperature, ultraviolet irradiation, aerobic conditions, and high organic matter in the soil. Once applied to the soil, methyl bromide volatilized into the air or adsorbed onto soil particles. Because of degradation, methyl bromide residues were not detected in the groundwater or commodities grown on fumigated soil. Residues were found in treated commodities after post-harvest fumigation.

I.B. TOXICOLOGY PROFILE

I.B.1. Pharmacokinetics

Pharmacokinetic studies showed that after inhalation, intraperitoneal, and oral administrations, methyl bromide was rapidly absorbed and radioactivity (¹⁴C) was distributed to all tissues. With inhalation exposure, the percentages of the administered doses absorbed were similar in several species; they were 48% in the rat, 40% in the dog, and 52 to 55% in human. In the rat, the highest levels in the tissues, principally in the lungs, were reached immediately after exposure. With oral and intraperitoneal administration to rats, more than 90% of the dose was absorbed, with the highest radioactivity levels measured in the liver, kidneys, and testes. Methyl bromide was extensively biotransformed into unidentified products and carbon dioxide. In the rat, within 1 hour after inhalation exposure, less than 10% of the radioactivity in the tissues was

intact methyl bromide. In humans, both methyl bromide and inorganic bromide were detected in the tissues 5 hours after a lethal dose exposure. The primary routes of excretion were the exhaled air for inhalation and intraperitoneal exposures, and the urine for oral exposure. Carbon dioxide accounted for almost 50% (inhalation and intraperitoneal routes), and 30% (oral route) of the radioactivity in the exhaled air. After oral administration, biliary metabolites of methyl bromide were reabsorbed from the gut.

I.B.2. Acute Toxicity

Methyl bromide is a Toxicity Category I compound because of its acute inhalation toxicity. Severe irritation to eyes, skin, and mucous membranes occur after acute exposure; therefore, acute oral, ocular and dermal studies are not required for registration. Neurotoxicity has been observed in humans and laboratory animals after inhalation exposure to methyl bromide. The severity of the effects depended on the dose and duration of exposure. In humans exposed to high concentrations, neurological effects included ataxia, convulsion, and tremors. The nonlethal effects observed in laboratory animals included changes in brain catecholamines and tyrosine hydroxylase activity, tissue degeneration (nasal, brain, and adrenal glands), and neurotoxicity (ataxia and paralysis). Signs of oral toxicity in the dog included prostration, increased heart rates, lesions in multiple organs including the stomach and brain, hypoactivity, hypothermia, and death. Human dermal exposure resulted in skin lesions.

I.B.3. Subchronic Toxicity

Subchronic inhalation exposure of laboratory animals to methyl bromide resulted in altered brain catecholamine levels, decreased brain tyrosine hydroxylase activity, neurotoxicity, tissue degeneration (brain, nasal cavity, heart, testes, adrenal glands, thymus, spleen, and kidneys), and death. Based on overt signs of neurotoxicity, the dog, rabbit, and monkey were more sensitive to methyl bromide than other species (rat, mouse, and guinea pig). The primary finding in the rat was hyperplasia of the forestomach after repeated oral exposure by gavage. A decrease in body weight gain and food consumption was observed in rats given micro-encapsulated methyl bromide mixed in the feed.

I.B.4. Chronic Toxicity

The nasal cavity, brain, and heart were major target organs in rodents after chronic inhalation exposure to methyl bromide. Olfactory epithelial damage (hyperplasia, metaplasia, and necrosis) and myocardial degeneration were observed in rats and mice. Cerebellar and cerebral degenerations were detected in mice while reduced brain weight was observed in rats. When rats were exposed to methyl bromide in microcapsules mixed in the feed, the primary effect was body weight reduction. Possible treatment-related lesions were found in the spleen, liver, pancreas, and lungs. In dogs given methyl bromide-fumigated feed, anemia and decreased serum calcium were observed.

I.B.5. Genotoxicity

Methyl bromide was genotoxic in several *in vitro* and *in vivo* assays. It was a base-pair substitution mutagen in the *Salmonella* assays. It was a direct-acting mutagen since a liver

S-9 fraction was not required for mutagenicity. It caused micronuclei formation in female mice and an increased frequency of sister chromatid exchanges in CHO cells and in mouse bone marrow cells *in vivo*. It did not induce unscheduled DNA synthesis in rat hepatocytes or cause sperm abnormalities in mice. DNA alkylation was detected in both rats and mice after *in vivo* exposure by oral, intraperitoneal, or inhalation routes while DNA damage was found in the germ cells of rats after inhalation exposure. There was some evidence of genotoxicity in workers exposed to methyl bromide. Elevated levels of sister chromatid exchanges in lymphocytes and S-methylcysteine adducts in the blood were measured in soil fumigators. An increased frequency of hypoxanthine-guanine phosphoribosyl transferase gene (*hprt*) mutations in the lymphocytes and an increased incidence of micronuclei in oropharyngeal cells were observed in structural fumigators.

I.B.6. Reproductive Toxicity

In a 2-generation reproductive toxicity study in rats by inhalation, methyl bromide reduced the fertility rate of the F₁ parents during the second mating trial. While the body weights of the treated pups at birth showed varied responses, their body weights were significantly lowered during lactation. Brain weight and cerebral cortex width were reduced in the F₁ parents.

I.B.7. Developmental Toxicity

Methyl bromide caused developmental effects in both rats and rabbits after inhalation exposure. The findings in the fetuses included delayed skull ossification in rats and fused sternbrae, gall bladder agenesis, and other effects in rabbits. Methyl bromide did not cause any significant developmental effects in rats and rabbits after oral exposure.

I.C. RISK ASSESSMENT

I.C.1. Hazard Identification

The evaluation of risks from exposure to methyl bromide followed the four steps of risk assessment: hazard identification, dose-response assessment, exposure evaluation, and risk characterization. In the hazard identification and dose-response assessment, a comprehensive review of the toxicology database from studies submitted by the registrant and published articles was conducted. From this review, the toxicity and the estimates of how much methyl bromide that could potentially cause an adverse effect as well as no-effect levels are identified for each study. Since human case reports did not provide sufficient details to derive the critical no-observed-effect levels (NOELs), results from experimental animal studies were used assuming that the effects observed in the animals would also be observed in humans. The NOELs were expressed as human equivalents (adult or child) to correct for the difference in respiration rates between humans and experimental animals. The studies with most relevant findings for risk assessment were then selected and the associated NOELs were expressed as critical NOELs to be used in the calculation of the margin of exposure (MOE) in the risk characterization step of the process. For methyl bromide, critical NOELs were determined for acute (one-time exposure), short-term (1-2 weeks), subchronic (more than one month, a season), and chronic (a year or more) exposures.

For acute exposure, neurotoxicity is the primary effect of concern and has been observed in both experimental animals and humans. The clinical signs observed include: decreased activity, ataxia, paralysis, convulsion, and tremors. Of the laboratory animals studied, there was a species sensitivity to the neurotoxicity of methyl bromide after short-term exposure. Based on the comparisons of the lowest-observed-effect level (LOEL) for neurotoxicity, the dog and rabbit showed greater sensitivity than the guinea pig, mouse and rat. For example, dogs exposed to 156 ppm (human equivalent level of 68 ppm) showed severe neurological effects in 2 to 7 days of exposure while rats exposed to the same concentration in terms of human equivalent level (65 ppm; 70 ppm actual air concentration) for the same exposure duration did not show any neurotoxicity. In pregnant animals, the rabbit was more sensitive to methyl bromide than the rat. For pregnant rabbits, severe neurotoxicity was observed at the LOEL of 70 ppm (Sikov *et al.*, 1981; Breslin *et al.*, 1990) while no neurotoxicity was reported in the pregnant rats at the same level (Sikov *et al.*, 1981).

The selection of results from the most sensitive species, in this case the dog, is consistent with the U.S. EPA Neurotoxicity Risk Assessment guidelines (U.S. EPA, 1998a). The critical NOEL was 103 ppm from short-term inhalation studies in the dog (Newton, 1994a and b). At this dose of 103 ppm, no effects were observed until the 8th day of exposure. Although the dog inhalation toxicity studies were not designed to be a neurotoxicity study as defined by the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) guideline, they were conducted under Good Laboratory Practices and DPR considered the results valid for hazard identification. These same data were used by the Methyl Bromide Industry Panel (MBIP) to support their position that a chronic inhalation toxicity study in the dog should not be required (CMA, 1994).

The selection of the 103 ppm dose as the acute NOEL considered three major factors: subjectiveness of the observations, severity of the neurotoxicity at higher concentrations, and possibility of delayed neurotoxicity. The finding of no effect at 103 ppm in the dogs was based on gross observations. Neurotoxicity may have been present but not detected unless more refined methods such as the Functional Observation Battery were used. Therefore it is possible that the actual NOEL may be lower than 103 ppm. Furthermore, severe neurotoxicity was observed at higher doses (1.5 times the NOEL) with a few additional days of exposure. At 156 ppm, one of two dogs showed lacrimation (tearing) on the first day. This finding by itself may arguably be considered less significant with respect to adversity. However, there were only two dogs in this group. With 2-3 days of additional exposure, there was significant toxicity as both dogs showed difficulty breathing and decreased activity. In another study, all dogs (8 in the group) exposed to 158 ppm showed decreased activity before the end of the second exposure day. With 5 additional days of exposure, all showed severe neurotoxicity and brain lesions. The selection of 103 ppm as the acute NOEL also addresses, indirectly, the possibility of delayed neurotoxicity which has been reported in humans after accidental poisonings. Since no effects were observed in the dogs at 103 ppm for 7 days of continuous exposure, it is unlikely that there would be delayed neurotoxicity within one week after a single exposure to the same level. The human equivalent NOEL (25 ppm) from the dog study was two-fold or less than those for the acute effects observed in the rats and guinea pigs.

Another endpoint DPR considered for acute exposure risk assessment is the developmental toxicity observed in experimental animals after methyl bromide exposure. In a developmental toxicity study, the pregnant animals were exposed continuously to methyl bromide during a specified period of gestation (when organ formation occurs). Any adverse

effect observed in the fetus is considered an acute effect under the current assumption that only a single exposure at a critical time is necessary for the induction of developmental adverse effects according to the U.S. Environmental Protection Agency Guidelines for Developmental Toxicity Risk Assessment. Since this endpoint is the result of exposure during pregnancy, it is only used for the assessment of exposure by women of childbearing age in the work force and the general population.

The critical NOEL for developmental toxicity was 40 ppm from a study with rabbits (Breslin *et al.*, 1990). The result from this study was the basis for the emergency regulation and permit conditions currently used in California. U.S. EPA also considered these endpoints of concern and has used the same study in a Section 18 evaluation on the use of methyl bromide on imported fruits at ports of entry. In this rabbit developmental toxicity study, fetuses exposed to 80 ppm *in utero* showed gall bladder agenesis (no gall bladders), fused sternebrae (early fusion of the sternebrae), and lowered body weights. The missing gallbladder finding was seen in Part I of the experiment, which by itself is a complete study and fulfilled FIFRA guidelines for an acceptable study. The investigator was concerned with the finding as it was rarely observed in the negative-control litters in the conducting laboratory as well as in other laboratories using the same rabbit strain. When the experiment (Part II) was repeated three months later, missing gall bladders were again observed in fetuses exposed to methyl bromide *in utero*. The fused sternebrae found in Part I was not confirmed since a skeletal examination was not performed in Part II.

The developmental toxicity effects observed in fetuses should not be discounted because of maternal toxicity (body weight changes and neurotoxicity) reported at the same dose level. Consideration must be given to when the effects were observed. First, the decrease in the body weight gain of the 80 ppm group does was not a consistent finding. Statistically significant decreases were reported for gestation days 13-16 in Part I and gestation days 7-20 and 10-13 periods in Part II. The reduced weight gain in the does of Part II occurred concomitantly with a reduction in the mean fetal body weight. Second, there was no significant difference in the strictly maternal parameter calculated as the terminal body weight minus gravid uterine weight. Third, body weight changes in pregnant rabbits are known to be more variable than in rodents. As a result, body weight changes often do not carry as much support as an indicator for maternal toxicity as for rodents as discussed in the U.S. EPA Developmental Toxicity Risk Assessment guidelines. Fourth, maternal neurotoxicity was characterized by clinical signs, including: lethargy, head tilt, slight ataxia and slight lateral recumbency. These signs were observed in only 3 of 43 does (7%) dosed at 80 ppm, and they did not appear until gestation days 19-20 (the last days of the 13-day exposure period). Based on the description and comparison with observations reported in other studies, DPR does not consider these signs as indicators of excessive toxicity.

Furthermore, the failure of gall bladders to form in some fetuses was independent of maternal neurotoxicity. In Part I, 6 of the fetuses with missing gallbladders were from 3 does without neurotoxicity while the remaining 7 affected fetuses were from 2 does with neurotoxicity. In Part II, none of the does showed neurotoxicity while 4 fetuses (from 4 does) had missing gallbladders. In addition, the development of the gall bladder in rabbits can be considered an acute event since it takes place in one to two days after its onset on gestation day 11.5 (Hoar and Monie, 1981). The maternal neurotoxicity reported on gestation days 19-20 would have occurred too late to have been a factor in the agenesis of the gall bladder.

Similar findings have not been reported in the rat developmental toxicity studies. While it is worth noting that rats do not have gall bladders, the absence of these findings in another species should not negate their significance as indicators of the potential for methyl bromide to cause developmental toxicity in humans. Species specificity in developmental effects has been demonstrated for some chemicals. Developmental toxicity testing under FIFRA guidelines requires two species to be tested, a rodent and a non-rodent species, typically the rabbit, for identifying species susceptibility. The need to test non-rodent species arose from the findings of thalidomide where it was demonstrated that this human teratogen did not exhibit significant teratological effects in rats but caused at least some significant effects in rabbits (Schardein, 1985). As stated in the U.S. EPA Developmental Toxicity Risk Assessment guidelines, developmental effects may not be evident in more than one species. The findings from the most sensitive species are appropriate to use to estimate human risk.

The significance of the developmental toxicity findings was discussed in a 1994 Proposition 65 meeting to determine whether methyl bromide should be listed for all uses. The emergency regulation in 1992 resulted in methyl bromide being listed as a chemical known to the State of California to be a reproductive toxicant. The Developmental and Reproductive Toxicity Identification (DART) Committee was presented with results from animal developmental toxicity (absence of gall bladders and fused sternbrae) and reproductive toxicity (decreased pup body weight) studies. After much discussion, the Committee voted not to expand the listing of methyl bromide from structural fumigation to all uses because there was not enough evidence to support the "clearly shown" criteria as mandated by the Proposition. However, the members expressed several concerns: the need for more experimental studies to clarify the findings, potential for exposure to methyl bromide via the milk during lactation, and the lack of information on human exposure especially during pregnancy.

After this meeting, DPR received additional data to support the consideration of reproductive or developmental toxicity as a pertinent endpoint for risk assessment and regulatory actions. First, supplemental data on the rat reproductive toxicity study showed that methyl bromide caused a reduction in the width of a certain part of the brain (cerebral cortex) in the F₁ adults exposed to methyl bromide *in utero* (American Biogenics Corp., 1986). Second, a study received by DPR in 1998 showed that methyl bromide caused a breakage of DNA in the testicular cells isolated from rats after inhalation exposure (Bentley, 1994). It is not known whether the effect was due to methyl bromide or a metabolite.

DPR also considered studies which showed that methyl bromide caused biochemical changes in the brain which may be associated with neurotoxicity. In the rat, acute exposure to methyl bromide has been shown to alter catecholamine (chemicals involved in the transmission of brain signals) levels and tyrosine hydroxylase (an enzyme involved catecholamine formation) activity in the brain. However, an extensive review of the published articles on this subject by DPR showed inconsistencies in the findings; thus, they were considered not appropriate for use in the determination of regulatory levels. The results of one of these study were used by the Agency for Toxic Substances and Diseases Registry of the Public Health Service to derive a minimum risk level as a screening tool for regulatory agencies to determine the need for regulation. As such, the study review did not critically analyze the results. This minimum risk level has not been adopted as an action level by any regulatory agency.

Therefore, two acute NOELs were selected to address the different human sub-populations. The NOEL of 40 ppm for developmental toxicity in the rabbit was most appropriate for workers and residents since women of child bearing age are in both groups. For children, the NOEL was 103 ppm for neurotoxicity in the dog. When these NOELs are converted to human equivalent NOELs taking into consideration of duration of exposure and the differences in the respiration rates between species, and between adults and children, the human equivalents were 21 ppm and 25 ppm, respectively, for developmental toxicity and neurotoxicity. The use of the lower human equivalent NOEL of 21 ppm compared to 25 ppm to address occupational and residential exposures would protect children from the effects of methyl bromide.

For short-term and subchronic exposures, neurotoxicity was also selected as the endpoint for the determination of the critical NOELs and was based on the same considerations as discussed for acute exposure. For short-term (1week) exposure, a NOEL was established to address the potential exposure of residents returning to fumigated homes, living near fumigated fields, and workers. The critical NOEL was 20 ppm based on neurotoxicity (convulsion, paresis) in the rabbit after exposure to 70 ppm for 1 week (Sikov *et al.*, 1981). Three of 26 does died after 9 to 10 days of exposure.

For subchronic exposures of longer duration (> 1 month, seasonal), the critical NOEL was an estimated NOEL (ENEL) of 0.5 ppm based on a lowest-observed-adverse-effect level (LOAEL) of 5 ppm for decreased responsiveness in two of eight dogs during a neurological examination after about 7 weeks of exposure (34 exposure days) and a default factor of 10 for the calculation of a NOEL from a LOAEL (Newton, 1994b). While the duration is shorter than the 13-week generally considered for subchronic exposure, it was chosen because of the endpoint (neurotoxicity) and species sensitivity (the dog is a more sensitive species than the rat to methyl bromide) considerations. It is possible that the NOEL may be lower if the dogs were exposed to methyl bromide for 13 weeks.

This ENEL was lower than the NOEL (3 ppm) for lowered body weights of rat pups from dams exposed to methyl bromide before mating and during part of the pregnancy in the reproductive toxicity study (American Biogenics Corp., 1986). Another study also showed a NOEL (estimated) of 3 ppm based on a dose-related decrease in brain weight at 30 ppm and higher concentrations in the female rats (Norris *et al.*, 1993 a and b). The brain weight was also significantly decreased in the 140 ppm male rats. This effect on the brain weight was considered biologically significant since the brain is a target organ of methyl bromide. The absence of neurotoxicity by Functional Observational Battery testing at the same dose (30 ppm) does not diminish the importance of the brain weight finding since the causes of the two effects are not necessarily related.

For chronic inhalation exposure, all chronic studies conducted with rodents (rats and mice), reproductive toxicity study, and subchronic dog inhalation toxicity study were considered in the determination of the chronic critical NOEL. After chronic inhalation exposure, tissue damage was noted in the nasal cavity, brain, and heart of rodents. The critical NOEL was an ENEL of 0.3 ppm based on a LOAEL of 3 ppm for the induction of an increase in the number of cells (hyperplasia) and change in cell type and function (degeneration) in the nasal cavity of rats after 24-29 months of exposure and a default factor of 10 for the calculation of a NOEL from a LOAEL (Reuzel *et al.*, 1987 and 1991). While the exposure duration was considered a life-time

for the rodents, the actual duration in the standard chronic toxicity studies is two years. Since humans may be exposed to methyl bromide on a yearly basis, not just one or two years in the lifetime, the NOEL from the chronic toxicity study after two years of exposure was, therefore, appropriate for use. This NOEL may underestimate the risk of repeated yearly exposure as there is evidence of cumulative toxicity, in particular, neurotoxicity. The LOEL (3 ppm) from this 29-month study for nasal olfactory epithelial damage (Reuzel *et al.*, 1987 and 1991) is further supported by the LOEL of 4 ppm from a 24-month study for lesions at the same site (Gotoh *et al.*, 1994). The U.S. EPA also used the same LOAEL from this study in the determination of the chronic reference dose (RfC).

The significance of the finding in the nasal cavity is that it showed methyl bromide not only injured the cells but also changed the normal function of the cells in the nasal cavity. Such damage may result in the loss of the animal's sense of smell. Tissue damage in other organs occurred at higher concentrations. With acute exposure to 200 ppm, the damage to the rat olfactory epithelium included epithelial disruption, fragmentation, and exfoliation (Hurtt *et al.*, 1988). Repair of the epithelium included replacement by a squamous epithelium, loss of sensory cells, and respiratory metaplasia (conversion of the olfactory epithelium to a ciliated respiratory type). In other short-term studies, the damage to the nasal epithelium was described as necrosis and degeneration (Eustis *et al.*, 1988) and dysplasia (NTP, 1992; Eustis, 1992). In the chronic inhalation toxicity study, nasal olfactory epithelial hyperplasia and degeneration were observed in the rat (Reuzel *et al.*, 1987 and 1991).

While the effect on the nasal cavity may generally be considered a finding confined to the rat due to anatomical considerations, it is not the case with methyl bromide. Dogs exposed to 156 ppm methyl bromide for only 6 days showed moderate to moderately severe olfactory degeneration (Newton, 1994b). In addition, the rodent studies are the only available studies to evaluate the chronic toxicity. The requirement for a non-rodent (dog) study was waived by DPR based on the evaluation of short-term studies in the dog which showed that a chronic study would have to be conducted at relatively low dose levels. For comparison, the ENEL of 0.3 ppm for nasal cavity effects when expressed as human equivalent level (0.1 ppm) was the same as the human equivalent level for neurotoxicity after subchronic exposure (ENEL of 0.5 ppm). This implied that the actual NOEL for chronic exposure if based on neurotoxicity could be lower than that based on the effects in the nasal cavity. However, it is not possible to extrapolate such a NOEL at this time because the subchronic NOEL was already an estimated NOEL based on a LOEL which was reduced by a 10-fold uncertainty factor.

The oncogenicity of methyl bromide can not be evaluated at this time because experimental studies either showed no increased incidence of tumors after treatment or the reports did not provide sufficient data to determine the incidences. There is evidence that methyl bromide causes damage to the genetic material in experimental animals and humans, which is generally considered to play a significant role in the process of tumor formation.

A summary of the critical NOELs for risk characterization is presented below:

Scenarios	Experimental NOEL	Human Equivalent NOEL ^a		Reference Concentration ^d	Effects in Animal Studies	Ref ^e
		Adult ^b	Child ^c			
Acute	40 ppm	21 ppm	na	210 ppb	Developmental toxicity (pregnant rabbit)	1*
	103 ppm ^f	45 ppm	25 ppm		Neurotoxicity (dog)	2
Subchronic 1 week	20 ppm	12 ppm	7 ppm	120 ppb(adult) 70 ppb (child)	Neurotoxicity (pregnant rabbit)	3
6 weeks	0.5 ppm (ENEL)	0.2 ppm	0.1 ppm	2 ppb (adult) 1 ppb (child)	Neurotoxicity (dog)	2
Chronic	0.3 ppm (ENEL)	0.2 ppm	0.1 ppm	2 ppb (adult) 1 ppb (child)	Nasal epithelial hyperplasia(rat)	4*

- ^{a/} Experimental NOELs were converted to human equivalents using equations in Appendix G. na= child equivalent NOEL were not calculated because the effects were observed in pregnant animals.
- ^{b/} The adult equivalent NOELs are appropriate to address worker exposures. They are also used for residential exposures when child equivalent NOELs were not calculated.
- ^{c/} The child equivalent NOELs are appropriate to address resident exposures (see footnote b).
- ^{d/} The reference concentration was the ratio of the human equivalent NOEL and a default uncertainty factor of 100 since the NOEL was derived from experimental animal studies.
- ^{e/} * indicates study was acceptable to DPR according to FIFRA guidelines. References: 1. Breslin *et al.*, 1990b; 2. Newton, 1994b; 3. Sikov *et al.*, 1981; 4. Reuzel *et al.*, 1987 and 1991.
- ^{f/} The NOEL and human equivalents are presented in this Table for comparison purposes only. They are not used for risk characterization.

I.C.2. Exposure Assessment

Exposure scenarios were developed in this document to address potential inhalation exposures from the use of methyl bromide. For each scenario, exposure levels were estimated for acute exposures. These levels were based on available monitoring studies conducted by DPR or provided by the registrants. When data were not available, default factors were used. The daily exposure levels based upon exposure duration were then used to estimate the short-term (7-day), seasonal (subchronic), and chronic exposures. Workers in pre-plant soil fumigation and residents in structural (home) fumigation were considered not to have chronic exposures to methyl bromide. While there is a potential for skin lesions from dermal exposure when the methyl bromide concentration in the air is relatively high (e.g. at 9000 ppm), there is insufficient absorption information to determine the absorbed dose.

For workers involved in the use of methyl bromide for preplant soil fumigation, the range of acute exposures was 0.6 ppb (shovelman of shallow shank nontarped-bed method) to 1659 ppb (tarp remover). Applicators, co-pilots, cutters, pullers, cultipackers, and other workers had exposures between these two levels. While the shovelman of shallow shank non-tarped-bed method had the lowest acute exposure (0.6 ppb), the shovelman for the shallow-shank tarp method had much higher exposures (high of 515 ppb and a mean of 147 ppb). Tarp removers of the latter method also had the highest exposures for all durations: 1659 ppb (mean of 835 ppb) for acute, 596 ppb for short-term, and 510 ppb for seasonal exposures.

For nursery and greenhouse workers, the use of the hot-gas method resulted in high exposure levels. The highest exposure was 724 ppb measured for the applicators of this method compared to 26-31 ppb for applicators not using such a method. The range of acute exposures were 1 (tarp remover of hot gas method) to 724 ppb (applicators of hot gas method). There were no seasonal or chronic exposures.

For workers involved in the fumigation of grain products, the range of acute exposures was 8 ppb (forklift driver of chamber) to 8458 ppb (aerator of sea container/trailer). These aerators also had the highest average exposures for short-term (4314 ppb), seasonal (3271 ppb), and chronic (2482 ppb) exposures. However, in comparison, aerators of tarp fumigation had much lower (less than one-tenth) exposures for all durations.

For workers involved in the fumigation of dried fruit and tree nut products, the range of acute exposures was 0.35 ppb (forklift operator of dried prunes in chamber) to 1463 ppb (worker involved in clear chamber for raisins). In general, fumigators and aerators had higher exposures than other workers. For workers involved in the fumigation of cherries for export, the range of acute exposures was 47 ppb (forklift driver) to 877 ppb (fumigator, start-up). As with dried fruit fumigation, the fumigator had the highest exposures.

Around non-certified chambers for walnuts, workers not directly involved in the application of methyl bromide had higher exposures than that (225 ppb) for the fumigator. The acute exposures of these workers ranged from 770 ppb (sorting worker) to 1789 ppb (cracking worker). For workers in a walnut processing plant, potential exposures were based on air sampling in work stations rather than specific tasks. Cleaning plant (448 ppb), vacuum chamber (636 ppb), and Dock 5 (500 ppb) had the highest methyl bromide levels for acute exposure compared to other areas. The lowest mean level (39 ppb) was measured in the sorting area.

For workers in a brewery, exposures were based on air concentrations at various locations. The exposure levels ranged from 28.9 to 173 ppb.

Workers in the rice warehouse had relatively low average exposures at 10 ppb and 7 ppb for acute and short-term exposures, respectively. The highest ambient air concentration was 55 ppm (mean of 32 ppb) for acute exposure. For other commodity fumigation facilities, high methyl bromide levels were generally detected near the chambers. For example, 13281 ppb detected in a leak check in a dried prune chamber, 2175 ppb in raisins aeration sheds, 1789 ppb in walnut vacuum chamber area, and 6800 ppb by the injection site in a walnut processing facility. Other areas in these facilities had various exposure levels probably depended on the distance from the source of methyl bromide, ventilation, and other factors. Each facility had areas with < 10 ppb methyl bromide.

For residential exposures, the ranges of acute exposure levels were: 406 to 2990 ppb for residents near houses during fumigation, 74 to 1064 ppb for residents downwind from a fumigated house during aeration, 210 ppb for residents after reentry into treated homes, and 210 ppb for residents living near commodity fumigation facilities. The highest exposure level was 2990 ppb measured for outdoor air 20 feet from a house during fumigation. Measurable levels of methyl bromide were detected in the indoor air of neighboring houses: 406 ppb (during fumigation) and 168 ppb (during aeration). Lower levels were detected during aeration process and declined with distance from the fumigated house. The levels were 1064 ppb at 10 feet, 208 ppb at 50 feet, and 74 ppb at 100 feet. The acute exposure for residents reentering fumigated homes and those living near commodity fumigation facilities was 210 ppb.

I.C.3. Risk Characterization

The NOEL at which adverse effects did not occur was used to assess the non-cancer hazard for potential human exposures to methyl bromide. The margin of exposure (MOE) was compared with a conventional benchmark level of 100.

The MOEs varied from <1 to greater than 1000 for workers. For the workers involved in the use of methyl bromide for preplant soil fumigation, the range of acute MOEs were 13 (tarp remover in shallow-shank tarp) to 35000 (shovelman of shallow shank nontarp method). Applicators, co-pilots, cutters, pullers, cultipackers, and other workers had exposures between these two levels. The methods with acute MOEs of less than 100 were: shallow-shank tarp (all workers), deep shank injection (disc driver), shallow shank non-tarped-bed (irrigation pipe tractor driver), and tarp shallow with Noble plow (cutter, puller, and some removers). The short-term MOEs for most of these workers were also less than 100. The MOEs for seasonal exposures were all less than 100, except for tractor driver for shallow shank tarped-bed fumigations, applicators of shallow shank non-tarped-bed fumigations, and end puller of tarp shallow with Noble plow shank fumigations.

For nursery potting soil workers, the use of the hot-gas method resulted in low acute MOEs of 29 and 63 for applicators and tarp venters, respectively. The MOEs were 158 to 2625 for workers not using such a method. However, the MOEs were greater than 100 for all workers after short-term exposures.

For workers involved in the fumigation of grain products, the range of acute MOEs was 2 (aerator of sea container/trailer) to 2625 (forklift driver of chamber). These aerators also had the lowest MOEs for short-term, seasonal, and chronic exposures. Applicators and aerators of tarp fumigation also had MOEs of less than 100 for all durations. The MOEs for forklift drivers of a container/trailer were 50 for seasonal and chronic exposures.

For workers involved in the fumigation of dried fruit and tree nut products, the range of acute MOEs was 14 (worker involved in clear chamber for raisins) to 60000 (forklift operator of dried prunes in chamber). For workers involved in the fumigation of cherries for export, the range of acute MOEs was 24 (fumigator, start-up) to 447 (forklift driver). The fumigators of start-up and closing-up had MOEs of 63 and 51, respectively, for short-term exposures. Short-term MOEs for other workers were greater than 100. For dried fruits, the fumigators and aerators had higher exposures than other workers; however, the acute and short-term MOEs were less than 100 only for these workers in dried prunes, raisins, and walnut fumigations. The

seasonal MOEs were less than 100 for almost all workers. For workers around non-certified chambers for walnuts, the acute MOEs ranged from 12 (cracking worker) to 263 (cleaning-fumigator). The MOEs for most of the workers, except a few, were less than 100 for all exposure durations. Depending on the location in a walnut processing plant, the range of acute MOEs for the workers were 33 (vacuum chamber) to 300 (sorting area). The MOEs were less than 100 for workers in the cleaning plant, Dock 5, and vacuum chamber areas for all duration of exposures.

For workers in a brewery, the acute MOEs ranged from 121 (area sample at entrance) to 840 (aerator). The MOEs were equal to or greater than 245 for short-term exposures.

In the rice warehouse, the MOEs were 2100 (acute) and 1714 (short-term) for workers in the rice warehouse with relatively low exposures. The acute and short-term MOEs were 382 and 522, respectively, for ambient methyl bromide concentrations. However, the MOE was only 13 for subchronic and chronic exposures. For exposure from working in commodity fumigation facilities, the acute MOEs near the chambers were 2 for leak check in a dried prune chamber, 10 in raisins aeration sheds, 12 in walnut vacuum chamber area, and 3 by the injection site of a chamber in a walnut processing facility. The acute and short-term MOEs were generally much greater than 100 for other areas in these facilities, except for walnut fumigation chambers where the MOEs were less than 100 for most areas.

The MOEs for residents living next to a house during fumigation were 7 (outdoor air) and 52 (indoor air) for acute, 13 (outdoor air) and 292 (indoor air) for short-term exposures. During aeration, only the MOE of 20 for outdoor air 10 feet from fumigated house was less than 100. The acute MOE for residents reentering fumigated homes and those living near commodity fumigation facilities or fumigated fields was 100 because the exposure was 210 ppb, the current regulatory limit. However, the MOEs were 20-78, 1, and 1 for short-term, seasonal, and chronic exposures, respectively.

I.D. RISK APPRAISAL

Certain limitations and uncertainties were incorporated into the hazard identification, exposure assessment, and risk characterization of methyl bromide. In the selection of the endpoints, these included: the use of developmental toxicity in rabbits as the critical endpoint with the assumption that methyl bromide may also cause developmental toxicity in humans, the use of the nasal cavity effect in the rat for chronic exposure, and the necessary extrapolation from the LOEL to estimate the subchronic and chronic NOELs.

The uncertainties involved in the estimation of occupational and residential exposures were due to the following factors: incomplete data from field studies, use of default physiological parameters and use patterns, and exposures based on limited monitoring. The uncertainties existed in both the monitoring portions as well as the modeling portions of the exposure estimates. These uncertainties included: the precision and accuracy of the sampling and analytical methods, influence of environmental factors on air concentrations, application variability, use of default weather conditions, and use of default assumptions in estimating air concentrations associated with overlapping applications. Actual exposure may be underestimated or overestimated because of these uncertainties.

For the risk characterization, the uncertainties included the use of uncertainty factors to address extrapolation of no-effects from the effect levels, data from experimental animals to humans, and intraspecies variations. Because of the known neurotoxicity of methyl bromide and the lack of data on developmental neurotoxicity, it would be prudent to incorporate an additional uncertainty factor to address the potential increased sensitivity of infants and children to methyl bromide. While this document addressed only inhalation exposure to methyl bromide, considerations should also be given to potential aggregate exposure from multiple routes.

I.E. CONCLUSIONS

The human health risk from potential inhalation exposure (occupational and residential) to methyl bromide was evaluated in this Risk Characterization Document. The critical toxicity endpoints were derived from experimental animals: developmental toxicity for acute exposure, neurotoxicity for short-term and seasonal exposures, and tissue damage to the nasal cavity for chronic exposures. For acute and chronic exposure endpoints, neurotoxicity was also considered in the determination of the critical NOELs.

The risks, expressed as the margins of exposure, were calculated for more than 180 exposure levels for workers involved in soil and commodity fumigations, and residents in the vicinity of structural and commodity fumigations. The acute exposure was based on the highest detected level when available. For longer exposure durations, the mean exposure level was used. The benchmark MOE traditionally considered as adequate for the protection of human health is a MOE of 100 based on an animal NOEL. Because of potential increased sensitivity of infants and children to methyl bromide, an additional uncertainty factor was recommended.

For exposure to methyl bromide, the MOEs varied from <1 to greater than 100,000. In general, the acute MOEs for workers close to the source of methyl bromide were less than 100. These workers included: copilots, shovelmen, tarp removers, puller, removers, fumigators, and aerators under certain work practices. In the commodity fumigation facilities, there was a wide range of MOEs for the monitored sites. Since these sites were specific to the particular facility, and the distance of the site to the source of methyl bromide is not known in most cases, it was not possible to make a conclusion about the magnitude of the MOEs.

For residential exposure from structural fumigation, the MOEs were less than 100 for residents living near (<50 feet) to a house being fumigated. For residents reentering fumigated homes and those living close to commodity fumigation facilities or fumigated fields, the acute MOE was 100 based on the permit conditions. However, the MOEs for short-term (1 week) exposures were less than 100.

The MOEs presented in this risk characterization document pertain only to the inhalation exposure from the air and does not include the concomitant exposures through dietary intake. It is essential that the significance of the MOEs be viewed in the context of the limitations and uncertainties discussed.

II. INTRODUCTION

A human health risk assessment for inhalation exposure to methyl bromide has been conducted because of adverse effects identified in chronic toxicity, oncogenicity, reproductive toxicity, developmental toxicity, genotoxicity, and neurotoxicity studies conducted with laboratory animals. The toxicity of methyl bromide in humans from occupational and accidental exposures also has been documented. This comprehensive risk assessment supersedes the Department of Pesticide Regulation (DPR) 1992 Preliminary Risk Assessment (Appendix A). The Preliminary Risk Assessment addressed only the acute and subchronic inhalation toxicity of methyl bromide. Some of the information in this document has been presented to the Proposition 65 Developmental and Reproductive Toxicant Identification Committee (Appendix B). Methyl bromide is regulated under the California Air Contaminant Act (AB 1807), The Food Safety Act (AB 2161), The Birth Defect Prevention Act of 1984 (SB 950), and Proposition 65. While the review of toxicology studies included all routes of exposure, the potential risk from dietary exposure to methyl bromide residues in the fumigated foods will be addressed in a separate document.

II.A. CHEMICAL IDENTIFICATION

Methyl bromide has been used commercially since the 1890s (review by Alexeeff and Kilgore, 1983; WHO, 1995). Early uses were as an anesthetic agent, methylating agent, refrigerant, fire extinguishing agent, and fumigant. Currently, methyl bromide is used as a multipurpose fumigant for pest control in structures (warehouses, ships, freight cars, and homes) and in post-harvest treatment of commodities (Farm Chemicals Handbook, 1998). It is also used in the preplant treatment of soil in fields and greenhouses to control nematodes, insects, weeds, bacteria, and fungi. Methyl bromide is the primary fumigant for quarantine use on commodities exported to other countries (Appendix C).

The primary effect of methyl bromide after inhalation exposure is neurotoxicity which has been observed in both humans and laboratory animals (detailed discussion of the studies is in **III. TOXICOLOGICAL PROFILE**, and Appendix D). The signs and symptoms are dependent on concentration and exposure duration (von Oettingen, 1946; Rathus and Landy, 1961; Greenberg, 1971; Grant, 1974; Anger *et al.*, 1986; Gehring *et al.*, 1991; Uncini *et al.*, 1990; De Haro *et al.*, 1997).

With dermal exposure, vesication and swelling of the skin and increased plasma bromide levels were observed in workers exposed to high concentration of methyl bromide (Butler *et al.*, 1945; Jordi, 1953; Zwaveling, *et al.*, 1987; Hezemans-Boer *et al.*, 1988).

II.A.1. Mechanism of Action

The exact mechanism for the toxicity of methyl bromide via the inhalation route is not known. Methyl bromide is an alkylating agent and has been shown to bind irreversibly to sulfhydryl groups *in vitro* (Lewis, 1948; Price, 1985). Gehring *et al.* (1991) suggested that the depletion of glutathione (GSH) observed in the study by Alexeeff and Kilgore (1985) was a result of methyl bromide-GSH conjugate formation. However, conjugated metabolites of methyl bromide have not been reported. The depletion of GSH may also be due to the inhibition of glutathione reductase activity (Jaskot *et al.*, 1988; Davenport *et al.*, 1992).

The effect of methyl bromide on brain GSH and GSH transferase has been proposed as the mechanism of neurotoxicity (Davenport *et al.*, 1992). Methyl bromide-induced locomotor effects (tremors, ataxia, and limb paralysis) may be a result of changes in brain GSH metabolism (Orlowski and Karkowsky, 1976). In a case study of two workers, methyl bromide conjugation with GSH by GSH transferase was hypothesized to account for the severity of neurotoxicity observed (Garnier *et al.*, 1996). Other investigators suggested that methyl bromide-induced neurotoxicity was due to the decrease of catecholamine (norepinephrine and dopamine) levels and the inhibition of tyrosine hydroxylase activity in the brain (Honma *et al.*, 1982, 1987, and 1991). The decrease in catecholamine levels may cause reductions in presynaptic neuronal activity, feeding, and body temperature. This effect would be consistent with the decreased locomotor activity and weight loss observed in rats. In addition, an increase in the sensitivity of dopamine receptors after methyl bromide exposure has been suggested as the mechanism for the neurotoxicity symptoms (hallucination, insomnia, and delusions) (Honma *et al.*, 1994). Other indications of neurotoxicity observed in the rats were conditioned taste aversion (Miyagawa, 1982) and increased thiopental-induced sleep time (Honma *et al.*, 1985).

The toxicity observed in human and animal studies with low concentrations of methyl bromide is likely due to methyl bromide *per se* and not bromide. First, methyl bromide is more toxic than bromide. The rat oral LD₅₀ (3,500 mg/kg) for sodium bromide is more than 15-fold higher than that (214 mg/kg) for methyl bromide (Smith and Hamburger, 1935; Danse *et al.*, 1984). For acute exposure, the lowest level of bromide in the blood that caused toxicity is 125 mg bromide/100 ml (Gosselin *et al.*, 1976). A no-effect level of 4 mg/kg based on electroencephalograph changes and increased thyroid activity in human volunteers and an average daily intake (ADI) of 1 mg/kg of sodium bromide has been proposed for humans (van Gelderen *et al.*, 1993). For chronic exposure to bromide, central nervous system effects (drowsiness, bizarre behavior, and hallucinations) were observed in a 60-year-old woman who ingested a bromide elixir daily for 7 years. The serum bromide level was 44.6 mEq/L (or 215 mg/100 ml based on 1 mEq=48 mg) (Blumberg and Neli, 1967).

Second, bromide level in the plasma, serum or urine is not an indicator of the severity of methyl bromide intoxication. The diet and past medicinal use of bromides contribute to the endogenous level of bromide (Harvey, 1985). Human studies and accidental poisoning cases showed that the bromide levels of affected individuals did not correlate with the symptoms (Drawneek *et al.*, 1964; Marraccini *et al.*, 1983; Squier *et al.*, 1992; Kishi *et al.*, 1991; Tanaka *et al.*, 1991). Hemodialysis did not alleviate the neurotoxic effects of methyl bromide in a worker accidentally exposed to methyl bromide while cleaning a rice silo (Moosa *et al.*, 1994). Hustinx *et al.* (1993) suggested that the severity of clinical signs is correlated more with previous exposures than serum bromide levels (more details about this study are in III.H. NEUROTOXICITY).

II.A.2. Chloropicrin

Chloropicrin, a lacrimator and a fumigant, is added to the formulations as a warning agent because methyl bromide is acutely toxic and is odorless. However, the efficacy of chloropicrin, which ranged from 0.25% to 67% depending on the formulation, has been questioned (WHO, 1995). Methyl bromide concentration may be 100-fold higher than that of chloropicrin because of the difference in vapor pressures and densities of these two compounds

(Van Assche, 1971). The vapor pressure and density for methyl bromide are 1380 mm Hg and 3.78, respectively, and those for chloropicrin are 18.3 mm Hg and 5.68, respectively. At 1.3 ppm chloropicrin, which causes irritation, the calculated methyl bromide level was 115 ppm. The air levels of these two compounds were measured simultaneously in a monitoring study conducted by the California Air Resources Board (Seiber *et al.*, 1987). After soil fumigation with methyl bromide (194 lbs/acre) and chloropicrin (95 lbs/acre), the maximum methyl bromide and chloropicrin levels were: 1133 ppt for methyl bromide and 681 ppt for chloropicrin in the ambient air, and 900 ppb for methyl bromide and 23.8 ppb for chloropicrin 20 yards from the field.

The use of chloropicrin in California has increased from 2.1 million to 2.8 million pounds from 1993 to 1997 (DPR, 1993-1997). Because of the increased use, in particularly as a replacement for methyl bromide, there is concern about the potential health effects from exposure. While an extensive review on chloropicrin is beyond the scope of this document for methyl bromide, a summary of use and toxicity information is provided in Appendix E.

II.B. REGULATORY HISTORY

The insecticidal activity of methyl bromide was first reported in 1932 (Le Goupil, 1932). Methyl bromide is a restricted use pesticide in the United States. Retail sale and uses are limited to certified applicators or persons under their direct supervision, and only for those uses covered by the applicator's certification.

II.B.1. Ozone Depletion

Methyl bromide is an ozone depleter with a calculated ozone depletion potential (ODP) of 0.7 (Watson *et al.*, 1992). In the 1994 Science Assessment of Ozone Depletion document, a panel of atmospheric scientists concluded that methyl bromide continued to be a significant ozone depleter with an ODP of at least 0.3 (NOAA/NASA/UNEP/WMO, 1995). The worldwide sources of methyl bromide include: anthropogenic (human made) agriculture (20-60 kilotons/year), biomass burning (forest fires, grass fires) (10-50 kilotons/year), leaded gasoline burning (0.5-22 kilotons/year), and oceans (60-160 kilotons/year). There is evidence that methyl bromide produced by open oceans is reabsorbed. Therefore, the release of methyl bromide from agriculture use is a significant source for human exposure. The amounts (as % applied) of methyl bromide used which eventually reach the atmosphere have been estimated to be: 50-95% for soil, 80-95% for post-harvest commodity, and 90% for structural uses (U.S. EPA, 1997a).

The concern of methyl bromide and other chemicals with ozone depleting potential (ODP > 0.2) is currently being addressed by the U.S. Clean Air Act and the United Nations Environment Programme Montreal Protocol on Substances that Deplete the Ozone Layer (Montreal Protocol). Under the U.S. Clean Air Act, the U.S. Environmental Protection Agency (U.S. EPA) was required in 1994 to freeze the U.S. production and importation of methyl bromide at the 1991 level (U.S. EPA, 1993). A complete ban was scheduled for the year of 2001. However, the 1998 U.S. Congress passed legislation to extend the use until 2005. Recently, U.S. EPA determined that pesticide applicators using methyl bromide would not be required to use impermeable tarps to reduce air emissions from field fumigation because not enough was known about how the use of the tarps would affect crop production (U.S. EPA, 1998b).

At the international level, the Parties (more than 125 nations) to the Montreal Protocol added methyl bromide to the list of depleters and agreed on deadlines for a freeze on the production and importation of methyl bromide (U.S. EPA, 1993). In 1997, Parties to the Montreal Protocol agreed to an extension of the use. The current deadlines for the 100% reduction in use are 2005 and 2015 for developed and developing countries, respectively (UNEP, 1997). With the potential cancellation of methyl bromide use, there are ongoing efforts to develop alternative approaches (GAO, 1996; USDA, 1993; CDFA, 1995; Braun and Supkoff, 1994).

II.B.2. Federal Regulations

The U.S. EPA established tolerances in commodities based on bromide level because of the assumption that methyl bromide is degraded completely to bromide (Federal Register, 1991). However, residue studies have shown that fumigated commodities contain detectable levels (in ppm range) of methyl bromide especially immediately after fumigation. In 1986, the Methyl Bromide Industry Panel (MBIP) petitioned the U.S. EPA for tolerances for methyl bromide *per se* (U.S. EPA, 1986a). The proposed levels ranged from 0.1 ppm for certain vegetables to 5.0 ppm for green cocoa beans.

The U.S. EPA has established an oral chronic reference dose (RfD) and inhalation reference concentration (RfC) for methyl bromide (U.S. EPA, 1992a). The RfD is 0.0014 mg/kg/day based on the no-observed-effect level (NOEL) of 1.4 mg/kg/day for forestomach epithelial hyperplasia in a rat oral subchronic study (Danse *et al.*, 1984) and an uncertainty factor of 1000. The inhalation RfC is 5×10^{-3} mg/m³ (1.3 ppb) based on the LOAEL of 3 ppm for nasal olfactory epithelial hyperplasia from a rat chronic inhalation study (Reuzel *et al.*, 1987 and 1991) and an uncertainty factor of 100.

For methyl bromide in the drinking water, the one-day, ten-day, and longer-term health advisory for a child is 0.1 mg/L assuming 1 L/day water consumption for a 10-kg child (U.S. EPA, 1992a). The longer-term health advisory for an adult is 0.5 mg/L assuming 2 L/day water consumption for a 70-kg adult. The lifetime health advisory is 0.01 mg/L assuming 20% of exposure by drinking water. Methyl bromide is classified as a "Group D" carcinogen (not classifiable as to human carcinogenicity) by U.S. EPA due to inadequate human and animal data (U.S. EPA, 1992a).

The Agency for Toxic Substances and Disease Registry (ATSDR) has established minimal risk levels (MRLs) for methyl bromide which may be of concern at hazardous waste sites and releases (ATSDR, 1992 and 1996). The acute MRL is 0.05 ppm based on an acute NOEL of 16 ppm for decreases of brain tyrosine hydroxylase activity in the rat after 8 hours of exposure to 31 ppm (Honma *et al.*, 1987 and 1991) and adjusted for daily exposure and a 100-fold uncertainty factor for interspecies and intraspecies extrapolation (16 ppm x 8/24 x 1/100). The intermediate inhalation MRL is 0.05 ppm (5 ppm x 1/100) based on a NOEL of 5 ppm for a decrease in monoamines in the rat after 3 weeks of continuous daily exposure to 10 ppm (Honma *et al.*, 1982). The chronic MRL is 0.005 ppm (2.3 ppm x 1/10 x 8/24 x 5/7 x 1/10) based on increased prevalence of muscle aching and fatigue, increased sensitivity threshold, and lowered recall ability in an epidemiological study of workers with an average exposure of 2.3 ppm for 8 hours per day and 5 days per week (Anger *et al.*, 1986) and an uncertainty factor of 100 for extrapolation of LOEL to NOEL and intraspecies variation. Respiration rate differences

between animals and humans are not accounted for in these MRLs. The intermediate oral MRL is 0.003 mg/kg/day based on gastrointestinal effects.

The National Institute for Occupational Safety and Health has determined an Immediately Dangerous to Life and Health Concentration (IDLH) for immediate evacuation of workers exposed to methyl bromide (NIOSH, 1997). The IDLH of 250 ppm was supposedly based on an acute inhalation toxicity data in humans (Clarke *et al.*, 1945). However, the air concentration was not measured in the cited human report. In the introduction, 220 ppm was cited as lethal in rats after 22 hours of exposure (no reference given). The federal Occupational Safety and Health Administration permissible exposure level (PEL) is 20 ppm (CFR, 1989). The American Conference of Governmental Industrial Hygienists (ACGIH) recommends a threshold limit value of 1 ppm (ACGIH, 1998).

II.B.3. California Regulations

For occupational exposure to methyl bromide, the current California permissible exposure limit (PEL) for methyl bromide is 5 ppm or 20 mg/m³ and a ceiling limit of 20 ppm (California Code of Regulations, 1998). A reentry level of 5 ppm following fumigation of homes was established under the Label Improvement Program for Fumigants, PR Notice 85-6, August 30, 1985 (U.S. EPA, 1986b). All persons are required to wear a self-contained breathing apparatus (SCBA) or combination air-supplied/SCBA respirators at concentrations higher than 5 ppm. In 1992, an evaluation of the monitoring data and toxicology studies showed that the 5 ppm reentry level in fumigated homes did not provide a sufficient safety margin for human exposure (DPR, 1992; Appendix A Preliminary Risk Assessment). The DPR promulgated emergency regulations to require a longer aeration period and lowered the reentry level to 1 ppm in the wall voids, and pest control operators had to hand out a Fact Sheet explaining the potential human hazards of methyl bromide fumigation. The Fact Sheet was prepared by the DPR in consultation with the Office of Environmental Health Hazard Assessment (OEHHA), California Department of Health Services, and U.S. EPA. A similar warning for developmental effects was subsequently required by U.S. EPA on methyl bromide product labels used for structural fumigation (U.S. EPA, 1992b).

On January 1, 1993, methyl bromide was administratively listed by OEHHA as a developmental toxicant under Proposition 65 via the provision for listing due to the federal label warning requirement. On December 21, 1993, OEHHA modified the listing from "methyl bromide" to "methyl bromide as a structural fumigant" because the Fact Sheet and label changes were limited to the use of methyl bromide as a structural fumigant. The Proposition 65 Developmental and Reproductive Toxicity Identification (DART) Committee of the OEHHA Science Advisory Board decided methyl bromide should not be listed for all uses since the evidence was considered equivocal^a and did not support the "clearly shown" criteria as mandated by the Proposition (OEHHA, 1994) (additional discussion under **IV. RISK ASSESSMENT A.1.a. Developmental Toxicity**).

^a As a reference, the definition for equivocal by the National Toxicology Program is "marginal evidence which may be chemical related."

II.B.4. Restricted Materials Permit Conditions

Methyl bromide when used in agricultural production is classified as a restricted material. Possession and use of restricted materials are allowed only under a permit from the county agricultural commissioner. Before issuing a permit, the county agricultural commissioner must evaluate the application to determine whether it will cause environmental harm. Depending on the results of this review, the commissioner may deny the permit or impose permit conditions including the use of specified mitigation measures. In evaluating permit applications, commissioners must consider and, where appropriate, use information provided by DPR. For methyl bromide, DPR provides this information as suggested permit conditions. In 1993, DPR began developing new use practice restrictions for agricultural applications that would incorporate a one-day (24-hour) exposure level of 0.21 ppm for workers and public. The suggested permit conditions include equipment modifications, restrictions on work hours, limits on application rates, limits on acreage or volume treated, tarpaulin specifications, restricted entry requirements, enclosed space requirements, and establishment of buffer zones. The minimum buffer zones for workers and residents are 30 feet and 100 feet, respectively. These buffer zones are currently being reviewed and revised to account for a change in the recovery rate calculation for the methyl bromide air concentration (Helliker, 1999; Appendix K).

II.C. TECHNICAL AND PRODUCT FORMULATIONS

In 1999, 51 methyl bromide-containing products were registered in California. The registrants of methyl bromide products are Ameribrom, Inc., Albemarle Corporation, Great Lakes Chemical Corporation, Soil Chemicals Corporation, Trical, Shadow Mountain Products Corp., and Bromine Compounds Ltd. The products are available as 100% methyl bromide and as mixtures with chloropicrin (range of 0.5% to 67% chloropicrin). Methyl bromide is used as a fumigant on raw and processed agricultural commodities, in structures, in soil, and on ornamentals.

II.D. USAGE

There are two major types of methyl bromide fumigations: applications to an area of soil or applications to an enclosed volume (e.g., chamber, building). All applications to soil occur prior to planting the crop for control of a wide variety of pests. Most soil applications involve the injection of methyl bromide beneath the soil surface. Injection can either be shallow (12 inches or less) or deep (20 inches or more), depending on the crop to be planted. In many cases, but not all, the soil is covered with a plastic tarpaulin to retard methyl bromide off-gassing. For typical agricultural field applications, a tractor injects the methyl bromide through a set of chisels or shanks, and lays the tarpaulin all at the same time. Applications can either be to a flat field (broadcast application) or to a field that has already had the beds and furrows formed (bed application). For a bed application, methyl bromide is applied only to the beds; the furrows are untreated.

Applications to post-harvest commodities or other enclosed volumes are very different. The types of enclosures and their efficiency in retaining methyl bromide are quite varied. The more typical enclosures include chambers, transportation containers, tarpaulins, and buildings. Regardless of the enclosure, most applications follow a general three-step process: 1) introduction or injection of methyl bromide into the fumigated space—this step usually takes a

few minutes; 2) a treatment or holding period—usually a few hours to a few days; 3) aeration of the fumigated space—also a few hours to a few days. The method of aeration can also be quite varied. Most chambers will have an elevated exhaust stack and release methyl bromide at a controlled rate, but tarpaulin applications are aerated simply by removing the tarpaulin.

From 1993-1997, 14-17 million pounds of methyl bromide were used each year for soil (approximately 85% of total), commodity and nursery (11-14%), and structural (3-4%) fumigations in California (DPR, 1993-1997). The major uses (% of total pounds used) in soil fumigation were strawberry fields (23-27%), preplant-outdoor sites such as seed beds (6-15%), grape orchards (9-13%), and nursery (11-14%). The use of methyl bromide in structural fumigation has declined significantly (< 5% of total use) in the last few years because of regulation that required extended aeration time before residents are permitted to reenter the homes.

II.E. ILLNESS REPORTS

Poisonings by methyl bromide have been reported as early as 1899 (von Oettingen, 1946). A review of the incidents published from 1939-1981 showed 115 incidences of fatalities and 843 incidences of nonfatal systemic or local injuries (Alexeeff and Kilgore, 1983).

In California, illnesses associated with methyl bromide have been reported and are described in detail in Appendix F. Approximately 50% of the cases also involved another compound or compounds. In the work place, the most common cause of the illnesses was equipment failure. Exposure of the general population to methyl bromide was due primarily to its release from fumigation chambers, leakage from fumigated fields, or off-gassing in fumigated homes.

Specific case studies and human epidemiological studies are presented in the **III.H. NEUROTOXICITY.**

II.F. PHYSICAL AND CHEMICAL PROPERTIES^b

Chemical name:	bromomethane, monobromomethane
CAS Registry number:	74-83-9
Common name:	methyl bromide
Trade names:	Brom, Brom-O-Gas, M-B-R, Metabrom, Meth-O-Gas, Methyl Bromide, Pic-Brom, Terr-O-Gas, Tri-Brom, Tri-Con, Tri-Pan.
Molecular formula:	CH ₃ Br
Molecular weight:	94.95 g/mole
Chemical structure:	CH ₃ -Br
Physical appearance:	colorless gas, usually odorless; sweetish, chloroform-like odor at high concentrations (odor threshold at 80 mg/m ³ or 20.6 ppm); burning taste. It is nonflammable in air but does burn in oxygen.
Solubility:	1.75 g/100 g in water (20EC, 748 mm), forms a crystalline hydrate, CH ₃ Br·20 H ₂ O, below 4EC; freely soluble in alcohol, chloroform, ether, carbon disulfide, carbon tetrachloride, benzene.
Boiling point:	3.56EC
Melting point:	-93.66EC
Octanol/Water partition coefficient:	log P=1.19 (15.5:1 octanol:water)
Vapor pressure:	1420 mm Hg (20EC), 2600 mm Hg (40EC)
Specific gravity:	3.3 g/ml (gas), 1.7 g/ml (liquid)
Conversion factor:	1 ppm = 3.89 mg/m ³ at 25EC

^b Farm Chemical Handbook, 1998; The Merck Index, 1989; U.S. EPA, 1986b

II.G. ENVIRONMENTAL FATE

Summary: Methyl bromide is degraded in the environment. The rate of hydrolysis was enhanced by elevated temperature, ultraviolet irradiation, aerobic conditions, and high organic matter in the soil. Once applied to the soil, methyl bromide volatilized into the air or adsorbed onto soil particles. Because of degradation, methyl bromide residues were not detected in the groundwater or commodities grown on fumigated soil. Residues were found in treated commodities after post-harvest fumigation.

II.G.1. Hydrolysis

Methyl bromide hydrolyzed to bromide in water (WIL, 1985a). The hydrolytic half-life was estimated to be 6 days for pH 9 at 35EC to 49 days for pH 7 at 25EC. There was leakage of methyl bromide from the flasks used in the experiments.

II.G.2. Photolysis or Photodegradation

Methyl bromide in solution (1.7×10^{-2} M) completely hydrolyzed to methanol, bromide, and hydrogen ion when irradiated at 254 nm for 40 days (Castro and Belser, 1981). The photolysis rate constant was $2 \times 10^{-6} \text{ sec}^{-1}$, 6.6-fold higher than without irradiation. However, experiments with incubation in the dark and in sterilized water were not performed.

In the upper stratosphere, methyl bromide was photo-dissociated, lost by diffusion, and reacted with hydroxyl radical (WHO, 1995). The products of the 3 reactions were probably carbon dioxide, carbon monoxide, and bromide species.

Photodegradation of methyl bromide was rapid in moist air (Gennari *et al.*, 1995). The half-lives were 22 minutes with irradiation at 250-420 nm and 15 mg/L moisture to 326 hours with irradiation at 366-750 nm and 15 mg/L moisture.

II.G.3. Microbial Degradation

There was no difference in the methyl bromide degradation half-lives between sterilized and unsterilized soils because methyl bromide decreased the microbial activity in both soils (Radian Corp., 1988). However, the half-lives under aerobic conditions were shorter than those for anaerobic conditions. The degradation in clay loam, with a higher organic content, was faster than in sandy loam. The average half-lives for sandy loam and clay loam soils were 35 and 3.8 hours, 47 and 2.5 hours, 144 and 39 hours, and 80 and 34 hours, under aerobic/ nonsterile, aerobic/sterile, anaerobic/nonsterile, and anaerobic/sterile conditions, respectively.

II.G.4. Mobility (Soil, Ground Water, Air, Plants)

II.G.4.a. Soil

The mobility of methyl bromide in soil, as with other soil fumigants, depends on many factors: chemical and adsorptive characteristics of the fumigant, temperature, moisture, organic matter, soil texture, and soil profile variability (Munnecke and Van Gundy, 1979; Kolbezen *et al.*, 1974; Abdalla *et al.*, 1974). The following section is a summary of some studies.

Thirty-two days after exposure to 200,000 ppm in the air, the soil contained less than 1.5 ppm methyl bromide (Radian Corp., 1988). This level was decreased by 10-fold after the soil was purged with nitrogen as methyl bromide was loosely associated with the soil. The proposed reaction of methyl bromide in the soil was: $\text{CH}_3\text{Br} + \text{O}_2 \rightarrow \text{HBr} + \text{H}_2\text{O} + \text{CO}$, then $2\text{CO} + \text{O}_2 \rightarrow 2\text{CO}_2$, with CO=carbon monoxide and CO_2 =carbon dioxide.

Under laboratory conditions, volatilization was the major route of dissipation from the soil (WIL, 1985b). The initial loss rate was $82.6 \text{ ug/cm}^2/\text{hour}$ and the steady state rate was $8.8 \text{ ug/cm}^2/\text{hour}$. Under similar conditions, the rate of loss for chloropicrin was much slower with a steady state rate of $1.8 \text{ ug/cm}^2/\text{hour}$.

Methyl bromide decomposed to bromide in all soil types (sand, peat, and loam) tested (Brown and Rolston, 1980). Decomposition was greatest in peat soil containing the most organic matter as the methyl group was transferred to the carboxyl groups and N- and S-containing groups of the amino acids and proteins of soil organic matter. In loam soil, the decomposition reaction was first order depending on the air concentration. The retention of methyl bromide in soil depended on soil water content. In dry sand, the adsorption and desorption rates of methyl bromide were similar. In moist sand, the desorption was slower (30% of adsorption) than adsorption.

Methyl bromide in water adsorbed to Canfield silt loam, Holly silt loam, and Wooster silt loam more than agricultural sand (WIL, 1986). The methyl bromide concentrations were 18 ppm in Canfield, 17 ppm in Holly, and 14 ppm in Wooster loam soils; but only 6 ppm was in sand. When the treated soil was mixed with water, 89 to 97% of the adsorbed methyl bromide were released from the soil and the equilibrium was reached in 24 hours.

Methyl bromide concentration and half-lives in Reiff (fine sandy loam) and Yolo loam were similar after preplant injection (6-8 inches deep) into the soil and then tarped, (MBIP, 1986). The highest concentration was at the level (1 foot) of injection. The dissipation half-life was the fastest at 1 foot (0.9-1.2 days) and the slowest at 3 feet deep (4.1 to 5.9 days). With deep injection of 18-24 inches and nontarped preplant injection, the dissipation half-lives ranged from 2.8 days at 1 foot to 9.6 days at 8 feet deep. When methyl bromide-treated soil was drenched with water and steamed upon removal of the tarp, methyl bromide was detected as far as 2 feet below the surface. The dissipation half-lives ranged from 2.5 to 3.0 days for the application rate of 1 lb/100 square feet, and 2.5 to 11.0 days for 2 lb/100 square feet.

Methyl bromide was retained after application beneath concrete slabs (McKenry and Secara, 1990). Methyl bromide (128 kg) was introduced into the ground via holes drilled through the floor and then the floor was tarped for 4 days. After treatment, the highest concentration was approximately 9,000 ppm on day 2 at 30 cm from the house and 90 cm deep underneath the floor. The concentration in the soil declined as the distance from the injection point was increased; however, even at 150 cm away, the concentration was as high as 5,700 ppm on day 3. The use of an air compressor to deliver air to the soil air space on day 15 enhanced the dissipation of methyl bromide. The concentration of methyl bromide at 15 cm beneath the slab remained below 21 ppm and averaged 8 ppm one week after the final air compressor treatment. With continuous aeration, the methyl bromide concentration in the open portion of the house did not exceed 88 ppb. The ambient concentrations of the backyard and neighboring houses were below 6 ppb during treatment and aeration.


II.G.4.b. Ground Water

Methyl bromide was not detected (detection limit of 1 ppb) in ground water samples from fields (muck and sandy type soils) in Florida (Pickrell *et al.*, 1985) and wells in Florida and California (Golder Associates, Inc. 1985a and b). The wells in California had a history of more than 10 years of methyl bromide use with the last fumigation 5 to 17 months before the study (Golder Associates, Inc. 1985b).

II.G.4.c. Air

The major sources of bromide and methyl bromide in the air are marine aerosols and natural marine biological processes, respectively (Lovelock, 1975; Wofsy *et al.*, 1975). Methyl bromide from fumigation uses was estimated at 25% of the total emissions (natural and anthropogenic) (Watson *et al.*, 1992). A minor source of methyl bromide is the exhaust from cars that use leaded gasoline (Harsch and Rasmussen, 1977). Measurements of methyl bromide over the Atlantic Ocean showed average concentrations of 15.4 ± 1.9 parts per trillion (ppt) and 10.6 ± 0.9 ppt for the Northern and Southern Hemispheres, respectively (Penkett *et al.*, 1985). The calculated total lifetime in the atmosphere was 1.16 years (Lobert *et al.*, 1995). In a survey conducted in 1979, the mean levels of methyl bromide in the ambient air were 244 ppt in Los Angeles, CA; 66.8 ppt in Phoenix, AZ; and 54.7 ppt in Oakland, CA (Singh *et al.*, 1981). Worldwide environmental levels are in the review by WHO (1995).

Methyl bromide degradation in the air involved reactions with hydroxyl radicals and ozone. Based on an estimated hydroxyl radical concentration of 2×10^6 /cm³ in the daytime, the estimated residence time in the air was 289 days with a daily rate of loss of 0.4% (Singh *et al.*, 1981). In the winter months when hydroxyl radical concentration could be lowered, the residence time would increase. The bromide from the degradation reactions reacted with ozone by the following reaction (Wofsy *et al.*, 1975; Watson *et al.*, 1992).



The release of methyl bromide after structural fumigation was monitored in close proximity and nearby fumigated homes (Gibbons *et al.*, 1996 a and b). During fumigation, air levels ranged from less than 0.019 to 1.495 ppm when monitored at 10 feet from the fumigated houses. At the neighboring houses, the air levels ranged from 0.012 ppm (detection limit) to 0.351 ppm. During aeration, the concentrations inside the houses at 50 and 100 feet downwind from the fumigated house depended on the aeration method and time of sampling. Compared to the Standard Method, the Pest Control Operators of California Aeration Method resulted in lower concentrations (7.5 ppm versus 13.6 ppm) at the 10-foot distance after 1 hour of aeration, but not those after 24-hour aeration, or at greater distances.

Liscombe, *et al.* (1995 and 1996) tested alternative tarping and aeration methods to mitigate methyl bromide air concentrations from structural fumigations. During treatment, air concentrations ranged from 0.035 to 1.08 ppm five feet from the fumigated structure, depending on the method of tarping. During aeration, air concentrations ranged from 0.021 to 0.11 ppm. Adding a second tarpaulin and aerating through an elevated stack were the most effective methods for reducing air concentrations.

Prior to 1992, limited monitoring of field fumigations was conducted. Methyl bromide levels in the air after tarpless fumigation of soil were determined in four farms (Soil Chemicals Corp., 1990). The shanks were placed 10 to 12 inches deep into the soil and the results are summarized below:

<u>Farms</u>	<u>Sampling site</u>	<u>Time</u>	<u>Methyl bromide level</u>
Belridge	60 ft downwind	several hours after fumigation	<30 ppb to 114 ppb
Kirschenman	150 ft downwind	0-11 hours after fumigation	<30 ppb (all times)
Major	200 ft downwind	during fumigation	<30 ppb to 211 ppb
Firini	in field	during fumigation	126 ppb
Firini	200 ft downwind	during fumigation	<30 ppb

In a monitoring study conducted by the Air Resources Board, Cal/EPA, air samples were collected on 3 off-sites close to strawberry fields and 4 sites in the nearby towns before and after application of methyl bromide (Seiber *et al.*, 1987). The fumigated field was tarped for 4 days. Methyl bromide levels were all below the MDL (1.1 ppb) for samples of ambient air in nearby towns. Methyl bromide was found in 3 off-sites (275, 76, and 175 meters from the field). The maximum levels (210-900 ppb) were detected on the day after application with a rapid decline on the next day. The highest average air concentration for an approximately 24-hour period was 450 ppb for site B, based on three 3-hour measurements. This level was used in the 1992 DPR Preliminary Risk Assessment (Appendix A).

Beginning in 1992, DPR, methyl bromide registrants, and academic researchers began more comprehensive monitoring of field and commodity fumigations. Details of the studies and the results are presented in Appendices F and H. The results were used to develop buffer zones for these uses.

II.G.5. Plant Residues/Metabolism

Methyl bromide residue in the food is generally analyzed by head-space gas chromatography (King *et al.*, 1981). This method requires the blending of macerated commodities in water. After the slurry is allowed to stand, methyl bromide which partitioned into the headspace is measured. The major uncertainty of this method is the time required for partitioning to reach an equilibrium that depends on the commodity composition and extent of methyl bromide interaction with endogenous components. The review of studies in the following section does not include residue data recently (July, 1998) submitted to the DPR as part of the methyl bromide dietary exposure assessment.

II.G.5.a. Soil Fumigation

Methyl bromide *per se* is generally not found in commodities grown on soil pretreated with methyl bromide because of its dissipation and degradation (MBIP, 1988a; Trical, Inc., 1985, 1990, and 1991; WIL, 1984). When methyl bromide was injected into the soil 15 days before planting and the field was tarped for 6 days, low levels of methyl bromide were found (Trical, Inc., 1985). The sampled crops and levels were: lettuce (< 3 ppb-29 ppb), turnip roots (< 23 ppb-644 ppb), broccoli (< 250 ppb-1452 ppb), carrots (<16 ppb), green tomatoes (< 16 ppb),

green beans (< 10 ppb). With cucumbers, two of the four samples contained residues at 5 and 34 ppb. In turnip leaves, methyl bromide levels ranged from < 9 ppb to 334 ppb; the control samples were all < 9 ppb. Matrix interference in the analysis was especially notable in broccoli and turnip roots. The levels of methyl bromide in two processed foods, tomatoes and dried beans, were below detection limits (10 ppb, and 6 ppb, respectively). Residues were also not detected in potatoes (< 5 ppb) and carrots in other studies (Trical, Inc., 1990 and 1991).

In a survey of methyl bromide residues after either broadcast application or bed treatment fumigation, 4 samples from each crop from 9 states (California, Florida, Oregon, Washington, Texas, Michigan, Arizona, Idaho, Hawaii, Wisconsin, and North Carolina) were analyzed (MBIP, 1988a). Most of the samples were collected in California. The crop groups and representative crops sampled were: root and tuber (carrot, potato, radish, and sugar beet); bulb vegetables (green onion, large bulb onion, small bulb onion, and garlic); leafy vegetables (head lettuce, leaf lettuce, celery, and spinach); Brassica vegetables (broccoli, cabbage, cauliflower, and mustard greens); legume vegetables (bush bean, green bean, soybean, succulent pea, and dry pea); fruiting vegetables (tomato, and pepper); cucurbit vegetables (cucumber, zucchini, watermelon, cantaloupe, and summer squash); citrus fruits (orange and grapefruit); pome fruits (apple); small fruits (raspberry, blueberry and strawberry); non-grass animal feed (alfalfa and clover); herbs and spices (basil, chives, dill, and marjoram); and miscellaneous (asparagus, ginger, grape, peanut, pineapple, corn, tobacco, and okra).

Almost all the samples contained residue levels below detection limits (ranged between 5 to 10 ppb). For mustard greens, the residue levels were 20.20 to 29.9 ppb (mean=24.9 ppb) for samples from California. In the Florida trial of cabbage, two of the four samples were 13.8 and 15.18 ppb and the other two were below the MDL (10 ppb). Since most of the samples did not contain residues, the levels above the MDL could be false positives due to dimethyl disulfide or other sulfur compounds. Bromide levels were determined in some crops and in general, were not indicative of methyl bromide application rates. For example, the mean bromide level in treated celery (442 ppm) was 4 times that of control (105 ppm). For pineapple, strawberry, and asparagus, bromide levels in the control and treated samples were similar (20 to 40 ppm).

II.G.5.b. Post-harvest Fumigation

The use of methyl bromide as a post-harvest fumigant on raw agricultural commodities and processed foods resulted in detectable levels of residues (Appendix C). Only residue data with adequate information of fumigation conditions and analysis method are included in Appendix C. In commercial practices, the fumigation condition is dictated by the pests of concern and the potential phytotoxicity by methyl bromide. In general, fresh fruits and vegetables are fumigated at short intervals (2 hours), and then stored at cool temperature.

The residue levels in fumigated foods were dependent on the nature of the commodity; load factor (volume of commodity to chamber size); application rate, temperature, frequency, and duration; as well as aeration and storage conditions. The half-lives of methyl bromide in fruits and vegetables were shorter than those for nuts and dried fruits. With cherries, there was an inverse relationship between load factor and initial residue levels such that an increase in the load factor from 1.6% to 32.0% resulted in lower residue levels (Sell *et al.*, 1988). Consequently, a shorter aeration period was needed to achieve a certain residue level. In addition, an increase in pulp temperature enhanced the desorption of methyl bromide from the cherries. For example,

the aeration period required to obtain 1 ppm methyl bromide remaining on the cherries was reduced from 18 hours to 9 hours when the pulp temperature was increased from 7EC to 27EC.

Likewise, an increase in temperature during fumigation or during aeration or storage resulted in the decrease of half-lives of methyl bromide in blueberries, walnuts, and strawberries (Appendix C). While an increase of the application rate from 2 to 4 lbs/1000 ft³ (plum, nectarine, or peach) or a doubling of exposure duration (avocado, peach, or pear) resulted in higher initial residue levels, there was no significant change in the half-lives. However, a 4-fold increase in an application rate (mango) resulted in an increase of both the initial residue levels and half-life.

Aeration lowered methyl bromide residues in fumigated strawberries (MBIP, 1984b). After 45 min, 1 hour, 2 hours, and 3 hours of aeration, the methyl bromide levels were 9.0 ppm, 7.0 ppm, 5.4 ppm, and 140 ppb, respectively. Samples which had been aerated for only 45 minutes before precooling contained 7.1 ppm of methyl bromide. The level then declined to 2.1 ppm and 1.6 ppb after 12 and 24 hours, respectively. Samples with 3 hours of aeration contained only 20-30 ppb after 2-3 hours at 34EF (1EC). No methyl bromide (MDL not specified) was detected in fumigated strawberries stored at 70EF (21EC) instead of 34EF (1EC). For asparagus, the mean residue level decreased from 2.14 ppm after fumigation to 0.016 ppm after 24 hours of aeration (Fieser and Conrath, 1993).

Because of the higher lipid content, the half-lives of methyl bromide in nuts (almonds, pistachio nut, and walnut) were longer than those for fruits (Appendix C). Methyl bromide absorbed into the almond shell and the kernel by dissolution of methyl bromide in the oil particles (Hartsell *et al.*, 1983). Some residues were released during aeration while others methylated amino acids or phenolic materials such as tannin and lignin. The shell generally contained higher levels of bromide and methyl bromide residues than the kernel. The highest bromide level was found in the Mission variety of almonds with the highest lignin and tannin contents, while lower levels were found in the Mercedes, Carmels, and Nonpareils varieties. Increase of temperature resulted in increased residues in the shell and kernels. During aeration, methyl bromide evaporated more rapidly from the shell than the kernel, but reached a similar concentration (< 2.5 ppm) in both compartments by about 50 hours.

Methyl bromide deposition in avocados was also related to the fat content (Singh *et al.*, 1982). The Fuerte variety absorbed 57% more methyl bromide and formed 22% more bromide than the Haas variety. Methyl bromide (21-36% conversion in Fuerte and 56-70% in Haas) was converted to bromide during storage.

Processing of fumigated grains reduced methyl bromide levels in the final product (MBIP, 1984a and 1988b; CMA, 1984). When wheat was processed to wheat flour, the levels ranged from below the detection limit (<1 ppb) to 2 ppb. For corn, grinding and baking at 350EC for 1 hour reduced the fumigated level (9.4 ppm) to 6.4 ppm and <0.01 ppm, respectively (MBIP, 1984a). Methyl bromide level in rice decreased from 5.2 ppm to < 0.01 ppm after 20 minutes of simmering. All flour and bakery mix samples had < 0.03 ppm methyl bromide.

Methyl bromide residues were higher (as much as 2-fold) in packaged dried fruits (dates, apricots, prunes) than in bulk form, except raisins (Appendix C). Because packaging provided limited off-gassing, the half-lives of methyl bromide in these commodities were also longer than those for bulk forms.

The dissipation half-lives provide only a qualitative estimate of the potential exposure by the consumers since multiple fumigations can occur to prevent reinfestation during processing and storage, either as bulk or packaged goods. A survey of nuts at the retail level showed methyl bromide levels ranged from 10 ppb to 3.7 ppm in 8 of 75 samples (Lindsay, 1985).

The U.S. Food and Drug Administration (FDA) collected samples of processed nuts, dried fruits, dried beans, and rice from the marketplace and analyzed them for methyl bromide (FDA, 1990 and 1991). Of the 1132 nut samples, only 2 pistachios samples contained detectable levels of residues (0.09 ppm and 0.03 ppm). Residues were not detected in 107 samples of rice, 347 samples of dried fruits, and 173 samples of dried beans; the MDL was 0.02 ppm (Ford *et al.*, 1992).

In a field-trial study, selected commodities were fumigated with methyl bromide at rates which ranged from 2-5 pounds/ 1000 ft³ for 2 to 24 hours according to the label uses (DFA, 1985; MBIP, 1985a). Methyl bromide residue levels were determined after 1, 3, or 7 days of aeration. The crop groups and selected commodities included: pome fruits (apple, pear), tree fruits (almond, walnut, and pistachios), root and tuber vegetables (carrot, potato, and sugar beet), legume vegetables (beans, peas, and soy beans), citrus fruits (orange, lemon, grapefruit), stone fruits (peach, plum), cereal grains (corn, rice, wheat), bulb vegetables (onion, garlic), fruiting vegetables (tomato, pepper), small fruits and berries (blackberry, blueberry, grape, and strawberry), herbs and spices (basil, chives, dill, and sage), leafy vegetables (broccoli and cabbage), cucurbits (cucumber, melon, and summer squash), processed (wheat, corn, rice), and miscellaneous (cocoa beans, dried fruits, and candy).

In wheat grown in sand containing ³⁵S-labeled sulfate and subsequently fumigated, the gluten or protein fraction contained N-methyl derivatives, dimethylsulfonium derivatives, as well as methoxy- and methylthio- derivatives (Winteringham *et al.*, 1955). Further study with fumigated wheat flour showed that methyl bromide methylated the basic nitrogen residues (Bridges, 1955). The histidine residue was the primary site of methylation resulting in the formation of 1-methylhistidine, 3-methylhistidine and 1,3-dimethylhistidine (Bridges, 1955). Studies with diets containing high levels of methylated histidines (0.36% of the diet) showed that these residues are not available for nutritional use resulting in weight loss in mice (Friedman and Gumbmann, 1979).

II.G.5.c. Structural Fumigation

Foods remaining in the house during fumigation may also acquire methyl bromide residues (Scheffrahn *et al.*, 1992). Packaged foods (unopened or reclosed) were fumigated with methyl bromide at approximately maximum labeled rates. Of the 23 food items, the highest residues were found in fatty commodities such as peanut butter (106 ppm) and margarine (151 ppm). Residue levels were below the detection limit in foods unopened with the factory seal intact or vacuum-packed.

III. TOXICOLOGY PROFILE

Pharmacokinetic and toxicity studies of methyl bromide are summarized in this section. Acceptability of the studies (except genotoxicity studies) by DPR, where noted, was based on the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) guidelines. The acceptability of the genotoxicity studies was based on the Toxic Substances Control Act guidelines (Federal Register, 1985 and 1987). The toxicology summary for studies reviewed for The Birth Defect Prevention Act of 1984 (SB 950) is included in Appendix D. In the toxicity studies, the no-effect levels may be expressed as NOELs or no-observed adverse effect levels (NOAELs). For the purpose of this document, endpoints under either designation are considered relevant for hazard identification.

Summary tables for selected toxicity studies considered for critical no-observed-effect levels (NOELs) and lowest-observed-effect levels (LOELs) are in Tables 4, 6, and 11 for acute, subchronic, and chronic exposures, respectively. Unless stated, concentrations are as measured concentrations from the studies. For comparison of toxicity between studies in these tables, the NOELs were also presented in terms of human equivalent NOEL. This human equivalent NOEL takes into consideration the air concentration, duration of exposure, and intake rate (respiration rate for inhalation exposure) differences between experimental animals and humans (adult and child) using equations in Appendix G. Since the experiments were conducted for various durations, the human equivalent NOEL is calculated on a per day basis to allow comparisons between studies. This approach follows the dose calculation methods outlined in the 1992 U.S. EPA Exposure Assessment guidelines, where the potential dose is a function of the concentration and intake rate (U.S. EPA, 1992c). It has generally been used for dietary exposure studies where the no-effect level is expressed as the dose (for example as mg/kg/day) accounted for consumption rate and duration of exposure, instead of concentration in the diet. Since the equivalent NOEL for children is lower than that for adults, only the children equivalents NOELs are presented in the summary tables. The only exceptions are the NOELs for developmental or reproductive toxicity studies since the endpoints are applicable only to adult exposures.

The internal doses for inhalation studies were assumed to be due to inhalation only. The inhalation pharmacokinetic studies were conducted with nose-only exposure while the toxicity studies (except where noted) were done with whole-body exposures. For laboratory animals in whole-body inhalation exposure studies, additional exposure was possible from oral ingestion due to licking of the fur and by dermal absorption. The internal dose from whole-body exposure may be higher than that calculated based only on the air concentration and respiration rates.

III.A. PHARMACOKINETICS

Summary: After inhalation, intraperitoneal, or oral administrations, methyl bromide was rapidly absorbed and radioactivity (¹⁴C) was distributed to all tissues. With inhalation exposure, the percentages of the administered doses absorbed were similar in several species; they were 48% in the rat, 40% in the dog, and 52 to 55% in humans. In the rat, the highest levels in the tissues, principally in the lungs, were reached immediately after exposure. With oral and intraperitoneal administration to rats, more than 90% of the dose was absorbed with the highest radioactivity levels measured in the liver, kidneys, and testes. Methyl bromide was extensively biotransformed into unidentified products and carbon dioxide. In the rat, within 1 hour after

inhalation exposure, less than 10% of the radioactivity in the tissues was intact methyl bromide. In humans, both methyl bromide and inorganic bromide were detected in the tissues 5 hours after a lethal dose exposure. The primary routes of excretion were the exhaled air for inhalation and intraperitoneal exposures, and the urine for oral exposure. Carbon dioxide accounted for almost 50% (inhalation and intraperitoneal routes), and 30% (oral route) of the radioactivity in the exhaled air. After oral administration, biliary metabolites of methyl bromide were reabsorbed from the gut.

III.A.1. Absorption

Fischer-344 rats were exposed to methyl bromide (^{14}C , >98% pure; nominal concentrations of 1.6, 9.0, 170, or 310 ppm) by nose only inhalation for 6 hours (Medinsky *et al.*, 1985). For 1.6, 9.0, and 170 ppm, the percentages of the absorbed dose (the ratio of radioactivity in the whole body homogenate after 6 hours to the amount inhaled x 100%) were 48%, 48%, and 38%, respectively. At 310 ppm, there were significant ($p \neq 0.05$) decreases in the tidal and minute volumes resulting in the absorption of only 27% of the total dose. At 66 hours after exposure, 17 to 26% of the total absorbed radioactivity remained in the body.

In beagle dogs, the calculated steady-state fractional, systemic uptake of the total inhaled methyl bromide (^{14}C , >98% pure, 174 to 361 ppb) was 39.5% after exposure for 3 hours by nose only, (Raabe, 1986). After the three-hour exposure, the blood concentration was 1.6% of the total amount inhaled and it declined to 1.2% at 117 hours after exposure.

In a human study, four volunteers (2 male and 2 female adults) were exposed to methyl bromide (^{14}C , >98% pure, 18 ± 6 ppb) for 2 hours by nose breathing and by mouth breathing (Raabe, 1988). Each subject was equipped with a two-stage demand regulator-based inhalation system that separated inhaled and exhaled air. The calculated steady-state, fractional systemic uptake of the total inhaled methyl bromide during nasal breathing and mouth breathing were 55.4% and 52.1%, respectively.

For oral and intraperitoneal administration in rats, the absorption was >90% when determined 72 hours after exposure (Medinsky *et al.*, 1984).

III.A.2. Distribution

After inhalation exposure to methyl bromide, radioactivity was found in the liver, kidneys, adrenal glands, lungs, thymus, brain, testes, and nasal turbinates in the rat (Medinsky *et al.*, 1985; Bond *et al.*, 1985; Jaskot *et al.*, 1988). The peak radioactivity levels in the tissues occurred immediately after exposure. The tissues with high radioactivity included the lungs, liver, and the nasal turbinates. In the study by Jaskot *et al.* (1988), the half-lives of tissue radioactivity were 2.71 hours (lung), 1.58 hours (liver), 0.58 hours (kidney), 1.37 hour (spleen), 6.54 hours (brain), and 5.29 hours (testes). In another study, the half-life of the radioactivity in the liver was 33 hours, while the half-lives were shorter in blood (7.7 hours), and in testes, small intestine, and brain (6 hours) (Bond *et al.*, 1985).

In a poisoning case, a man ingested and inhaled an unknown amount of methyl bromide for approximately 1.5 hours and died 4 hours later (Michalodimitrakis *et al.*, 1997). Tissues were sampled 1 hour after death during the autopsy. Except for the spleen, methyl bromide was

found in all tissues. The methyl bromide levels were: subclavian blood (3.8 $\mu\text{g/ml}$), brain (3.5 $\mu\text{g/g}$), adrenal gland (3.4 $\mu\text{g/g}$), peripheral blood (3.3 $\mu\text{g/ml}$), lung (2.9 $\mu\text{g/g}$), testis (2.8 $\mu\text{g/g}$), kidney (2.6 $\mu\text{g/g}$), liver (1.9 $\mu\text{g/g}$), bile (1.2 $\mu\text{g/g}$), and epididymis (1.2 $\mu\text{g/g}$). The inorganic bromide levels were: subclavian blood (530 $\mu\text{g/mL}$), brain (30 $\mu\text{g/mL}$), peripheral blood (480 $\mu\text{g/ml}$), lung (410 $\mu\text{g/g}$), kidney (310 $\mu\text{g/kg}$), and bile and epididymis (none).

After intraperitoneal or oral administration of methyl bromide (^{14}C , >98% pure, 250 $\mu\text{mol/kg}$ or 23.7 mg/kg), the radioactivity levels (in decreasing order) in Fischer-344 rats tissues were: liver > kidney, testes > lung, heart, stomach > spleen (Medinsky *et al.*, 1984).

III.A.3. Biotransformation/Excretion

Methyl bromide was rapidly biotransformed and readily excreted in rats after inhalation exposure (Bond *et al.*, 1985). In all tissues examined, over 90% of the radioactivity was metabolites. The elimination half-life of radioactivity from the tissues was 1.5 to 8 hours. Almost 50% of the absorbed dose was excreted by the lungs as carbon dioxide. The pulmonary excretion was biphasic with half-lives of 3.9 hours and 11.4 hours. The half-lives of radioactivity were 9.6 hours and 16.1 hours in the urine and feces, respectively.

In rats exposed to methyl bromide by inhalation, 47 to 50% of the absorbed dose was exhaled as CO_2 and 0.4 to 4% as methyl bromide (Medinsky *et al.*, 1985). Excretion in the exhaled air was biphasic with a half-life of 4.1 hours in the initial phase and 17 hours in the second phase. Approximately 20% of the absorbed radioactivity was excreted in the urine, while only about 1% was found in the feces. The urinary half-life for radioactivity was 9.9 hours.

In another inhalation study with rats, nearly 75% of the absorbed dose was excreted by 65 hours with exhalation as the primary route (Bond *et al.*, 1985). The excretion of $^{14}\text{CO}_2$ in the exhaled air was biphasic with the initial half-life at 3.9 hours, and a later half-life of 11.4 hours. The urinary half-life was 9.6 hours. The percentages of the absorbed dose excreted in the expired air as CO_2 , in the expired air as methyl bromide, in the urine, feces, and retained in the body were 47%, 1%, 23%, 2%, and 28%, respectively. A similar excretion pattern was reported by Jaskot *et al.* (1988).

In dogs, the excretion of radioactivity was 1% (total inhaled) in the urine, 0.04% in the feces, and 9.86% in the lungs when measured 0 to 21 hours after exposure to methyl bromide (Raabe, 1986). At 69 hours after exposure, 5.7% and 0.7% of the total amount inhaled were excreted in the urine and feces (Raabe, 1986). The estimated total clearance half-life was 41 hours.

In humans, the amount exhaled as $^{14}\text{CO}_2$ ranged from 0.2 to 1.0% of the dose for mouth breathing, and 0.2 to 0.4% of the dose for nose breathing exposure when measured at the end of 2 hours of exposure and after 0.5 hour for clearance (Raabe, 1988). Radioactivity levels excreted in the urine ranged from 0.08 to 0.24% for mouth breathing, and from non-detected to 0.32% for nose breathing. The determination of urinary clearance was complicated by the variability in the volume. The net body retention for both exposure routes was 51.1% with a clearance half-life of 72 hours based on the amounts in the exhaled air and in the urine at 0.5 hour after inhalation exposure.

In rats after oral exposure, the distribution (as % of an absorbed dose) was 32% as $^{14}\text{CO}_2$ and 4% as intact methyl bromide in exhaled air, 43% in the urine, and 14% in the carcass at 72 hours after exposure to methyl bromide (^{14}C , >98% pure, 250 $\mu\text{mol}/\text{kg}$ or 23.7 mg/kg) (Medinsky *et al.*, 1984). Only 2% of the dose was found in the feces. In cannulated rats, biliary excretion was a major pathway as 46% of the dose was found in the bile and only 12% and 7% in the exhaled CO_2 and urine, respectively, at 24 hours after dosing.

With intraperitoneal administration, the cumulative percentages of the doses in rats measured after 72 hours were: 45% as $^{14}\text{CO}_2$ and 20% as intact methyl bromide in exhaled air, 16% in the urine, 1% in the feces, and 17% in the carcass (Medinsky *et al.*, 1984).

III.B. ACUTE TOXICITY

Summary: Methyl bromide is a Toxicity Category I compound because of its acute inhalation toxicity (Federal Register, 1991)^c. Severe irritation to eyes, skin, and mucous membranes occur after acute exposure; therefore, acute oral, ocular and dermal studies are not required for registration. Neurotoxicity has been observed in humans and laboratory animals after inhalation exposure to methyl bromide with severity depending on the dose and duration of exposure. In humans exposed to high concentrations, neurological effects included ataxia, convulsion, and tremors. The nonlethal effects observed in laboratory animals included changes in brain catecholamines and tyrosine hydroxylase activity, tissue degeneration (nasal, brain, and adrenal glands), and neurotoxicity (ataxia and paralysis). Signs of oral toxicity in the dog included prostration, increased heart rates, lesions in multiple organs including the stomach and brain, hypoactivity, hypothermia, and death. Human dermal exposure resulted in skin lesions. The acute lethal toxicity of methyl bromide has been reviewed (Sayers *et al.*, 1929; Irish *et al.*, 1940; Alexeeff and Kilgore, 1983 and 1985; WHO, 1995). Studies considered for risk assessment are summarized in Table 4. NOELs and LOELs were determined only for those studies with adequate descriptions of the experimental protocol and results.

III.B.1. Inhalation - Rat

The first comprehensive study on the acute toxicity of methyl bromide in rats was conducted by Irish *et al.* (1940). Rats (strain not specified) were exposed to methyl bromide (99% pure, nominal concentrations of 0.42 to 50 mg/L or 100 to 13,000 ppm) continuously by inhalation, and the acute toxicity was determined. The lethal concentrations and exposure durations which resulted in 100% mortality rate are listed in Table 1. At 260 ppm for 20 hours, rats were excitable and jumped when stimulated. At concentrations < 2,600 ppm, they showed roughening of the fur, hunching of the back, drowsiness, heavy breathing, and occasionally lacrimation. After prolonged exposure, there were kidney damage, lung congestion and edema, and bronchopneumonia leading to death.

Fischer-344 rats were exposed to methyl bromide (99.9% pure; nominal concentrations of 0, 90, 175, 250, or 325 ppm) for 6 hours per day for 5 days (Hurtt *et al.*, 1987). At 325 ppm, severe tissue degeneration in the nasal cavity, brain (cerebellar and cerebral), liver, and adrenal glands as well as minor alteration in testicular histology (delayed spermiation) were observed. Three of seven rats in the 325 ppm group died after 3 exposures. In addition, diarrhea, hemoglobinuria, ataxia, tremors and convulsions occurred in these animals (325 ppm). Ataxia and diarrhea were also observed in the 250 ppm treated animals. No death or tremors were observed in animals at this and lower concentrations. The degeneration in the nasal cavity and other tissues was concentration dependent. There was partial or complete destruction of the olfactory epithelium in the 325 and 250 ppm groups. At 175 ppm, there was moderate degeneration of the cerebellum, adrenal glands, and nasal cavity. The NOEL was 90 ppm.

^c The criteria for Toxicity Category I include an inhalation LD₅₀ of less than or equal to 0.05 mg/L, and fatal if swallowed, inhaled, or absorbed through the skin.

Table 1. The acute lethal toxicity of technical methyl bromide.

Species	Gender	Concentration ppm	Dosage mg/kg	Reference ^a
<u>Oral LD₅₀</u>				
Rat	M		214	1
Rat	M		104-133	2
<u>Subcutaneous LD₅₀</u>				
Rat	M		135	3
<u>Inhalation LC₁₀₀</u>				
Rat (6 min)	- ^b	13000		4
Rat (24 min)	- ^b	5200		4
Rat (42 min)	- ^b	2600		4
Rat (4 hr)	M	900		5
Rat (6 hr)	- ^b	520		4
Rat (26 hr)	- ^b	220		4
Rabbit (30 min)	- ^b	13000		4
Rabbit (1.4 hr)	- ^b	5200		4
Rabbit (11 hr)	- ^b	520		4
Rabbit (24 hr)	- ^b	260		4
Rabbit (32 hr)	- ^b	220		4
Guinea Pig (15 min)	- ^b	50000		6
Guinea Pig (1.5 hr)	- ^b	6950		6
Guinea Pig (3 hr)	- ^b	2290		6
Guinea Pig (8 hr)	- ^b	490		6
<u>Inhalation LC₅₀</u>				
Rat (15 min)	- ^b	5480		7
Rat (30 min)	- ^b	2700		8
Rat (4 hr)	M	780		5
Rat (8 hr)	M	310		8
Mouse (1 hr)	M	1200		9
Guinea Pig (5 hr)	- ^b	310		6
<u>Lethality</u>				
Human (5.5-7.5 hrs) ^c	M	8160		10
Human-adult (2 hr) ^d	M	60000		11

^{a/} References: 1. Danse *et al.*, 1984; 2. Kiplinger, 1994; 3. Tanaka *et al.*, 1988; 4. Irish *et al.*, 1940; 5. Kato *et al.*, 1986; 6. Sayers *et al.*, 1929; 7. AmeriBrom, Inc. 1983; 8. Honma *et al.*, 1985; 9. Alexeeff and Kilgore, 1985; 10. Miller, 1943; 11. Wyers, 1945.

^{b/} Sex was not specified in the study.

^{c/} This man was found in convulsions in a refrigerator car fumigated with methyl bromide 5.5 to 7.5 hours earlier. He was provided with life support but died 80 hours after removal from the car.

^{d/} This worker used methyl bromide to extinguish a fire and then continued to work for 2 hours. He was hospitalized and died one hour later.

In a subsequent study, Fischer-344 rats were exposed to methyl bromide (99.9% pure; nominal concentrations of 0, 90, or 200 ppm) for 6 hours per day for 1-5 days to determine regeneration and recovery of olfactory function (Hurtt *et al.*, 1988). No clinical signs of toxicity or olfactory epithelial damage in the 90 ppm group were observed. At 200 ppm, there was a transient significant decrease (9%) in body weight at the end of the fifth exposure. The body weights of the treated group returned to control values by day 47 after exposure. Extensive destruction of the olfactory epithelium of the dorsal meatus, nasal septum and lateral walls, and the complex ethmoid turbinates was evident after a single 6-hour exposure to 200 ppm. The olfactory degeneration was characterized by epithelial disruption, fragmentation, and exfoliation. The basal cell layer remained intact. Regeneration of the epithelium with the replacement by a squamous epithelium was evident by day 3 of exposure and was essentially complete by 10 weeks after exposure. However, some minor defects were not repaired such as: adhesions between the turbinates and adjacent structures, thinning of the olfactory epithelium due to a paucity of sensory cells, and respiratory metaplasia (conversion of the olfactory epithelium to a ciliated respiratory type). Olfactory function, as determined by the ability of food-deprived rats to find buried food pellets, was affected only in rats treated at 200 ppm. The impairment of the olfactory function was temporary with recovery by 4 to 6 days after exposure but before complete regeneration of the olfactory epithelium. The NOEL was 90 ppm based on olfactory damage at 200 ppm.

In a similar experiment, rats (strain not specified) were exposed to methyl bromide (purity not specified, a nominal concentration of 200 ppm) for only 4 hours per day, 4 days per weeks for 2 weeks (Hastings, 1990). After a single 4-hour exposure, the olfactory epithelium was extensively damaged, and olfactory function was impaired. After the end of exposure, no overt signs of toxicity were observed. The epithelium began to repair immediately after exposure but required more than 30 days for the restoration to a near normal appearance. Consistent with the findings of Hurtt *et al.* (1988), olfactory function returned to normal before complete epithelial regeneration.

CD rats (15/sex/group) were exposed to methyl bromide (>99% pure; nominal concentration of 0, 30, 100, or 350 ppm) for a single 6-hour inhalation exposure (Driscoll and Hurley, 1993). Testing was done at pre-exposure, within 3 hour post-exposure, 7 days post-exposure, and 14 days post-exposure using an automated assessment of motor activity and a Functional Observation Battery. Rats were sacrificed 16 to 19 days post-exposure. There were no effects on survival, body weight, and brain weight. No histological lesions were noted in the nervous system or the nasal tissues of the 350 ppm rats. Neurobehavioral effects were only seen in the 350 ppm group tested 3 hours post-exposure. Findings included: decreased arousal (both sexes); increased incidences of drooping or half-shut eyelids (both sexes); increased urination (females only); decreased rearing (both sexes); decreased tail pinch response (males only); increased incidences of piloerection (both sexes); decreased rectal temperature (both sexes); abnormal air righting reflex (females only); and decreased motor activity (both sexes). The NOAEL was 100 ppm based on neurobehavioral effects at 350 ppm.

The biochemical effects of methyl bromide were studied by Hurtt and Working (1988), Jaskot *et al.* (1988), and Davenport *et al.* (1992). Adult male Fischer 344 rats were exposed to methyl bromide (99.9% pure; 0, or a nominal concentration of 200 ppm) by inhalation for 6 hours per day for 5 days (Hurtt and Working, 1988). Rats were sacrificed on days 1 (first day of exposure), 3, 5, 6, 8, 10, 17, 24, 38, 52, and 73. Methyl bromide did not affect testis weight,

testicular and epididymal histology, daily sperm production, cauda epididymal sperm count, sperm morphology, sperm motility, and linear sperm velocity, or cause any observable toxicity. At day 5, the methyl bromide treated group weighed approximately 10% less than the control group and continued to weigh less until day 52. The nonprotein sulfhydryl levels of the testis and liver were significantly ($p \# 0.05$) decreased after 1 and 3 days of treatment. The depression was transient as the levels returned to control values by day 8 (3 days after treatment). There was a transient decrease in the testosterone level during the 5-day exposure, as well as the day after exposure. The testosterone level was back to the control level by day 8.

CD rats were exposed to methyl bromide (99.5% pure, 0 or a nominal concentration of 30 ppm) for 5 or 10 days (Jaskot *et al.*, 1988). After either 5 or 10 days of exposure, there were significant ($p \# 0.05$) decreases in the enzyme activities of glutathione (GSH) reductase and GSH transferase in the liver and increases of GSH transferase and glucose-6-phosphate dehydrogenase in the lung. The decreases were no more than 17% of the respective control values. In addition, serum chemistry showed significant ($p \# 0.05$) decreases in the levels of blood urea nitrogen (BUN), uric acid, cholesterol, and erythrocyte cholinesterase activity, as well as an increase in leucine aminopeptidase activity.

Davenport *et al.* (1992) proposed that methyl bromide-induced neurotoxicity was due to an effect on GSH and glutathione-S-transferase in the brain. Fischer-344 rats were exposed to methyl bromide (99.9% pure; 150 ppm) for 6 hours per day for 5 days. The concentration was chosen because it did not induce brain lesions or signs of toxicity. The inhibition of GSH transferase ranged from 45 to 56% of control values and the depletion of GSH ranged from 51 to 86% for the different regions (frontal cortex, caudate nucleus, hippocampus, brain stem, and cerebellum). Pretreatment and post-treatment of rats with BW 755C (3-amino-1-[m-(trifluoromethyl)phenyl]-2-pyrazoline), an inhibitor of monohalomethane toxicity, protected against GSH transferase inhibition in all brain regions and gave partial protection against GSH depletion. Monoamines (dopamine and serotonin in the frontal cortex, caudate nucleus, and hippocampus) and amino acids were not affected by the treatment.

Another proposed mechanism for the neurotoxicity of methyl bromide was the alteration of catecholamine levels in the brain (Honma *et al.*, 1987). Male Sprague-Dawley rats (5/group) were exposed to methyl bromide (purity not specified; nominal concentrations 0, 31, 63, 100, 125, or 250 ppm) for 8 hours. Catecholamine levels (dopamine, DA; norepinephrine, NE; homovanillic acid, HVA; 3-methoxy-4-hydroxyphenylglycol, MHPG; serotonin, and 5-hydroxyindoleacetic acid, 5HIAA) were determined in the striatum, hypothalamus, frontal cortex, midbrain, and medulla oblongata after exposure (Table 2). Dopamine levels were decreased in all regions with a lowest-effect level (LEL) of 100 ppm for the striatum. Norepinephrine was decreased (87% of control, $p < 0.05$) with a LEL of 31 ppm in the hypothalamus. However, the decrease (84% of control) at the next dose of 63 ppm was not statistically significant. Homovanillic acid and MHPG were increased with a LEL of 63 ppm in the striatum, hypothalamus, and midbrain (MHPG only). Serotonin and 5HIAA were not significantly affected in any brain segment. Time course studies showed that the maximal effect was detected immediately or 2 hours after exposure for most catecholamines, and the levels returned to control levels 24 hours after exposure was stopped. There were inconsistencies in this study. First, decreased dopamine was measured in the striatum of rats exposed for 8 hours to 100 or 125 ppm methyl bromide whereas exposure at these same levels for a much longer period, 24 hours, did not affect dopamine content in Honma *et al.* (1982) (**III.C. SUBCHRONIC TOXICITY**).

Second, if tyrosine hydroxylase was inhibited as proposed in Honma *et al.* (1991), one would expect that the metabolites such as HVA, “downstream” from tyrosine hydroxylase would be decreased. However, the HVA level was increased in this study.

The inhibition of tyrosine hydroxylase has been proposed as the mechanism for the reduction in dopamine levels. Male Sprague-Dawley rats (3-5/group) were exposed to methyl bromide (purity not specified; nominal concentrations of 0, 16, 31, 63, 125, or 250 ppm) for 8 hours (Honma *et al.*, 1991). Brain tyrosine hydroxylase activity (TH) in different brain segments was determined by *in vitro* (all doses) and *in vivo* (31 ppm) methods at 0, 1, 2, and 24 hours after exposure. Both assays indicated dose-responses for decreases in DOPA (3,4-dihydroxy phenylalanine) production. The segment with the LEL in the “*in vitro*” assay was the hypothalamus; the LEL was 16 ppm, the lowest dose tested (Table 2). The segments with the LEL in the “*in vivo*” assay were the striatum and hypothalamus; the LEL was 63 ppm, with a possible incipient effect at 31 ppm. The maximal inhibition of TH in both assays was seen with the rats sacrificed immediately after the 8 hour exposure period; significant recovery took place within two hours post-exposure and was complete by 24 hours post-exposure. The authors interpreted their findings as evidence that methyl bromide directly caused changes in the enzyme structure, presumably by methylation. However, DPR had significant questions about the findings and its relationship to other studies (additional discussion in **IV.A.1.b. Brain Monoamines and Enzyme Activity**).

The possible relationship between methyl bromide and dopamine was further investigated by testing whether rats exposed to methyl bromide were more sensitive (responsive) to the dopamine agonist apomorphine, which causes hyperactivity in rats (Honma *et al.*, 1994). Increased sensitivity to a dopamine agonist was expected if methyl bromide had damaged presynaptic neurons that use dopamine as the neurotransmitter. Also, the experiment tested whether methyl bromide affected the hypoactivity induced by the norepinephrine agonist clonidine. In the first of two assays, stereotypic oral behavior caused by an intraperitoneal injection of apomorphine was determined in male Sprague-Dawley rats (5/group) 7 days before exposure to methyl bromide and on days 1, 4, 7, 14, and 28 post-exposure. There were two types of inhalation exposure: 8 hours to 0, 25, 50, 100, or 200 ppm; and 8 hours per day for 7 consecutive days to 0, 5, 10, 25, or 50 ppm. There were no consistent dose responses for the supposed methyl bromide-induced increases in stereotypic behavior. The second assay involved measuring locomotor activity (automated-counting apparatus) after an intraperitoneal injection of apomorphine or clonidine. This assay (2 rats/dose) was conducted 7 days after exposure to 0, 10 or 50 ppm methyl bromide (8 hours/day for one day or for 7 consecutive days). Testing also was done the day before exposure to methyl bromide; in these instances, neither apomorphine nor clonidine were administered before the locomotor activity was recorded. The effect of methyl bromide on locomotor activity was uncertain as the data were only for 2 rats per dose and the variability in the mean value was not indicated.

Table 2. Alteration of catecholamine levels and tyrosine hydroxylase activity in the rat brain after acute inhalation exposure to methyl bromide^a.

Exposure (ppm)	16	31	63	100	125	250	63	125	250
Brain Region	% control (0 hour post-exposure)						% control (2 hours post-exposure)		
Striatum									
DA		92	94	76**	81*	70**			
HVA		106	144**	145**	132	137*			
NE		89	86	66*	89	78			
MHPG		107	127*	129*	139**	142**			
TH <i>in vitro</i>	98	87	72*		61**	60**	104	85*	77*
TH <i>in vivo</i>		90	62**		40**	26**	80	69*	81*
Hypothalamus									
DA		89	89	81**	82*	73**			
HVA		101	133*	137**	114	127			
NE		87*	84	79**	73**	70**			
MHPG		107	130*	126*	126	134*			
TH <i>in vitro</i>	82*	65**	65**		59**	57**	92	85	75*
TH <i>in vivo</i>		90	72*		50**	47**	104	96	79
Frontal cortex									
DA		87	92	90	91	75*			
HVA		102	121	144**	125*	125*			
NE		86	78	68**	72*	74*			
MHPG		101	127	110	115	146**			
TH <i>in vitro</i>	100	92	87*		84*	80**	105	89	80*
TH <i>in vivo</i>		103	90		62**	39**	106	101	97
Midbrain									
DA		95	88	86	87	77			
HVA		114	133	161**	148**	150**			
NE		96	80	73*	79**	73**			
MHPG		102	121*	132**	151*	140**			
TH <i>in vitro</i>	103	97	92		87	64**	106	100	97
TH <i>in vivo</i>		96	81*		68**	53**	104	102	99
Medulla									
DA		92	90	88	89	76*			
HVA		104	120	140**	149**	137*			
NE		94	91	86	83*	86			
MHPG		100	114	108	112	139**			
TH <i>in vitro</i>	97	89	91		85	72**	106	105	89
TH <i>in vivo</i>		111	92		69*	48**	104	99	104

* ** Significance level at $p < 0.05$ (*) or $p < 0.01$ (**) based on statistics as reported.

^{a/} Data from Honma *et al.* (1987) for dopamine (DA), norepinephrine (NE), homovanillic acid (HVA), and 3-methoxy-4-hydroxyphenylglycol (MHPG) levels; and Honma *et al.* (1991) for tyrosine hydroxylase activity (TH). Measurements were made immediately, 2 hours, or 24 hours after 8 hours of exposure. Some data were not shown (see details of the studies in the text).

III.B.2. Oral- Rat

Albino rats (5/sex/group) were given either methyl bromide (99.5% pure) in corn oil or in microcapsules (Kiplinger, 1994). In the liquid methyl bromide testing, methyl bromide was given once by gavage at 50, 100, and 150 mg/kg in initial testing and at 0, 80, 120 and 160 mg/kg in retesting. The initial dose levels were chosen on the basis of range-finding testing which also was discussed in the report. Rats were fasted for 18-20 hours prior to dosing and feed was made available 3-4 hours after dosing. Rats were observed for mortality and clinical signs at approximately 1, 3, and 4 hours after dosing (post-dosing day 0) and once in the morning and once in the afternoon on post-dosing days 1 through 14 (day of scheduled sacrifice). All rats in the initial testing and retesting were necropsied. In the retesting, microscopic examination of the stomach, duodenum, jejunum and ileum also was performed. With one exception, rats that died did so on or before post-dosing day 2. At necropsy, the main organ affected was the stomach. The findings were consistent with severe irritation of the lumen surface. The mortality data indicated a slight sex difference; for example, LD50 values for males and females in the initial testing were, respectively, 139 mg/kg (125-155 mg/kg as 95% confidence interval) and 107 mg/kg (97-119 as 95% confidence interval). The lowest LD50 value was 86 mg/kg (77-96 mg/kg as 95% confidence interval); this was seen in the females in the retesting. The testing of the microencapsulated methyl bromide can not be evaluated pending clarification of the following: (1) whether the microcapsules dissolved before dosing; (2) whether the procedure for the methyl bromide content analyses was appropriate; and (3) whether the microencapsulated material was comparable to the material used in the two-year microencapsulated methyl bromide feed study (Mertens, 1997; described in **III.D.2. Dietary - Rat**). This study was considered supplemental.

III.B.3. Inhalation - Rabbit

Rabbits were exposed to methyl bromide (99% pure; nominal concentrations of 0.42 mg/L to 50 mg/L or 100-13,000 ppm) by inhalation (Irish *et al.*, 1940). Paralysis was observed in some rabbits exposed at 1 mg/L (260 ppm) for 20 hours. There was an indication of lung irritation at higher concentrations, though less pronounced than that observed in the rat. The concentrations and exposure times which resulted in 100% mortality are listed in Table 1.

III.B.4. Inhalation - Mouse

Swiss-Webster mice were exposed to methyl bromide (>99.5% pure) from 0.87 to 5.93 mg/L (223-1518 ppm) for 1 hour by nose only inhalation (Alexeeff and Kilgore, 1985). At all concentrations tested, the 24-hour weight gain of the treated mice was significantly ($p \neq 0.05$) decreased by 4-16% when compared to control group. Transient hyperactivity was observed in the 2.72 and 3.82 mg/L (696 and 978 ppm) groups. Abnormal gait, passivity, and lack of grooming were evident at 3.5 mg/L (896 ppm) and higher with the earliest onset at 3 hours for 4.7 to 5.93 mg/L. Additional clinical signs included: increased depth of respiration, decreased respiration rate, tremors, fasciculation, loss of righting reflex, splayed limbs, tonic seizures, and muscular rigidity. Death occurred at 3.82 mg/L and higher levels. There was also rectal hemorrhaging with diarrhea in those animals treated at 5.77 and 5.93 mg/L, occurring within 6 hours after treatment. One week after treatment, damage was observed in the kidneys

(\$3.50 mg/L), liver (\$3.82 mg/L), brain (\$3.82 mg/L), and colon (\$3.82 mg/L). Glutathione levels in livers of the 4.7 and 5.93 mg/L groups were significantly ($p \neq 0.05$) lower than the control group. The NOEL was 2.72 mg/L (696 ppm) based on mortality and other effects at 3.82 mg/L.

III.B.5. Inhalation - Dog

Dogs were exposed to methyl bromide for one day, two days, and four days in range-finding studies to select doses for a one-year exposure study (Newton, 1994a). The average measured methyl bromide concentrations ranged from 55 to 394 ppm. The dogs were observed every 15 minutes for the first hour and hourly thereafter. The results for short-term exposures are summarized in Table 3.

In the one-day experiment, dogs (1/concentration, except 2 dogs used for 314 ppm) were exposed for 7 hours to methyl bromide (233, 314, 345, 350, and 394 ppm). All dogs exhibited neurotoxicity (including tremors, hunched appearance, and labored breathing) by the seventh hour, and the earliest onset was 3 hours in the 345, 350, and 394 ppm groups. The exposure of the 394 ppm dog ended at the 6th hour because of clinical signs (mucoïd nasal discharge and labored breathing). One day after exposure, the 345 and 350 ppm dogs showed moderate white nasal discharge, while the 394 ppm dog was lethargic with excessive nasal discharge, salivation, and panting. At 314 ppm, one dog showed decreased activity after 4 hours of exposure. All the dogs were observed for 3 or more days and appeared normal before they were used for the two-day study. The NOAEL for the one-day exposure was < 233 ppm for neurotoxicity.

In the two-day exposure study, dogs (3/group) were exposed to 268 ppm and 283 ppm of methyl bromide (Newton, 1994a). These dogs had been used in the one-day experiment: 268 ppm group were previously exposed to either 233, 314, or 394 ppm; while the 283 ppm group were exposed to either 314, 350, or 345 ppm. At the start of this study, all dogs appeared clinically normal. On the first day, one of the 283 ppm dogs showed excessive salivation, labored breathing, and emesis at the 6th hour of exposure. By the 7th hour, the dogs in this group showed labored breathing (3/3), excessive salivation (2/3), and emesis (2/3). By the second day, the 283 ppm group exposure was terminated because of the following observations: severe neurotoxicity (delirium, thrashing and vocalization, tremors, traumatizing behavior, depression, ataxia, irregular gait) and abnormal respiratory sounds. The 268 ppm group dogs appeared normal on the first day. One of the dogs showed labored breathing at the 3rd to 7th hour on the second day. All dogs showed decreased activity at the 7th hour of the second day. Also, increased blood urea nitrogen and serum aspartate aminotransferase were detected in both exposure groups. The NOAEL for the two-day exposure was < 268 ppm for neurotoxicity.

The four-day exposure study used dogs that had not been exposed to methyl bromide previously (Newton, 1994a). Beagle dogs (1/sex/group) were exposed to 55 ppm and 156 ppm by inhalation for four days (7 hours/day). In the 156 ppm group, one dog showed lacrimation at the fifth hour of exposure. Both dogs exposed to 156 ppm showed decreased activity, lacrimation and labored breathing during the third and fourth day of exposure. Irregular gait was observed in both dogs on the fourth day post-exposure. No abnormal signs were observed during or after exposure for the 55 ppm dogs. The NOAEL for neurotoxicity observed after 3 days of exposure was 55 ppm.

Table 3. The neurotoxicity of methyl bromide in dogs after acute exposure^a.

Concentration mean \pm standard deviation (ppm)	Duration of exposure ^a	First signs of neurotoxicity and Incidence ^b	Clinical signs with additional exposure
394 \pm 20	3 hours	hunched appearance and tremors (1/1)	hunched appearance and tremors, mucoid nasal discharge, labored breathing
350 \pm 13	3 hours	labored breathing (1/1)	labored breathing, decreased activity, hunched appearance, tremors, excessive salivation and swallowing response
345 \pm 8	3 hours	tremors (1/1)	tremors, labored breathing, hunched appearance, excessive salivation, and gasping
314 \pm 6	4 hours	decreased activity (1/2)	tremors, hunched appearance, tremors, and restlessness (2/2)
283 \pm 13	6 hours	salivation, labored breathing, emesis (1/3)	excessive salivation (2/3), labored breathing (3/3), and emesis (2/3)
268 \pm 19	7 hours	no effects	day 2: labored breathing (1/3) and decreased activity (3/3)
233 \pm 21	5 hours	trembling (1/1)	panting, rapid eye blinks, and tremors
158 \pm 7 ^c	7 hours ^c	decreased activity (8/8)	day 3: decreased activity; day 6: severe neurotoxicity; brain, and adrenal lesions, olfactory degeneration (8/8)
156 \pm 15	5 hours	lacrimation (1/2)	day 3 and 4: lacrimation and labored breathing (2/2), prostrate (1/2), and decreased activity (2/2); day 4 post-exposure: irregular gait (2/2)
103 \pm 9 ^c	8 days	no effects	day 9: start of decreased activity (3/8) and emesis (1/8); day 10: tremor (1/8); week 5: cerebellar lesions (1/8) at sacrifice
53 \pm 4 ^c	13 days	no effects	day 14: decreased activity (2/8)
55	4 days	no effects	(experiment terminated after 4 days)

^{a/} Data from Newton (1994 a and b). Hours of exposure for onset of neurotoxicity.

^{b/} Incidences as number of dogs affected/ total are shown in parentheses.

^{c/} Data (Newton, 1994b) is under **III.C. SUBCHRONIC TOXICITY**. The first exposure was on a Friday with no effects reported. However, decreased activity was observed during the second exposure, on the following Monday.

III.B.6. Oral - Dog

Beagle dogs (1/sex/group) were given a single oral dose of methyl bromide (100% pure; 1, 3, 5, 50, or 500 mg/kg) in capsules in a corn oil solution (Naas, 1990). There was no control group. Animals treated at 500 mg/kg showed prostration, increased heart rates, and were dead within one day. Necropsy showed damage in the stomach (reddened mucosa), kidneys (reddened cortico-medullary junction), adrenal glands (dark red, bilateral), and brain (hydrocephaly). At 50 and 500 mg/kg, hypoactivity, hypothermia, and no pain reflex were observed. Some animals treated at 3 mg/kg and higher doses vomited, in some cases the vomitus contained reddish material. No vomiting was observed during one week post-dosing observation period. The NOEL was 5 mg/kg based on toxicity and mortality observed at the higher dosages. This study was considered supplemental information by DPR.

III.B.7. Inhalation - Guinea Pig

Guinea pigs were exposed to methyl bromide (purity not specified, nominal concentrations of 0.01 to 5%, with 1% =10,000 ppm) for 5 to 810 minutes (Sayers *et al.*, 1929). No tissue damage or death was observed when the guinea pigs were exposed to 0.06% for 90 minutes, or 0.03% for 270 minutes. The mortality was 100% at 5% of methyl bromide for 15 minutes, 1.3% for 68 minutes, 0.7% for 90 minutes, or 0.23% for 170 minutes. Death occurred within 21 hours after exposure at 2.2% of methyl bromide for 35 minutes, 0.23% for 90 minutes, and 0.05% for 480 minutes. Before death, lacrimation, difficulty in breathing, and weakness were observed. Necropsy showed congestion and tissue degeneration of multiple organs (including lungs, liver, heart, kidneys, and brain).

Guinea pigs (16/group) were exposed to methyl bromide (purity not specified; nominal concentration of 100 or 220 ppm for 7.5 to 8 hours per day for 1 to 3 days (Irish *et al.*, 1940). Guinea pigs at 220 ppm showed respiratory difficulty before death and lung damage. No effect was observed at 100 ppm.

III.B.8. Inhalation - Human

The effects of methyl bromide in humans after acute exposure is well-documented. A summary of the clinical signs and symptoms is provided in this section. Those studies which reported actual exposure levels are described in detail in **III.H. NEUROTOXICITY**. The signs and symptoms after methyl bromide exposure are dependent on concentration and exposure duration (von Oettingen, 1946; Rathus and Landy, 1961; Greenberg, 1971; Grant, 1974; Anger *et al.*, 1986; Gehring *et al.*, 1991; Uncini *et al.*, 1990; De Haro *et al.*, 1997). Acute exposures to lethal concentrations result in early symptoms of malaise, headache, visual disturbances, nausea, and vomiting. After the early symptoms, delirium, disorientation, and excitability may also occur. There is a delayed onset of symptoms that include numbness, ataxia, tremor, myoclonus, exaggerated (or absent) deep reflexes, positive Romberg's signs, paroxysmal abnormalities of the EEG, agitation, change of personality, coma, as well as clonic and tonic convulsions. Death usually occurs within 48 hours of exposure, due to pulmonary edema leading to respiratory failure or cardiovascular collapse. Postmortem examination of the brain showed generalized swelling and lesions, including neuronal loss (Squier *et al.*, 1992).

Nonlethal exposures to methyl bromide result in neurological effects that include the early symptoms seen in lethal exposures as well as confusion, muscular weakness, tremors, convulsions, euphoria, delirium, and psychoses. Some symptoms may persist after exposure depending on the severity of toxicity. Ataxia and myoclonus continued to be experienced by a worker one year after exposure (Rathus and Landy, 1961). In severe chronic poisoning, endogenous chloride is replaced by bromide, the ionic form of bromine, from the biotransformation of methyl bromide in the body (Blumberg and Neli, 1967).

III.B.9. Dermal - Human

Six workers (1 woman and 5 men) were exposed to methyl bromide (estimated 35 g/m³, or 9,000 ppm) dermally during the fumigation of a castle (Hezemans-Boer *et al.*, 1988; Zwaveling *et al.*, 1987). They wore tight-fitting face masks, overalls over their daily clothing, work shoes, and polyvinylchloride gloves. Shortly after exposure, the primary complaint was burning sensation under the armpits and in the groin. No neurotoxicity was observed. Redness of the skin was noted. Approximately 8 hours after exposure, all workers developed skin lesions which consisted of sharply demarcated erythema with multiple vesicles and large bullae in areas of moisture (from perspiration) such as the axillae, groin, and submammary areas. Histopathological examination showed necrosis of keratinocytes, edema of the papillary dermis, subepidermal blistering, and preferential infiltration by neutrophils. Six days after exposure, there were signs of regeneration of the epidermis, but there was still evidence of cell necrosis and presence of neutrophils. One month after exposure, the skin lesions had disappeared with no significant scarring.

III.B.10. Additional Acute Studies

Acute effects were also observed in studies described (in detail) in **III.C. SUBCHRONIC TOXICITY** and **III.G. DEVELOPMENTAL TOXICITY** and are included in Table 4.

Table 4. The no-observed-effect levels (NOELs) and lowest-observed-effect levels (LOELs) for acute effects of methyl bromide from selected studies^a.

Route/ Species	Exposure Duration ^b	Study NOEL/LOEL ppm	Human Equivalent ^c NOEL/LOEL ppm	Effects	Ref. ^d
<u>INHALATION</u>					
Rat	6h/dx5d	90/175	47/91	tissue degeneration (brain, adrenals, nasal cavity) neurotoxicity at higher doses (250 ppm)	1
Rat	6 h	90/200	47/104	olfactory epithelium degeneration, 9 olfactory function	2
Rat	8h	31/63 <16/16	22/44 <11/11	altered brain catecholamines and tyrosine hydroxylase activity altered tyrosine hydroxylase activity (hypothalamus)	3
Rat	6 h	100/350	52/183	changes in neurobehavioral battery	4
Mouse	1 h	696/896	113/146	abnormal gait, passivity, no grooming	5
Dog	7h 7h/dx3d	<233/233 ^e 55/156	<58/58 14/39	neurotoxicity 9 activity, labored breathing, prostrate	6
Dog^f	7h/dx1d	103/158	25/39	9 activity on 2nd exposure day; brain lesions after 6 exposures (had previous exposure of 11 ppm for 24 days).	7
Guinea Pig	8h/dx1-3d	100/220	31/61	respiratory difficulty, death	8
<u>INHALATION-DEVELOPMENTAL TOXICITY STUDIES^g</u>					
Rat	7h/d, gd1-19	20/ 70	22/75	progeny-delayed skull ossification	9*
Rabbit	6h/d, gd7-19	40/ 80	21/42	progeny-fused sternebrae, gall bladder agenesis and other effects	10*
<u>ORAL</u>					
Dog	1 dose (capsule)		na	hypoactivity, hypothermia, no pain reflex	11
<u>DERMAL</u>					
Human	few hours	-/9000 in air	na	skin lesions (erythema, vesicles)	12

^{a/} Bolded studies are those selected to derive the critical NOELs for risk characterization.

^{b/} Duration: min=minutes, h=hours, d=days, w=weeks, and gd=gestation days.

^{c/} Human equivalent NOEL/LOELs were calculated by adjusting for respiration rate differences and duration of exposure (Appendix G). The equivalent levels were those for children only (except references 9 and 10). The adult equivalent levels would be 2-fold higher since the adult respiration rate is lower (0.26 m³/kg/day compared to the child's 0.46 m³/kg/day). For references 9 and 10, the equivalent levels were those for adults since the effects were observed in pregnant animals.

^{d/} * after reference number indicates the study was acceptable according to FIFRA guidelines. References: 1. Hurtt *et al.*, 1987; 2. Hurtt *et al.*, 1988; 3. Honma *et al.*, 1987 and 1991; 4. Driscoll and Hurley, 1993; 5. Alexeeff and Kilgore, 1985; 6. Newton, 1994a; 7. Newton, 1994b; 8. Irish *et al.*, 1940; 9. Sikov *et al.*, 1981; 10. Breslin *et al.*, 1990b; 11. Naas, 1990; 12. Hezemans-Boer *et al.*, 1988.

^{e/} The lowest dose tested.

^{f/} Study described in **III.C. SUBCHRONIC TOXICITY**.

^{g/} Studies described in **III.G. DEVELOPMENTAL TOXICITY**.

III.C. SUBCHRONIC TOXICITY

Summary: Subchronic inhalation exposure of laboratory animals to methyl bromide resulted in altered brain catecholamine levels, decreased brain tyrosine hydroxylase activity, neurotoxicity, tissue degeneration (brain, nasal cavity, heart, testes, adrenal glands, thymus, spleen, and kidneys), and death. Based on overt signs of neurotoxicity, the dog, rabbit, and monkey were more sensitive to methyl bromide than other species (rat, mouse, and guinea pig). The primary finding in the rat was hyperplasia of the forestomach after oral exposure by gavage. A decrease in body weight gain and food consumption was observed in rats given microencapsulated methyl bromide mixed in the feed. A summary of selected subchronic studies is presented in Table 6.

III.C.1. Inhalation - Rat

Rats (strain not specified, 10-12/group) were exposed to methyl bromide (99% pure, nominal concentrations of 17 to 220 ppm) by inhalation for 7.5 to 8 hours per day, 5 days per week for 6 months or until toxicity was observed (Irish *et al.*, 1940). Methyl bromide at 220 ppm was lethal; all the rats died after 3 to 4 days. At 100 ppm after 6 or more exposures, the rats were in poor general appearance and some developed convulsions. Lung lesions were observed in 66% of the rats in the 100 ppm group. Except for the one death at 66 ppm (time of death not specified), no other effect was observed at this or lower doses.

Sprague-Dawley rats were exposed to methyl bromide (99.9% pure; nominal concentration of 65 ppm for 7.5 hours per day, 4 days per week for a total of 100 hours or 55 ppm for 6 hours per day, 5 days per week for 36 weeks (Anger *et al.*, 1981). There were no effects on sciatic and ulnar nerve conduction velocities, open field activity, or coordination.

Male Sprague-Dawley rats (5-6/group) were exposed to methyl bromide (99.94% pure; nominal concentrations 0, 1, 5, or 10 ppm) for 3 weeks (daily exposure duration not specified) (Honma *et al.*, 1982). Data were presented in graphs. There was no treatment-related change in the levels of acetylcholine in the whole brain or striatum, or cyclic AMP in the striatum. A marked decrease in cyclic GMP in the cerebellum at 5 and 10 ppm was not noted as statistically significant. The other effect at 5 ppm was an increase in dopamine in the striatum (~125% of control, $p < 0.05$). However at the next dose (10 ppm), the increase was only ~111% of control and was not statistically significant. Other monoamines were affected at 10 ppm. There was a decrease of norepinephrine in the hypothalamus (~35% of control, $p < 0.05$), and in the cortex and hippocampus (~30% of control, $p < 0.05$). Serotonin was decreased in the cortex and hippocampus (~70% of control, $p < 0.05$). The NOEL was 5 ppm for decreased monoamines in the brain at 10 ppm.

Sprague-Dawley rats were exposed to methyl bromide (purity not specified) for 4 hours per day, 5 days per week at nominal concentrations of 150 ppm for 11 weeks (Kato *et al.*, 1986). No clinical signs were observed. However, there was an increase in spleen weight (113% of control). Lesions in the heart were noted as small focal necrosis of heart muscle and fibrous replacement of heart muscle. In another experiment, rats were exposed to 200, 300, or 400 ppm for 6 weeks. Paralysis of the hind limb and death were observed in both the 300 and 400 ppm groups. The first death occurred after 3 weeks of exposure at 400 ppm. Voluntary movement was inhibited in all animals of the 400 ppm group by 2 weeks of exposure. Organ weights (brain, thymus, heart, liver and spleen) were significantly ($p \neq 0.05$) lower in one or both

dose groups. Histological examination of the brain of the 400 ppm group showed spongy nerve tissue and glial cell proliferation. Atrophy of the testes, sometimes accompanied by the appearance of giant cells in the seminal tubules, was observed in the 400 ppm group. Multiple and small focal necrosis or fibrosis were observed throughout the left and right ventricles and papillary muscle of the heart in all treated groups. A dose-related increase in bromide concentration was detected only in the liver.

The brain and heart were also target organs in Fischer-344 rats exposed to methyl bromide (99.8% pure, nominal concentration of 160 ppm) for 6 hours per day, 5 days per week for 6 weeks (Eustis *et al.*, 1988). Because there was more than 50% mortality, the male rats were sacrificed after 14 exposures. At the end of the 30 exposures, the survival of the female rats was 50%. There were no treatment related findings in the parameters measured in the clinical chemistry or urinalysis. The treated rats showed curling and crossing of hindlimbs, forelimb twitching, and tremors. There were significant ($p \# 0.05$) decreases in the body weight (68% of control), and the weights of lungs, heart, spleen, right kidney, brain, liver, and right testes of the males. For the females, significant ($p \# 0.05$) decreases in body weight (82% of control), and the weights of lungs, right kidney, brain, and liver were observed. Pathological lesions were found in the brain (necrosis and loss of neurons in the cerebral cortex, hippocampus, and thalamus), testes (degeneration and atrophy of the seminiferous tubules), nasal olfactory epithelium (necrosis and degeneration), heart (cardiomyopathy), adrenal cortex (cytoplasmic vacuolation), liver (hepatocellular necrosis), thymus (atrophy), and spleen (lymphoid depletion).

In a longer-term study, Fischer-344 rats were exposed to methyl bromide (99.8% pure, nominal concentrations of 0, 30, 60, or 120 ppm) for 6 hours per day, 5 days per week for 13 weeks (NTP, 1992; Eustis, 1992). No mortality was observed. The body weights were decreased as early as week 6 of exposure. By 13 weeks, the body weights of the 60 and 120 ppm males were 90% and 84% of control, respectively. The brain weights of the 120 ppm groups were significantly ($p \# 0.01$) decreased to 92% (males) and 93% (females) of control values. There were occasional neurobehavioral effects (such as decreased startle response amplitude, increased startle response latency, and decreased grip strength) in the 120 ppm groups which were significantly ($p \# 0.05$) different from control values; however, they were not related to exposure duration. Significant ($p \# 0.05$) decreases in the mean hematocrit, hemoglobin, and erythrocyte counts were detected in the 120 ppm females only. A significant decrease in the erythrocyte count also was noted in the 60 ppm females. An increase in the incidences of olfactory epithelial dysplasia and cysts (irregularity in mucosal thickness and focal cavity spaces) was found in both sexes of the 120 ppm groups. The NOEL was 60 ppm for neurotoxicity and olfactory epithelial dysplasia at 120 ppm.

CD rats (15/sex/group) were exposed to methyl bromide (>99% pure; nominal concentrations of 0, 30, 70, or 140 ppm) by inhalation for 6 hours per day, 5 days per week for 90 days (Norris *et al.*, 1993 a and b). Neurobehavioral testing was done pre-exposure and at the end of study weeks 4, 8 and 13. Testing included automated assessments of motor activity and a Functional Observation Battery (FOB). Two 140 ppm males died on test (days 12 and 27); the latter had convulsions and tremors before dying. One other 140 ppm male that survived until the

end of the study also exhibited clonic convulsions and tremors. The 140 ppm groups (both sexes) weighed significantly less than the controls, starting at study week 4 while the 70 ppm female group exhibited a body weight reduction, starting about study week 9. The mean body weights were 87% of control values on week 13 for 140 ppm groups, and 92% for the 70 ppm females.

FOB testing identified effects only in the 140 ppm groups; some effects were evident as early as study week 4. Findings included: ataxia (5 females, 1 male); decreased arousal (females only); decreased rearing activity (females only); increased hind leg splay (males only); and (possibly) abnormal air righting (males only). Motor activity testing identified effects only in the 70 ppm and 140 ppm female groups. Findings included decreased total motor activity and decreased rearing activity; both were evident as incipient effects in study week 8. Female groups exposed to methyl bromide exhibited a dose response for reduced absolute brain weight. The reductions were statistically significant ($p \neq 0.01$) for all dose levels and were 96%, 95%, and 90% of control for 30 ppm, 70 ppm, and 140 ppm, respectively. For the males, the brain weight of the 140 ppm group only was reduced (94% of control, $p \neq 0.05$). Histological findings included: brain lesions at multiple sites (four 140 ppm males; neuronal loss, neuronal necrosis, malacia); peripheral nerve degeneration and/or vacuolation (140 ppm, 2/sex affected; one 30 ppm female affected); and olfactory epithelium dysplasia (140 ppm, 3/sex affected). The NOAEL was < 30 ppm based on reduced brain weight at the lowest dose tested. Neuro-behavioral testing effects were observed at $\$ 70$ ppm. Submission of "validation" training and positive control data (Gill, 1989 a, b, c, and d) generated 5 years before the methyl bromide testing using different personnel than those in the methyl bromide testing (Driscoll *et al.*, 1994) were considered inadequate. Therefore, this study is considered unacceptable and not upgradeable by DPR.

III.C.2. Oral - Rat

In a chronic toxicity range-finding study, Sprague-Dawley rats (15/sex/group) were given microcapsules containing methyl bromide (0, 0.1, 1.0, 10, or 100 ppm) mixed in the diet daily for 4 weeks (Tompkins, 1995). The calculated average dosages for both sexes were 0.009, 0.09, 0.8, and 8 mg/kg/day. No treatment-related effects were noted in clinical signs, hematology, serum chemistry, or at necropsy. There was decreased food consumption and body weight gain throughout the study in the 100 ppm group. Food consumption was 91 to 92% of control for the 100 ppm male for each of the 4 weeks. The body weight gains of the 100 ppm males were 85%, 87%, and 88% of control for weeks 0 to 1, 0 to 2, and 0 to 3, respectively. The only significant ($p \neq 0.05$) decrease (76% of control) in the body weight gain of females was at 0 to 2 weeks for the 100 ppm group. The NOEL was 10 ppm (0.8 mg/kg/day) based on decreased food consumption and body weight gain at 100 ppm (8 mg/kg/day).

Wistar rats were given methyl bromide ($>98\%$ pure; 0, 0.4, 2, 10, or 50 mg/kg/day) by gavage 5 days per week for 13 weeks (Danse *et al.*, 1984). At 50 mg/kg/day, there were significantly ($p \neq 0.05$) reduced body weight gain (69% and 73% of control values for 6 and 12 weeks, respectively) in the male, decreased food consumption (77-82% of control values) in both sexes, and decreased red blood cell concentration (93% of control values) in the males. Histological examination showed that there was an increase in the incidence and severity of hyperplasia of the stratified squamous epithelium of the forestomach. At 10 mg/kg/day, the hyperplasia was described as slight in 9 of 10 males and 6 of 10 females. At 50 mg/kg/day, the

hyperplasia was predominantly strong and occurred in 8 of 10 of males and all females. In the original report, squamous cell carcinoma of the forestomach was reported for 13 of 20 animals treated with 50 mg/kg/day. However, re-analysis of the histological slides by a National Toxicology Program Panel concluded that the lesions were non-neoplastic (Boorman, 1984). Histological changes in the lungs were focal interstitial pneumonia in the 10 and 50 mg/kg/day groups, and slight atelectasis in the 50 mg/kg/day groups. The NOEL was 2 mg/kg/day (1.4 mg/kg/day as daily dose from adjustment for 5 days per week dosing) based on forestomach epithelial hyperplasia.

The reversibility of the lesions in the stomach was studied by Boorman *et al.* (1986). Wistar rats were given methyl bromide (>99% pure, 0 or 50 mg/kg) by gavage 5 days per week for 13, 17, 21 or 25 weeks. Pseudoepitheliomatous hyperplasia of the forestomach was observed in all treatment periods; however, there was no significant increase in the incidence with prolonged exposure. Evidence of early squamous cell carcinoma was detected in one animal after 25 weeks. In another experiment, rats were treated for 13 weeks and were sacrificed immediately afterward, 4 weeks, 8 weeks, or 12 weeks after treatment. There was an apparent regression of the hyperplasia formed at 13 weeks, as no incidence of hyperplasia was observed in animals sacrificed 12 weeks after treatment. However, adhesions, fibrosis, and mild acanthosis remained in the forestomach. The dosage for lowest-observed-effect level (LOEL) was 50 mg/kg/day (35.7 mg/kg/day as an adjusted daily dose).

In a study similar to that conducted by Boorman *et al.* (1986), Wistar rats were given methyl bromide (purity not specified) by gavage at 0, 25, or 50 mg/kg/day for 4 to 17 weeks and allowed a recovery period of 4 to 9 weeks before necropsy was performed (Hubbs, 1986). In the non-glandular stomach, the squamous epithelial portion showed ulceration and pseudoepitheliomatous hyperplasia characterized by hyperkeratosis, acanthosis, and epithelial peg formation. Fibrosis was found in the muscularis mucosa or tunica muscularis. In the glandular stomach, there were submucosal lymphoid aggregates and non-lymphoid mononuclear cell infiltrates in the lamina propria. At 4 to 9 weeks after treatment, there was regression of the hyperplasia in the stomach but not the muscularis fibrosis. The LOEL was 25 mg/kg/day (17.9 mg/kg/day as an adjusted daily dose).

III.C.3. Inhalation - Mouse

In a range-finding study for a 90-day study, B6C3F1 mice (10/sex/group) were exposed to methyl bromide (99.8% pure; nominal concentrations of 0, 12, 25, 50, 100, or 200 ppm) for 6 hours per day, 5 days per week, for 14 days (NTP, 1992; Drew, 1983). Mortality occurred only at the 200 ppm groups with 4 of 10 females and 1 of 10 males surviving until terminal sacrifice. Bloody urine (both sexes) and mild hyperemia in the lungs, liver, and kidneys (females) were observed in the 200 ppm groups. No urinary bladder lesion was detected to account for the hematuria. Neurotoxicity, manifested as trembling, jumpiness, and paralysis was noted for all exposed groups but was more prominent for the groups treated at 50 ppm and higher. There are questions as to the validity of the observations at the lower doses (12 and 25 ppm) since the observer had been cautioned to look for behavioral changes (Drew, 1983).

A decrease in body weight and neurotoxicity were also observed when B6C3F1 mice were exposed to methyl bromide (99.8% pure, a nominal concentration of 160 ppm) for 6 hours per day, 5 days per week for 2 weeks in a target organ toxicity study (Eustis *et al.*, 1988). The

experiment was intended for 6 weeks but was terminated after 2 weeks since more than 50% of the animals were dead after 10 and 8 exposures for the males and females, respectively. Clinical signs included red urine, lethargy, and neurological effects (curling and crossing of the hindlimbs, forelimb twitching and tremors). Treated male and female body weights at termination were only 74% and 82% of the control values ($p \# 0.001$, both sexes), respectively. Absolute organ weights which were significantly ($p \# 0.05$) lower in both sexes (unless noted) included: lungs, heart, spleen (males), right kidneys (males), thymus, brain, and liver. Histological examination of the tissues showed lesions in the brain (necrosis and loss of neurons in the cerebellum and cerebral cortex), kidney (nephrosis, dilatation and increased cytoplasmic basophilia), testes (degeneration), nasal cavity (degeneration and atrophy, males), heart (cardiomyopathy), adrenal gland (atrophy of the inner-zone of the adrenal cortex, females), thymus (atrophy), and spleen (lymphoid depletion and hematopoiesis).

When B6C3F1 mice were exposed to methyl bromide (99.8% pure; nominal concentrations of 0, 10, 20, 40, 80, or 120 ppm) for 6 hours per day, 5 days per week for 13 weeks, the survival rates were 83% for the 120 ppm males and 100% for the females (NTP, 1992; Eustis, 1992). In the 120 ppm group, body weight gain was 58% of control ($p \# 0.05$, males), and brain weights were 92 to 94% of control ($p \# 0.01$, both sexes). Clinical signs (severe curling, crossing of the hindlimbs, and twitching of the forelimbs) were observed in the 120 ppm groups, with greater severity occurring in the males than in the females. Alterations in a few of the neurobehavioral responses were observed, primarily in the 80 ppm groups; no data were reported for the 120 ppm groups. Significant ($p \# 0.05$) changes in the hematological parameters which were dose related included: decreased mean cell hemoglobin (40 to 120 ppm males, 120 ppm females), decreased mean cell volume (40 to 120 ppm males), increased erythrocyte count (40 to 120 ppm males), and increased hemoglobin (120 ppm males). No compound-related lesions were observed by histological examination. The NOEL was 20 ppm for changes in blood parameters at 40 ppm and neurobehavioral responses at 80 ppm.

III.C.4. Inhalation - Rabbit

Rabbits (strain unspecified) were exposed to methyl bromide (99% pure, nominal concentrations of 17, 33, or 66 ppm) by inhalation for 7.5-8 hours per day, 5 days per week during a 6-month period (Irish *et al.*, 1940). No adverse effects were observed at 17 ppm. Paralysis and lung damage occurred at the next dose (33 ppm) and higher. At 66 ppm after 14 to 46 exposures, 28 of 42 rabbits showed severe paralysis, and 14 of 42 rabbits died. Rabbits appeared to recover from the paralysis if they were removed from exposure at the first sign of paralysis.

New Zealand white rabbits (2/control, 6/treated) were exposed to methyl bromide (99.9% pure; a nominal concentration of 65 ppm for 7.5 hours per day, 25 hours per week for 4 weeks (Anger *et al.*, 1981). The data were presented in graphs. After 4 weeks of exposure, the methyl bromide group showed a decrease in body weight which continued to decline at test week 5 (1 week after continuous exposure). After 4 days of exposure, eye blink amplitude were decreased in both control and treated groups. The decline in the control returned to pre-test level in the following week. However, in the treated group, eyeblink amplitude was decreased after additional exposure. Furthermore, during the fourth week of exposure, the rabbits did not use the hind limbs or groom themselves. The recovery of normal functions was complete for sciatic

nerve conduction velocity and partial for ulner nerve conduction velocity and eyeblink amplitude when the treated rabbits were tested 6-8 weeks after exposure (Russo *et al.*, 1984).

Male New Zealand white rabbits (2 control, 6 treated) were exposed to methyl bromide (99% pure, a nominal concentration of 27 ppm) at 7.5 hours per day, 25 hours per week, for 30 weeks (Russo *et al.*, 1984). There were no treatment-related changes in sciatic and ulnar nerve conduction velocities and eyeblink reflex amplitude.

III.C.5. Inhalation - Dog

Beagle dogs (2-4 dogs/sex/group) were exposed to methyl bromide (100% pure) by whole body inhalation at 7 hours per day, 5 days per week, for three exposure durations. The durations of exposure were: 23 to 24 exposure days (0, 26, 53, or 103 ppm), 30 exposure days (24 exposure days at 11 ppm, then 6 exposure days at 158 ppm), or 34 exposure days (0 or 5 ppm) (Newton, 1994b). Air concentrations stated were measured concentrations. Serum bromide levels increased with the dose at \$ 26 ppm. Body weight loss (25%) and neurotoxicity were seen in the dogs exposed to 158 ppm. These dogs showed decreased activity on the second exposure day (the first dose was on a Friday and the second dose was on the following Monday). They were reported in poor condition during the final (6th) exposure (Table 5). The next day, three 158 ppm males had to be sacrificed due to severe toxicity (opisthotonos, irregular gait, opening and closing of the jaws and convulsions). The remaining 158 ppm dogs exhibited: nystagmus, intention tremors, ataxia, irregular gait and depression. Elevated levels of protein and bilirubin were measured in the urine of the 158 ppm dogs. Histological examinations showed that each of the 158 ppm dogs had cerebellar lesions (vacuoles in the granular layer) and olfactory degeneration; the males also had adrenal cortex findings (zona fasciculata, cytoplasmic vacuoles).

A decrease in body weight gain and less severe neurotoxicity (tremors, emesis, decreased activity during exposure but not post-exposure) were seen in the 103 ppm dogs. The loss in body weight was statistically ($p \neq 0.05$) significant for all weekly measurements for males and from week 2 on for females. A decrease in activity was noted starting on exposure day 9 involving most or all of the animals. Tremors were observed in 1 of 8 dogs on day 10. Emesis was observed in 1 dog on days 9 and 10. One 103 ppm male exhibited a cerebellar lesion similar to that seen in the 158 ppm dogs. In the 53 ppm group, a decrease in activity during exposures also was noted (in 2 dogs), starting on exposure day 14. However, no abnormal findings were reported for the 53 ppm group in post-exposure examinations. The NOAEL for 23-24 exposure days was 53 ppm based on neurotoxicity observed at 103 ppm and 158 ppm.

The female dogs which were exposed the longest to methyl bromide (5 ppm group) had reduced absolute spleen weights (55% of control compared to 75% of control for the male group) and two 5 ppm females were observed by an animal neurologist at the end of test week 6 to be less responsive than expected. The neurologist had examined the dogs on two other occasions: pre-test and at the end of week 4. The LOAEL for 34 exposure days was 5 ppm for decreased spleen weight (females) and decreased responsiveness (females).

Acute and short-term NOELs were also established from these experiments. The acute NOEL was 103 ppm for decreased activity seen in the 158 ppm group after 2 exposures and

severe neurotoxicity after 6 exposures (Table 3). For 14 days of exposure, the NOEL was 26 ppm for decreased activity observed at 53 ppm (Table 5).

Table 5. The neurotoxicity of methyl bromide in dogs after subchronic exposure^a.

Concentrations mean \pm sd (ppm)	Onset	Clinical Signs and Incidences ^c	Clinical Signs with Additional Exposure
158 \pm 7 ^b	day 2	decreased activity (8/8)	severe neurotoxicity, cerebellar lesions (8/8)
103 \pm 9	day 9	decreased activity (3/8)	day 9 to 10: emesis (1/8), tremor (1/8), decreased activity (3/8); week 5: cerebellar lesions (1/8)
53 \pm 4	day 14	decreased activity (2/8)	
26 \pm 1	23-24 exposures	no effects observed	
5 \pm 0.4	34 exposures	decreased responsiveness (2/8)	

a/ Data from Newton, 1994b.

b/ The dogs were exposed to 11 ppm for 24 exposure days, then 158 ppm for 6 exposure days.

c/ Incidences as number of dogs affected/total are shown in parentheses.

III.C.6. Inhalation - Guinea Pig

Guinea pigs exposed to methyl bromide (99% pure; nominal concentrations of 0, 33, 66, 100, or 220 ppm) for 7.5 to 8 hours per day and 5 days per week showed severe intoxication only at 100 ppm and higher concentrations (Irish *et al.*, 1940). At 100 ppm, 4 of 11 animals died after 64 to 91 exposures. However, pulmonary damage was insignificant. At 200 ppm, 14 of 16 animals died after only 1 to 3 days of exposure. All showed difficulty in breathing and lung damage.

III.C.7. Inhalation - Monkey

Monkeys were exposed to methyl bromide (99% pure; nominal concentrations of 0, 33, 66, 100 ppm) for 7.5 to 8 hours per day and 5 days per week (Irish *et al.*, 1940). At 100 ppm after 11 exposures, severe convulsions were observed in the one monkey tested. At the lower concentration (66 ppm), six monkeys were tested: one became paralyzed after 25 exposures and two others became paralyzed after 45 to 57 exposures. No toxicity was observed in four monkeys exposed to 33 ppm methyl bromide for 116 to 259 exposure days.

III.C.8. Additional Studies

Additional studies for the consideration of subchronic toxicity are described in the **III.D. CHRONIC TOXICITY**, **III.F. REPRODUCTIVE TOXICITY**, and **III.G. DEVELOPMENTAL TOXICITY** and are summarized in Table 6.

Table 6. The no-observed-effect levels (NOELs) and lowest-observed-effect levels (LOELs) for subchronic effects of methyl bromide^a.

Route/ Species	Exposure Duration ^b	Study NOEL/LOEL ppm	Human equivalent NOEL/LOEL ppm	Effects	Ref. ^d
INHALATION					
Rat	8h/dx5d/w \$6 exposures	66/100	31/47	convulsions, lung lesion	1
Rat	7.5h/dx4d/w 100 h	65/- ^d	24/-	no effects on neurobehavioral tests	2
Rat	24 h/dx3w	5/10	10/21	9 brain monoamine levels	3
Rat	4h/dx5d/wx6w	200/300	50/75	paralysis, death	4
Rat	6h/dx5d/wx3w	<160/160 ^e	<60/60	death, neurotoxicity, and tissue damage	5
Rat	6h/dx5d/wx13w	60/120	22/45	startle response, 9grip strength olfactory epithelial dysplasia	6
Rat	6h/dx5d/wx13w	<30/30 30/70	<11/11 11/26	9 brain weight 9 motor activity	7
Rat ^f	6h/dx5d/w 132-145 days	3/30 3/30	2/20 2/20	F ₁ parental-9 fertility progeny-9 body weight, 9 brain weight (F ₁), 9 cerebral cortex width (F ₁)	8*
Mouse	6h/dx5d/w x 2w	<160/160 ^d	<112/112	death, tissue degeneration, 9 body and organ weights, neurotoxicity	5
Mouse	6h/dx5d/wx13w	20/40	14/28	hematology and neurobehavioral changes at higher doses	6
Mouse ^g	6h/dx5d/wx20w	33/100	23/98	brain, heart, sternum, and olfactory epithelium lesions, neurotoxicity	6

a/ Bolded studies are those selected to derive the critical NOELs for risk characterization.

b/ Duration of exposure is indicated as : h=hours, d=days, w=weeks, m=months.

c/ Human equivalent NOEL/LOELs were calculated by adjusting for respiration rate differences and duration of exposure (Appendix G). The equivalent levels were those for children only (except reference 8). The equivalent levels would be 2-fold higher since the adult respiration rate is lower (0.26 m³/kg/day compared to the child's 0.46 m³/kg/day). For reference 8, the equivalent levels were those for adults since the effects were related to pregnancy.

d/ * after the reference number indicates the study was acceptable to DPR according to FIFRA guidelines. References: 1. Irish *et al.*, 1940; 2. Anger *et al.*, 1981; 3. Honma *et al.*, 1982; 4. Kato *et al.*, 1986; 5. Eustis *et al.*, 1988; 6. NTP, 1992 and Eustis, 1992; 7. Norris *et al.*, 1993a and b; 8. American Biogenics, Corp., 1986.

e/ The only dose studied or the dose at which an effect was observed at the specified duration but not necessarily the LOEL.

f/ Study described in **III.F. REPRODUCTIVE TOXICITY.**

g/ Study described in **III.D. CHRONIC TOXICITY.**

Table 6. The no-observed-effect levels (NOELs) and lowest-observed-effect levels (LOELs) for subchronic effects of methyl bromide (continued)^a.

Route/ Species	Exposure Duration ^b	Study NOEL/LOEL ppm	Human equivalent NOEL/LOEL ppm	Effects	Ref. ^c
INHALATION					
Rabbit	8h/dx5d/w x<6 mon	17/33	4/9	paralysis, lung lesion	1
Rabbit	8h/dx5d/w x 14-46 exp	33/66	9/17	paralysis, death	1
Rabbit	7.5h/dx4d/wx4w	<65/65 ^e	<14/14	9 nerve conduction velocity, 9 eyeblink reflex, paralysis	2
Rabbit	7.5h/dx4d/wx30w	27/>27	7/>7	no change in nerve conduction velocity, or eyeblink reflex 3	
Rabbit^f	7h/d, gd1-15	20/ 70	7/24	convulsion, paresis, death after 1 week	4
Rabbit ^f	6h/d, gd 7-17	70/140	21/41	neurotoxicity	5
Rabbit ^f	6h/d, gd 7-19	40/80	12/23	neurotoxicity on gd 19	6*
Dog	7h/dx14d	26/53	5/9	9 activity	7
Dog	7h/dx34d (6w)	<5/5^e	<1/1	9 responsiveness, 9 spleen weight	7
Guinea Pig	8h/dx5d/wx47 exp	33/66	7/14	death	1
Monkey	8h/dx5d/wx11 exp	66/100	14/21	convulsion	1
Monkey	8h/dx5d/wx25 exp	33/66	7/14	convulsion, paralysis	1
ORAL					
Rat	capsules, dailyx1w	NOEL/LOEL in <u>unadjusted</u> 0.8/8	mg/kg/day <u>adjusted^g</u> 0.8/8	9 body weight gain and food consumption	8
Rat	gavage, 5d/wx13w	2/10	1.4/7.1	forestomach-hyperplasia	9
Rat	gavage, 5d/wx13-25w	<50/50	-/35.7	forestomach-hyperplasia, tumor	10
Rat	gavage, 5d/wx4-17w	<25/25	/17.9	forestomach-hyperplasia	11

a/ Bolded studies are those selected to derive the critical NOELs for risk characterization.

b/ Duration of exposure is indicated as : h=hours, d=days, w=weeks, exp=exposures, gd=gestational day.

c/ Human equivalent NOEL/LOELs were calculated by adjusting for respiration rate differences and duration of exposure (Appendix G). The equivalent levels were those for children only. The equivalent levels for adults would be 2-fold higher since the adult respiration rate is lower (0.26 m³/kg/day compared to the child's 0.46 m³/kg/day).

d/ * after the reference number indicates the study was acceptable to DPR according to FIFRA guidelines.

References: 1. Irish *et al.*, 1940; 2. Anger *et al.*, 1981; 3. Russo *et al.*, 1984; 4. Sikov *et al.*, 1981; 5. Breslin *et al.*, 1990a; 6. Breslin *et al.*, 1990b; 7. Newton, 1994b; 8. Tompkins, 1995; 9. Danse *et al.*, 1984; 10. Boorman *et al.*, 1986; 11. Hubbs, 1986.

e/ The only dose studied or the dose at which an effect was observed at the specified duration.

f/ Studies described in **III.G. DEVELOPMENTAL TOXICITY**.

g/ The dosages were adjusted for experiments with dosing 5 days per week to 7 days per week.

III.D. CHRONIC TOXICITY AND ONCOGENICITY

Summary: The nasal cavity, brain, and heart were major target organs in rodents after chronic inhalation exposure to methyl bromide. Olfactory epithelial damage (hyperplasia, metaplasia, and necrosis) and myocardial degeneration were observed in rats and mice. Cerebellar and cerebral degenerations were detected in mice while reduced brain weight was observed in rats. When rats were exposed to methyl bromide in microcapsules mixed in the feed, the primary effect was body weight reduction. Possible treatment-related lesions were found in the spleen, liver, pancreas, and lungs. In dogs fed methyl bromide-fumigated feed, anemia and decreased serum calcium were observed. A summary of the chronic toxicity studies is presented in Table 11.

III.D.1. Inhalation - Rat

Wistar rats (90/sex/group) were exposed to methyl bromide (98.8% pure; nominal concentrations of 0, 3, 30, or 90 ppm) 6 hours per day, 5 days per week (Reuzel *et al.*, 1987 and 1991). The main group for each dose consisted of 50 rats of each sex and were exposed to methyl bromide for 29 months. There were 4 satellite groups (10/sex/group except noted): week 13-14 hematology and blood chemistry analyses, week 41 behavioral effects, 1 year interim sacrifice, and 2 years interim sacrifice.

In the 90 ppm group, the 2 year and 2.5 year mortality rates for both sexes (male/female) were 52%/46% and 84%/86%, respectively. These rates were considered higher than those for the control groups which were 32% for 2 years and about 72% for 2.5 years, for both sexes. The mean body weight of the 90 ppm female group was significantly ($p \# 0.05$) lower than that of the controls throughout most of the study with the maximal reduction (12%) at the end of the study. The mean body weight of the 90 ppm male group was significantly ($p \# 0.05$) decreased (maximum of 6%) on occasion. Absolute kidney weights were significantly ($p \# 0.05$) decreased in the 30 ppm (89% of control) and 90 ppm (84% of control) females, and in the 90 ppm males (85% of control) when compared to control values at the 1 year sacrifice. Absolute brain weights of the 90 ppm females were reduced at the 1-year and 2-year sacrifices (Table 7). At 29 months, the absolute brain weights were significantly decreased in the 30 ppm males and females, and the 90 ppm females. Brain weight reduction also was seen in the 90 ppm male group at 29 months but was not statistically significant probably due to the small number of survivors ($n=8$). The NOAEL was 3 ppm for brain weight reduction.

There were increased incidences of heart thrombi and myocardial degeneration in rats that died or were killed when moribund (Table 7). These lesions may be the cause of increased mortality in the high dose groups. Statistical comparison to the control group did not identify any increased tumor incidences. Few tumors were observed: brain glioma in the 30 ppm group (1 male and 1 female), granular cell myoblastoma in the 30 ppm group (2 males), and 90 ppm group (1 male, and 2 females), spinal cord glioma in the control (1 male) and 90 ppm group (1 female). The historical control data (1974-1988) for Wistar rats showed the following incidences (male and female, respectively): 8/873 and 3/876 for brain glioma, 2/685 and 1/701 for spinal cord glioma, and 2/873 and 8/876 for granular cell myoblastoma.

At 12-24 months, the incidences of nasal cavity lesions of the 30 and 90 ppm groups were significantly ($p \# 0.05$) different from those of the control group (Table 7). At 24-29 months, there was a dose-related increase in the incidences of nasal cavity degeneration/hyperplasia, heart lesions (thrombus, myocardial degeneration, and cartilaginous metaplasia), esophageal hyperkeratosis, and stomach hyperkeratosis in all treatment groups. The finding of epithelial cell degeneration/basal cell hyperplasia in the olfactory epithelium of the nasal cavity was both dose- and time-related in incidence and severity (Table 7). The lesions were described as very slight at the lower doses to moderate at the higher doses. There was thinning (atrophy) of the epithelial layer and the formation of cyst-like glandular structures in the submucosa layer. The LOAEL for the basal cell hyperplasia/degeneration were >90 ppm, 30 ppm, and 3 ppm for exposures lasting 12 months, 12-24 months, and 24-29 months, respectively. The results from the reexamination of the nasal-cavity histological slides (Hardisty, 1997) did not change the LOAELs because: (1) the rereading of the slides was not conducted in accordance with standard procedures for a peer review, (2) dose response for incidence and severity remained the same with effects observed at 3 ppm. This study was considered acceptable to DPR according to FIFRA guidelines.

Fischer-344/DuCrj rats (50/sex/group) were exposed to methyl bromide (99.9% pure; 0, 4, 20, or 100 ppm) by inhalation 6 hours per day, 5 days per week for 104 weeks (Gotoh *et al.*, 1994). Results in this study were reported as summary data in a brief publication. There were no effects on survival. The mean body weights of the 100 ppm (both sexes) and 20 ppm (males) were stated to be lower than controls; however, no data were included in the report. The primary effect was increased incidences ($p \# 0.05$) of necrosis (100 ppm males), inflammation (4 ppm males; 100 ppm females), and respiratory metaplasia (100 ppm males; 4 ppm females) of the olfactory epithelium. These effects occurred in the males at lower doses than in the females. The LOEL for nasal cavity inflammation and respiratory metaplasia was 4 ppm. Histological examination of the tissues showed increased incidences of pituitary adenoma (100 ppm males) and adrenal gland pheochromocytoma (4 ppm females). The significance of these findings as well as non-neoplastic findings require the evaluation of individual data, historical control data, and subchronic studies. The study was considered unacceptable and upgradeable by DPR.

Table 7. The effects of methyl bromide in rats after chronic inhalation exposure^a.

Exposure duration	Methyl Bromide Concentration (ppm)				
	0	3	30	90	
MALE					
Mean Absolute Organ weight (g)^b					
Brain Weight:	1 year	2.01±0.03	2.12±0.01*	2.03±0.03	1.93±0.03
	2 year	2.09±0.07	2.15±0.02	2.08±0.03	1.99±0.06
	29 months	2.15±0.02	2.11±0.03	2.03±0.04*	2.02±0.05
Incidences					
Nasal Cavity:	12-24 months	6/30++ (20%)	1/26 (4%)	11/35 (31%)	22/39** (56%)
	24-29 months	4/36++ (11%)	12/37* (32%)	16/32*** (50%)	20/28*** (71%)
Incidences					
Heart:	died before end of study ^c	1/33++	2/25	5/34	12/41**
	Cartilaginous metaplasia				
	Myocardial degeneration -moderate/severe	21/33++	16/25	8/34	36/41*
	Thrombus	4/33++	3/25	10/34	20/41**
<hr style="border-top: 1px dashed black;"/>					
FEMALE					
Mean Absolute Organ weight (g)^b					
Brain Weight:	1 year	1.94±0.02	1.95±0.03	1.85±0.04	1.81±0.03**
	2 year	1.97±0.02	1.91±0.01	1.88±0.03	1.84±0.05*
	29 months	2.01±0.02	1.96±0.02	1.92±0.02*	1.77±0.06**
Incidences					
Nasal Cavity	12-24 months	7/38++ (18%)	5/34 (15%)	16/39* (41%)	26/42*** (62%)
	24-29 months	5/40++ (13%)	16/42** (38%)	15/38** (39%)	25/35*** (71%)
Incidences					
Heart:	died before end of study ^c	4/41	10/33	2/35	14/51*
	Cartilaginous metaplasia				
	Myocardial degeneration -moderate/severe	13/41++	5/33	5/35	38/51**
	Thrombus	5/41++	8/33	1/35	20/51*

* , ** , *** ; + , ++ Level of statistical significance, p # 0.05 (* or +), p # 0.01 (** or ++), or p # 0.005 (***). Significance for incidences was based on a dose-weighted chi-square trend test and the Fisher's Exact Test. For brain weights, significance was based on ANOVA and Dunnett tests.

^{a/} Data from Reuzel *et al.* (1987, 1991). Incidence rates were the number of animals affected/number of animals examined. Rats in the 12-24 month group were those in the 1-year and 2-year sacrifice groups, and in the main group which died before two years. Rats in the 24-29 month group were those in the main group which died between days 736 and terminal sacrifice, and those at terminal sacrifice.

^{b/} The number of animals per group for the 29 months data were: 15, 25, 16, and 8 for the males; and 18, 25, 24, and 9 for the females for 0, 3, 30, and 90 ppm, respectively.

^{c/} Incidences of heart lesions in those animals dead before the end of the study.

III.D.2. Dietary - Rat

Sprague-Dawley rats (70/sex/group, except for 0.5 and 2.5 ppm with 50/sex/group) were given methyl bromide in microcapsules dispersed into the granular feed for presentations to the animals (Mertens, 1997). Corn oil containing methyl bromide was microencapsulated using starch and sucrose. Two types of microcapsules were produced. One was a blend of 7 production runs; it had a methyl bromide content of 0.48% w/w. The second type was a blend of five production runs; its methyl bromide content was 3.44% w/w. The two types of microcapsules differed also in terms of corn oil, starch, and sucrose contents and age of the material at start of testing. Nominal methyl bromide concentrations in the diet were as follows: 0 (basal diet), 0 (diet containing placebo microcapsules), 0.5, 2.5, 50 and 250 ppm. The blend containing 0.48% methyl bromide was used to prepare the two low doses while the blend containing 3.44% was used to prepare the two high doses. The highest dose tested was selected on the basis of a two-week range-finding study. The daily ration of feed varied as follows: for test weeks 0-65, males and females each received 30 and 23 g, respectively; for test weeks 66 -104, males and females received 35 and 30 g, respectively. One outcome of this feeding strategy appeared to have been that a fraction of the animals in the control and 0.5 to 50 ppm groups had their feed consumption restricted during the first 65 weeks of the study. In test week 53, interim sacrifices were performed on 18-20 rats/sex for the following dose levels: 0 (basal diet), 0 (placebo microcapsules), 50 and 250 ppm. Survival was statistically increased in the 250 ppm male group and in the 50 and 250 ppm female groups when compared to the placebo-microcapsule groups. Body weight was reduced in the 250 ppm groups; the reduction reached a maximum in the first weeks of testing in both sexes; a further reduction in body weight relative to the controls (placebo-microcapsule groups) did not occur despite continued exposure. Since a reduction in feed consumption occurred in the 250 ppm groups (both sexes) starting with the first exposure week, the body weight reduction would appear to be due mainly to the reduced feed consumption. No treatment-related effects were reported in the following areas: clinical observations, ophthalmology, hematology, serum chemistry or urinalysis. Effects on absolute organ weights (brain, kidneys, liver, testes/ovaries) and organ weights relative to body weight appeared to be due to the body weight reduction in the 250 ppm groups; this was true for animals sacrificed at test week 52 as well as for the survivors at the end of the study. Possible, treatment-related findings at necropsy were an increasing incidence of splenomegaly in the males (0 ppm, basal: 2/50; 0 ppm, placebo: 2/50; 0.5 ppm: 7/50; 2.5 ppm: 10/50; 50 ppm: 11/50; and 250 ppm, 3/50) and an increased incidence of dark red areas on the liver in the 50 ppm females surviving to test week 104 (0 ppm, basal: 5/20; 0 ppm, placebo: 3/19; 0.5 ppm: 8/22; 2.5 ppm: 4/24; 50 ppm: 14/27; and 250 ppm, 8/29). No statistical analyses were supplied for the histology data. Also, the lesion-incidence summary table did not present autolysis and lesion-grade data and may not have been corrected for tissues lost to autolysis. Possible treatment-related effects include: increased incidence of pancreatic acinar atrophy at 250 ppm (both sexes), increased incidence of adrenal cortical hypertrophy at 250 ppm (females), and increased incidence of pulmonary arterial mineralization at 50 ppm (females). Two rare tumor types, adenocarcinoma of the prostate and endometrial stromal sarcoma of the cervix, were seen at 4% incidence at 250 ppm. By experimental design, the histological examinations of the pancreas, prostate, spleen, adrenal glands, cervix, and uterus at the 0.5 to 50 ppm dose levels re limited to those rats that did not survive to terminal sacrifice. Autolysis was a frequent observation in the gastrointestinal organs in rats that did not survive to the end of the study (all groups, both sexes). While an increased incidence of spongiosis hepatis was seen in the 50

ppm females, the relationship of this lesion to angiectasis and the necropsy finding of dark red liver spots that also occurred at the 50 ppm dose level needs clarification. When first reviewed, the study was considered unacceptable pending the submission of the supplemental information regarding: range-finding study; analytical methods; cause and extent of autolysis; histological examinations for the lower dose groups; and clarification of liver gross and histological findings. Additional information were submitted and this study is considered marginally acceptable to DPR. The LOEL, as a conservative call, was 0.5 ppm based on the following: in the absence of histological data to the contrary, the instances of splenomegaly in the 0.5, 2.5 and 50 ppm male groups that have not been examined histologically were assumed to be due to lymphoma.

Fischer-344 rats were fed daily for 2 years feed fumigated with methyl bromide (99.5% purity) (Mitsumori *et al.*, 1990). After fumigation, the feed was exposed to air for 21 days until the methyl bromide level was < 20 ppm and the total bromide level was approximately 500 ppm. Diets containing 80 ppm and 200 ppm total bromide were prepared from the 500 ppm feed. Another group of rats was maintained on a diet containing 500 ppm potassium bromide. In the methyl bromide groups, there were no major adverse effects reported, except for a significant ($p \# 0.05$) decrease in body weight (3-6% from week 60 onward). Significant increases ($p \# 0.05$) in food and water consumption and urobilinogen were observed in both 500 ppm methyl bromide and potassium bromide treated diet groups. Since the purpose of the study was to determine the effects of residual bromide, not methyl bromide itself, this study was considered supplemental information by DPR.

III.D.3. Inhalation - Mouse

B6C3F1 mice (86/sex/group) were exposed to methyl bromide (99.8% pure; nominal concentrations of 0, 10, 33, or 100 ppm) 6 hours/day, 5 days/week, for 2 years (NTP, 1992; Eustis, 1992). There were interim sacrifices (~ 10/sex/group) at 6 and 15 months for the control, 10, and 33 ppm groups; and at 15 months for the 100 ppm females. Another group (10 mice/sex/group) was used for neurobehavioral testing every 3 months. The exposure of the 100 ppm group was stopped after only 20 weeks because of neurotoxicity and mortality. In this group, clinical signs indicative of neurotoxicity (tremors, paralysis, unusual gait, abnormal posture) were observed in 78% of the males and 43% of the females. The surviving animals in the 100 ppm group were observed for the duration of the study (2 years). Neurological signs in the 100 ppm groups often began to appear well after their exposure had stopped.

At the interim sacrifice of 6 months, there were no significant findings in the 10 and 33 ppm groups. At the 15-month interim sacrifice, treatment-related lesions in the brain, sternum, and heart were observed. These changes were similar to those observed after 2 years. While there was no evidence of carcinogenic effects, methyl bromide caused increased incidences of cerebellar and cerebral degeneration, myocardial degeneration, cardiomyopathy, sternal dysplasia, and olfactory epithelial necrosis and metaplasia (Table 8). Cerebellar degeneration, cerebral degeneration, myocardial degeneration, and olfactory epithelium necrosis were considered fatal lesions since higher incidences occurred in animals which died before terminal sacrifice. Heart lesions, either cardiac degeneration or chronic cardiomyopathy, were observed in 80% of the males and 69% of the females exposed to 100 ppm methyl bromide; also, the incidence of chronic cardiomyopathy in the male 33 ppm group (20%) was greater than that seen in the controls (8%). Sternal dysplasia was observed at low incidences (4-6%) in the 10 ppm and

33 ppm female groups and the 33 ppm male group but higher incidences (15-20%) in the 100 ppm groups (both sexes). Because of the rarity of this lesion, the finding at 10 ppm was considered significant. Degenerative lesions in the cerebellum were observed in 44% and 18% of the male and female 100 ppm groups, respectively. Cerebellar degeneration, sternal dysplasia, chronic cardiomyopathy, and olfactory metaplasia were observed in the 100 ppm survivors which were sacrificed at study week 104 (Table 9).

Despite the neurotoxicity observed in the 100 ppm mice before week 20, neurobehavioral testing at 3 months resulted in significant ($p \leq 0.01$) findings for only 5 and 3 of the 9 endpoints studied in the 100 ppm male and female groups, respectively. However, testing of the 100 ppm female group at 6 months resulted in significant ($p \leq 0.01$) findings in 3 of the endpoints though the survivors had not been exposed to methyl bromide for about one month. Neurobehavioral testing also found a significant ($p \leq 0.05$) decrease in locomotor activity in the 10 ppm and 33 ppm groups (both sexes) when tested at 6 months as well as 12 months (females only). Based on the neurobehavioral testing changes in locomotor activity and sternal dysplasia, the LOAEL was established at 10 ppm, the lowest test dose. This study was considered acceptable to DPR according to FIFRA guidelines.

BDF1 mice (50/sex/group) were exposed to methyl bromide (99.9% pure; 0, 4, 16, or 64 ppm) by inhalation 6 hours per day, 5 days per week for 104 weeks (Gotoh *et al.*, 1994). There were no significant differences in survival between the groups. Results in this study were reported as summary data in a brief publication. The mean body weights of the 64 ppm (both sexes) were stated to be lower than controls; however, no data were included in the report. The primary effect was increased incidences (30% compared with 0% in controls; $p \leq 0.05$) of atrophy (slight) of the granular layer of the cerebellum of the 64 ppm group. The NOEL for cerebellar lesion was 16 ppm. Histological examination of the tissues showed increased incidences of liver adenoma (4 ppm females) and lymphoma in lymph nodes (4 ppm females). However, the significance of these findings as well as non-neoplastic findings require the evaluation of individual data, historical control data, and subchronic studies. The study was considered unacceptable and upgradeable to DPR.

Table 8. The incidences of histological lesions in mice after chronic inhalation exposure to methyl bromide^a.

	Methyl Bromide Concentration (ppm)			
	0	10	33	100 ^b
Male				
Cerebellar degeneration	0/50++	0/50	0/50	31/70** (44%)
Cerebral degeneration	0/50++	0/50	0/50	11/70** (16%)
Sternal dysplasia ^c	0/50++	0/50	3/50 (6%)	14/70** (20%)
Myocardial degeneration	0/50++	0/50	0/50	32/70** (46%)
Chronic cardiomyopathy	4/50++ (8%)	7/50 (14%)	10/50 (20%)	24/70** (34%)
Olfactory epithelium, metaplasia	0/50	0/50	1/50 (25%)	2/69 (3%)
Olfactory epithelium, necrosis	0/50++	0/50	0/50	6/69* (9%)
% Survival at the end of the study ^d	82	74	80	23
Female				
Cerebellar degeneration	0/50++	0/50	0/50	11/60** (18%)
Cerebral degeneration	0/50+	0/50	0/50	2/60 (3%)
Sternal dysplasia ^c	0/50++	2/50 (4%)	2/50 (4%)	9/60** (15%)
Myocardial degeneration	1/50++ (2%)	0/50	0/50	7/59 (12%)
Chronic cardiomyopathy	2/50++ (4%)	4/50 (8%)	2/50 (4%)	34/59** (57%)
Olfactory epithelium, metaplasia	0/50++	0/50	0/50	5/60* (9%)
Olfactory epithelium, necrosis	0/50	0/50	0/50	1/60* (2%)
% Survival at the end of the study ^d	71	82	90	65

^{a/} Data were from NTP, 1992 and Eustis, 1992. Overall incidence was the number of animals with lesions/number of animals examined at site. Level of statistical significance, p # 0.05 (* or +) or p # 0.01 (** or ++), is indicated after each incidence. Significance indicated at the control value was based on a dose-weighted chi-square trend test; significance at the dosed group was based on the logistic regression test or the life table test when a lesion was considered to be fatal.

^{b/} Because of high mortality, exposure of the 100 ppm group was stopped at 20 weeks. The incidence rates are for all animals in this group and included incidences before and after the termination of exposure.

^{c/} Sternal dysplasia also was observed at the 15 month sacrifice (one male and one female from the 33 ppm group and one female in the 100 ppm group).

^{d/} The percentage of survival represented the survival rates adjusted for interim evaluation, neurobehavioral study animals, and accidental deaths.

Table 9. The incidences of histological lesions in survivors of chronic inhalation exposure to methyl bromide^a.

	Methyl Bromide Concentration (ppm)			
	0	10	33	100 ^b
Male Mice				
Cerebellar degeneration	0/40++	0/37	0/40	3/16** (19%)
Sternal dysplasia	0/40++	0/37	2/40 (5%)	12/16** (75%)
Chronic cardiomyopathy	4/40++ (10%)	4/37 (11%)	9/40 (23%)	9/16** (56%)
Olfactory epithelium, metaplasia	0/40++	0/37	1/40 (3%)	2/16 (13%)
Female mice				
Cerebellar degeneration	0/36++	0/41	0/45	4/40 (10%)
Sternal dysplasia	0/36++	2/41 (5%)	2/45 (4%)	7/40** (18%)
Chronic cardiomyopathy	1/36++ (3%)	4/41 (10%)	2/45 (4%)	27/39** (69%)
Olfactory epithelium, metaplasia	0/36++	0/41	0/45	5/40* (13%)

^{a/} Data were from NTP, 1992 and Eustis, 1992. Overall incidence was the number of animals with lesions/number of animals examined at terminal kill. Level of statistical significance, p # 0.05 (* or +) or p # 0.01 (** or ++), is indicated after each incidence. Significance indicated at the control value was based on a dose-weighted chi-square trend test; the pair-wise significance at the dosed group was based on the Fisher's Exact Test.

^{b/} Because of high mortality, exposure of the 100 ppm group was stopped at 20 weeks. Incidence rates indicated are those of the survivors at 2 years.

III.D.4. Dietary - Dog

Methyl bromide (purity not specified) fumigated feed with bromide levels of 0, 35, 75, or 150 mg/kg/day was fed to beagles (4/groups) daily for 1 year (Rosenblum *et al.*, 1960). Concentrations of methyl bromide *per se* were not determined. Lethargy and lower weight gain were observed in the high dose group; these findings were absent in the sodium bromide (100 mg/kg/day) group. Salivation, diarrhea, and death occurred in both the high dose methyl bromide and sodium bromide treated diet groups. This study was considered unacceptable to DPR due to too few animals, as well as the lack of feed analysis and necropsy/pathology data.

Beagle dogs (4/sex/dose except 8 dogs/sex at high dose) were given feed fumigated with methyl bromide 5 days per week for one year (Newton, 1996). Granular feed containing 10% corn oil was fumigated with methyl bromide at concentrations of 0, 7092, 20,000 or 116,279 ppm for one hour and degassed for one hour. Nominal residual methyl bromide levels in the feed-corn oil admixture one hour after the feed had been presented to the dogs were: 0, 0.5, 1.5 and 5.0 ppm. In addition, test feeds presumably contained fumigation-derived products (bromide, methylation adducts, methyl chloride). While the concentrations of reaction products were not measured, because of the experimental design, their concentrations in the low-dose feed versus high-dose feed may have varied by a factor of 16. Residual methyl bromide levels were selected on the basis of discussions between the Registrant and the U.S. EPA to achieve a "safety" study (i.e., the high dose was not set on the basis of toxicity data). A new analytical procedure was developed to determine residual methyl bromide; however, the adequacy of the new procedure could not be assessed pending submission of supplemental information.

There were no clear effects on survival, cage side observations, body weight or food consumption. Possible treatment-related effects included: decreased hemoglobin and (or) hematocrit at 3, 6 and (or) 12 months in the high-dose male group; and decreased serum calcium at 6 and 12 months in the mid- and high-dose male group (Table 10). The incidence of thyroid C-cell hyperplasia in the male control group was 1/4 versus 5/8 in high-dose male group. Statistically reduced absolute kidney weight was seen in the mid- and high-dose female groups; when viewed relative to terminal body weight or brain weight, kidney effects were not statistically significant. Due to the experimental design, the effects seen in this study may be due to residual methyl bromide and (or) its reaction products. The NOEL was 1.5 ppm based on decreased hemoglobin and/or hematocrit (anemia) at higher doses. When first reviewed, this study was considered unacceptable and upgrading would require the submission of the following: 1) supplemental information regarding the analytical method; 2) historical control data for thyroid C-cell hyperplasia in males; 3) histological examination of the thyroid in the low- and mid-dose male groups and the parathyroid in three high-dose females whose tissues were not examined originally; and 4) the statistical analyses of the hemoglobin, hematocrit and serum phosphate data. Subsequently, the registrant submitted data on the analytical method, histological data for the thyroid and parathyroid, and historical control data for the thyroid (CMA Methyl Bromide Industry Panel, 1998; Auletta, 1998). Based on these data, C-cell hyperplasia was no longer considered a possible adverse effect. Validation for the analytical method used in this study has been requested. This study is considered supplemental information by DPR.

Table 10. Selected hematology and clinical chemistry parameters in dogs exposed to methyl bromide fumigated feed^a.

Parameters Males	months	Nominal concentration in the diet (ppm)			
		0	0.5	1.5	5.0
Hematocrit %	3	51.7	50.7	50.5	47.7*
	6	50.8	51.0	49.1	46.8*
	12	57.2	54.9	51.7	51.5*
Hemoglobin g/dL	3	17.4	17.1	17.0	16.2*
	6	17.7	18.0	17.3	16.6*
	12	19.1	18.3	17.5	17.3*
RBC 10 ⁶ / μ L	3	7.69	7.41	7.41	7.24
	6	7.64	7.62	7.40	7.23
	12	8.22	7.81	7.45	7.59
Potassium mEq/L	3	4.3	4.3	4.3	4.4
	6	4.2	4.4	4.3	4.3
	12	4.6	4.6	4.5	4.5
Calcium mg/dL	3	9.9	9.9	9.5	9.8
	6	9.9	9.9	9.4**	9.6*
	12	10.0	9.8	9.4**	9.4**
Females					
Hematocrit %	3	45.9	48.4	47.9	49.6
	6	45.4	45.3	49.5	48.9
	12	48.0	49.2	53.7*	48.3
Hemoglobin g/dL	3	15.6	16.4	16.3	16.8
	6	15.9	16.0	17.4	16.9
	12	15.8	16.3	18.0*	16.2
RBC 10 ⁶ / μ L	3	6.76	7.21	7.26	7.46
	6	6.80	6.82	7.53	7.33
	12	6.79	7.04	7.85*	7.03
Potassium mEq/L	3	4.2	4.4	4.4	4.7**
	6	4.3	4.2	4.2	4.4
	12	4.4	4.5	4.3	4.5
Calcium mg/dL	3	9.7	10.1	10.1	10.0
	6	9.8	9.7	9.6	9.7
	12	9.6	9.7	9.9	9.5

^{a/} Data from Newton, 1996. *, ** Statistically different from control value at p<0.05 and p<0.01, respectively. There were 4 dogs/sex at all dose levels except 8 dogs/sex at the high dose. The estimated dosages (males/females) were: 0.06/0.07 mg/kg/day, 0.13/0.12 mg/kg/day, and 0.27 mg/kg/day for 0.5, 1.5, and 5.0 ppm, respectively.

Table 11. The no-observed-effect levels (NOELs) and lowest-observed-effect levels (LOELs) of methyl bromide from chronic oral and inhalation studies^a.

Species	Duration ^b	Study NOEL/LOEL ppm	Human equivalent ^c NOEL/LOEL ppm	Effects	Ref. ^d
<u>INHALATION</u>					
Rat	6h/dx5d/w x24-29m x12-24m	<3/3 3/30	<1/1 1/11	olfactory epithelial hyperplasia/ degeneration	1*
Rat	6h/dx5d/w x24m	<4/4	<1/1	nasal inflammation and respiratory metaplasia	2
Mouse	6h/dx5d/w x24m	<10/10	<7/7	neurobehavioral changes and sternal dysplasia	3*
Mouse	6h/dx5d/w x 104w	16/64	11/45	cerebellum atrophy	2
<u>ORAL</u>					
		mg/kg/day			
Rat ^e	2 years	not determined		potential treatment-related lesions in multiple organs	4
Dog ^f	1 year	1.5/5.0		decreased hemoglobin and hematocrit	5

a/ Bolded studies are those selected to derive the critical NOELs for risk characterization.

b/ Duration of exposure is indicated as: d=days, w=weeks, m=months.

c/ Human equivalent NOEL/LOELs were calculated by adjusting for respiration rate differences and duration of exposure (Appendix G). The equivalent levels were those for children only. The equivalent levels would be 2-fold higher since the adult respiration rate is lower (0.26 m³/kg/day compared to the child's 0.46 m³/kg/day).

d/ * after the reference number indicates the study was acceptable to DPR according to FIFRA guidelines. References: 1. Reuzel *et al.*, 1987 and 1991; 2. Gotoh *et al.*, 1994; 3. NTP, 1992; Eustis, 1992; 4. Mertens, 1997; 5. Newton, 1996.

e/ Methyl bromide was in microcapsules mixed in the feed. Additional information is needed to determine the NOEL and LOEL, in terms of mg/kg/day, and to confirm that the lesions are treatment related.

f/ The feed was fumigated with methyl bromide and then allowed to offgas.

III.E. GENOTOXICITY

Summary: Methyl bromide was genotoxic in several *in vitro* and *in vivo* assays. It was a base-pair substitution mutagen in the *Salmonella* assays. It was a direct-acting mutagen since a liver S-9 fraction was not required for mutagenicity. It caused micronuclei formation in female mice and an increased frequency of sister chromatid exchanges in CHO cells and in mouse bone marrow cells *in vivo*. It did not induce unscheduled DNA synthesis in rat hepatocytes or cause sperm abnormalities in mice. DNA alkylation was detected in both rats and mice after *in vivo* exposure by oral, intraperitoneal, or inhalation routes while DNA damage was found in the germ cells of rats after inhalation exposure. There was some evidence of genotoxicity in workers exposed to methyl bromide. Elevated levels of sister chromatid exchanges in lymphocytes and S-methylcysteine adducts in the blood were measured in soil fumigators. An increased frequency of hypoxanthine-guanine phosphoribosyl transferase gene (*hprt*) mutations in the lymphocytes and an increased incidence of micronuclei in oropharyngeal cells were observed in structural fumigators. The genotoxicity studies are summarized in Table 12.

III.E.1. Gene Mutation

Methyl bromide was mutagenic to *Salmonella typhimurium* strains TA100 and TA1535, but not TA98, TA1537 or TA1538 (Simmon *et al.*, 1977; Kramers *et al.*, 1985; Moriya *et al.*, 1983; NTP, 1992). It was also mutagenic to *Escherichia coli* strains Sd-4 and WP2 hcr (Djalali-Behzad *et al.*, 1981; Moriya *et al.*, 1983), and *Saccharomyces cerevisiae* (Mortelmans and Shepherd, 1980). The positive mutagenic response was independent of the presence of rat liver S9 fraction (Kramers *et al.*, 1985; Mortelmans and Shepherd, 1980; NTP, 1992) or hamster liver S9 fraction (NTP, 1992). Methyl bromide induction of sex-linked recessive lethality in *Drosophila melanogaster* was dependent on the exposure duration (Kramers *et al.*, 1985; and McGregor, 1981). Methyl bromide was also positive for the induction of forward mutations in the mouse lymphoma L5178Y assay at both the TK and HGPRT loci (Kramers *et al.*, 1985). Of all the studies summarized above, only the study by Mortelmans and Shepherd (1980) was considered acceptable to DPR.

III.E.2. Structural Chromosomal Aberrations

B6C3F1 mice were exposed to methyl bromide (0, 12, 25, 50, 100, and 200 ppm) for 6 hours per day, 5 days per week for 10 exposure days (NTP, 1992). Peripheral blood from female mice of the 100 and 200 ppm groups showed increased frequencies of micronuclei formation. No effect was observed in the male mice. In the second experiment, mice were exposed to methyl bromide (0, 10, 20, 40, 80, and 120 ppm) for 6 hours per day, 5 days per week for 4, 8, or 12 weeks. No increase in the frequency of micronucleated red blood cells was observed for either sex at any sampling times. This study was considered acceptable to DPR.

No induction of micronuclei formation was observed in rats (both sexes) given methyl bromide up to 123 mg/kg by a single intraperitoneal injection (Putman and Morris, 1991). This study was considered acceptable to DPR.

Methyl bromide did not cause dominant lethal mutations in male rats nor structural chromosomal aberrations in the bone marrow cells of rats (both sexes) exposed to methyl bromide by inhalation up to 70 ppm for 5 days (McGregor, 1981).

III.E.3. Other Genotoxic Effects in Experimental Animal Studies

Methyl bromide did not induce unscheduled DNA synthesis in rat hepatocytes (Kramers *et al.*, 1985) or human embryonic intestinal cells (McGregor, 1981) under *in vitro* conditions, or cause sperm abnormalities in mice after inhalation exposure (McGregor, 1981).

Methyl bromide caused a dose-related increase in the frequency of sister chromatid exchanges in Chinese hamster ovary cells *in vitro* (Rounds, 1980) and in bone marrow cells in female mice after inhalation exposure (6 hours/day and 5 days/week) to methyl bromide at 100 and 200 ppm for 10 exposures (NTP, 1992). However, no increase in the frequency of sister chromatid exchanges was observed for either sex exposed to methyl bromide up to 120 ppm for 12 weeks (NTP, 1992). The NTP (1992) study was considered acceptable to DPR.

In vivo exposure of mice to methyl bromide resulted in the alkylation of tissue DNA. After exposure to methyl bromide (>98% pure) by either inhalation or intraperitoneal routes, 7-methyl-guanine in liver and spleen DNA and methylated cysteine in hemoglobin and liver protein were detected (Djalali-Behzad *et al.*, 1981).

In the study by Gansewendt *et al.* (1991), rats were exposed to methyl bromide (96% pure) by inhalation for 6 hours (263 ppm for females, and 131 ppm for males), or by gavage (8.3 μ moles or 0.8 mg/kg for females and 0.58 mg/kg for males). From both routes of exposure, 7-methyl guanine and O⁶-methyl guanine were detected in the DNA from the liver, lung, stomach, and forestomach. The DNA adducts were considered a systemic genotoxic effect since high concentrations of the adducts were found in the stomach and forestomach DNA for both routes.

Methyl bromide (>99% purity; 0, 77, 153, or 258 ppm) was administered to Fischer 344 rats (5/group) by whole body inhalation (6 hours/day for 5 days) (Bentley, 1994). Animals were sacrificed one hour or one day after the 5th exposure. Damage to testicular DNA was determined by the alkaline-elution assay. The positive control group consisted of rats given methyl methanesulfonate. Mean body weights for the 150 and 250 ppm groups were reduced during the experiment. On the day after the 5th exposure, the mean body weight of the 150 and 250 ppm groups were 97% and 78%, respectively, of the pretreatment levels. On days 5 and 6, two rats in the 250 ppm group died, and a third rat was moribund. Signs of neurotoxicity seen in the 250 ppm group included: ataxia, spasms, diarrhea, lethargy, and prostration. Colored nasal discharge was reported for all groups (including 40% incidence in the control), but was not explained. DNA from the 250 ppm group (both sacrifice times) eluted faster (breakage of DNA into smaller pieces) than DNA from the 0 ppm group but at a comparable rate to that for the positive control. While it is clear that inhalation exposure to methyl bromide at 250 ppm resulted in DNA damage in male germ cells, the results for DNA elution rates for the 150 and 75 ppm groups was not consistent. The elution of DNA from the 150 ppm group (1 hour post-treatment sacrifice) was significantly slower (DNA as more intact or cross-linking) than the rate seen with the 0 ppm group (both sacrifice times). In the 75 ppm group, DNA from the one-day post-treatment sacrifice groups, but not DNA from the one-hour post-treatment group, eluted faster

than the DNA from the control. Also, the amount of DNA from the one-day group retained on the filters at the end of the elution was significantly less than that seen with the control. A NOEL cannot be determined at this time pending submission of the following information: protocol and raw data; historical control data; explanation of the time frame per group for inhalation exposures, sacrifices, and alkaline elution runs; and data analysis. This study is considered unacceptable by DPR according to FIFRA guidelines. The U.S. EPA concurred with the study author that the result for the 75 ppm group (one-day) did not represent a treatment-related effect (Hansen, 1994). The result at 250 ppm was considered positive for genotoxic potential to the DNA of testicular cells after inhalation exposure.

III.E.4. *In vitro* and *In vivo* Human Studies

Hallier *et al.* (1993) showed a polymorphism in human blood for glutathione-S-transferase activity and methyl bromide. Of the individuals studied, 75% of them were considered conjugators; that is, there was an apparent enzyme-mediated disappearance of methyl bromide when their erythrocyte cytoplasm was incubated with methyl bromide (99% pure; 5,000 ppm) and glutathione. Individuals whose blood did not show such a reaction were considered non-conjugators. The conjugation reaction was apparently a detoxification mechanism because sister chromatid exchanges in the peripheral lymphocytes of non-conjugators were increased by approximately twofold over untreated control levels. Under identical testing conditions, lymphocytes from conjugators showed little or no increase in sister chromatid exchanges.

In a report on biomonitoring, S-methylcysteine adducts in the blood and sister chromatid exchanges in lymphocytes were measured in methyl bromide soil fumigators (Goergens *et al.*, 1994). Methyl bromide exposure levels and duration of exposure were not provided. The adduct levels in the blood ranged from 23 to 42 nmoles/g protein for 8 soil fumigators and 13-18 nmoles/g protein for 4 controls with no methyl bromide exposure. For 14 soil fumigators, there was an increase of sister chromatid exchange rates in the lymphocytes collected in September (the end of use season) compared to those collected in June (beginning of the season).

In another biomonitoring study, hemoglobin S-methylcysteine levels in 14 methyl bromide workers showed a range of 5 to 35 nmoles/g protein compared to 5-10 nmoles/g protein (estimated from graph in the report) (Iwasaki *et al.*, 1989). The methyl bromide concentration in the work place was less than 2 ppm, the detection limit of the gas detector tube.

In a study on the genotoxicity of methyl bromide in humans, blood and oropharyngeal cells were collected from 32 workers involved in structural fumigation (Calvert *et al.*, 1998a). Oropharyngeal cells were used to indicate recent exposure since the average lifespan for these cells is 14 days. Compared to individuals with no history of methyl bromide exposure (28 referents), samples from workers showed an increased incidence of micronuclei in oropharyngeal cells, and an increased frequency of hypoxanthine-guanine phosphoribosyl transferase gene (*hprt*) mutations in lymphocytes. No consistent differences were observed for the frequencies of kinetochore-negative lymphocyte micronuclei, or kinetochore-positive lymphocyte micronuclei. The limitations of the study included: small sample size, inadequate exposure information, and short duration of exposure (median length was 4 hours). The authors did conclude that the findings provided some evidence of genotoxicity in humans after short-term exposure to methyl bromide.

Table 12. The genotoxicity of methyl bromide.

Test types	Route/ Exposure Duration ^a	Dose ^b	Effects/Comments	References ^c
I. Gene Mutation				
<u>Bacterial mutagenicity tests</u>				
S. typhimurium, TA100	air, x 21 h	\$ 0.01%	+, dose-related increase in revertants	1
S. typhimurium, TA100	air, x 5 d	\$ 1900 mg/m ³	+, ± rat liver S9 fraction	2
S. typhimurium, TA100	air	\$ 500 mg/m ³	+	3
S. typhimurium, TA100	air	\$ 0.004 moles/L	+, ± rat or hamster liver S9 fraction	4
S. typhimurium, TA98	air	> 50,000 mg/m ³	!	2
S. typhimurium, TA98	air	> 2.4 moles/L	!	4
S. typhimurium, TA1535	air, x 2 d	\$ 5000 ug/plate	+	3
S. typhimurium, TA1537, TA1538,TA98	air	> 5000 ug/plate	!	3
E. coli Sd-4	solution	\$ 4 mM	+	5
E. coli WP2 hcr	air, x 2 d	5000 ug/plate	+	3
<u>Mitotic recombination</u>				
S. cerevisiae	solution, x 4 d	\$ 0.2%	+, ± rat liver S9 fraction	6*
<u>Sex-linked recessive lethal test</u>				
D. melanogaster	air, x 5 h	> 70 ppm	!	7
D. melanogaster	air, x 6 h/d x 5 or 15 d	\$ 200 mg/m ³	+	2
<u>Forward mutation test (Mouse lymphoma L5178Y)</u>				
TK locus and HGPRT locus	solution, x 24 h	0.3 mg/L	+	2
II. Structural Chromosomal Aberrations				
<u>Micronucleus test</u>				
Mouse	intraperitoneal	> 123 mg/kg	!, in bone marrow cells	8*
Mouse	inhalation, 6h/dx5d/wx2w	\$ 100 ppm	+, in peripheral red blood cells (females)	4*
Mouse	inhalation, 6h/dx5d/wx12w	> 120 ppm	!, in peripheral red blood cells	4*

a/ The duration of exposure was: h= hrs, d=days, and w=weeks.

b/ Dose was the concentration of methyl bromide which resulted in a positive response or the highest dose tested with a negative response.

c/ * indicates the study was acceptable to DPR according to FIFRA guidelines. References: 1. Simmon *et al.*, 1977; 2. Kramers *et al.*, 1985; 3. Moriya *et al.*, 1983; 4. NTP, 1992; 5. Djalali-Behzad *et al.*, 1981; 6. Mortelmans and Shepherd, 1980; 7. McGregor, 1981; 8. Putman and Morris,1991.

Table 12. The genotoxicity of methyl bromide (continued).

Test types	Route/ Exposure Duration ^a	Dose ^b	Effects/Comments	References ^c
<u>Chromosomal aberration test</u>				
Rat	inhalation, 7 h/dx5d	> 70 ppm	! , in bone marrow cells	1
<u>Dominant lethal test</u>				
Rat	inhalation, 7 h/dx5d	> 70 ppm	! , no genotoxicity or reproductive effects	1
III. Other Genotoxic Effects				
<u>Unscheduled DNA synthesis</u>				
Rat hepatocytes	solution, x 24 h	> 0.3 mM	! , no increase in nuclear grain counts	2
Human embryonic intestinal cell	air, x 3 h	> 70%	! , no increase in nuclear grain counts	1
<u>Sister chromatid exchange</u>				
CHO cells	air, x 18 h	\$ 1 ppm	+, dose related 8 in frequency	3
Mouse	inhalation, 6h/dx5d/wx2w	\$ 100 ppm	+, dose related 8 in frequency (females)	4*
Mouse	inhalation, 6h/dx5d/wx12w	> 120 ppm	!	4*
Human lymphocytes	air, x 1hr	5,000 ppm	+, non-conjugators	5
<u>Alkylation</u>				
Mouse	inhalation x 4 h	36 ppm ^d	+, N-7-methylguanine in liver and spleen DNA and methylated cysteine in hemoglobin and liver protein	6
Mouse	intraperitoneal	417 ug/kg	+, alkylation protein in hemoglobin and liver	6
Rat	inhalation x 6 h or single dose by gavage	131-263 ppm 0.58-0.8 mg/kg	+, methylated guanine in liver, lung, stomach and forestomach	7
<u>Sperm abnormality</u>				
Mouse	inhalation, 7h/dx5d	> 70 ppm	! , no increase in frequency of abnormally shaped sperm	1
<u>Micronuclei and hprt mutation</u>				
Humans	inhalation, not reported	not reported	+ (weak), micronuclei and hprt mutations	8

a/ The duration of exposure was: h= hrs, d=days, and w=weeks.

b/ Dose was the concentration of methyl bromide which resulted in a positive response or the highest dose tested with a negative response.

c/ * after the reference number indicates the study was acceptable to DPR according to FIFRA guidelines. References: 1. McGregor, 1981; 2. Kramers *et al.*, 1985; 3. Rounds, 1980; 4. NTP, 1992; 5. Hallier *et al.*, 1993; 6. Djalali-Behzad *et al.*, 1981; 7. Gansewendt *et al.*, 1991; 8. Calvert *et al.*, 1998a.

d/ Static exposure.

III.F. REPRODUCTIVE TOXICITY

Summary: In a 2-generation reproductive toxicity study in rats by inhalation, methyl bromide reduced the fertility rate of the F₁ parents during the second mating trial. While the body weights of the treated pups at birth showed varied responses, their body weights were significantly lowered during lactation. Brain weight and cerebral cortex width were reduced in the F₁ adults.

III.F.1. Inhalation - Rat

Methyl bromide (99.9% pure) was administered to Sprague Dawley rats of both sexes by whole-body inhalation 6 hours per day and 5 days per week at the nominal concentrations of 0, 3, 30, or 90 ppm (American Biogenics Corp., 1986; Hardisty, 1992; Busey, 1993). Parental animals were exposed for about 40 or 55 days and 90 to 105 days before their first and second matings, respectively, and were exposed for a total of 132-145 days before they were sacrificed. There was no exposure to methyl bromide between gestation day 21 and lactation day 4 in any of the four birthing periods. The pups were not directly exposed to methyl bromide until after weaning (post-natal day 28). Body weights (91-95% of control values) and body weight gain (76% of control values) during the pre-mating periods were significantly decreased only in the F₀ males of the 90 ppm group. For the 30 and 90 ppm F₁ groups, the body weights at pre-mating and during reproduction were consistently lower than those of controls. The absolute brain (wet) weights of the F₀ males, F₁ males, and F₁ females in the 90 ppm groups were significantly ($p \leq 0.05$) decreased by 5%, 6%, and 6%, respectively, compared with controls (Table 13). The brain weight of the F₁ females of the 30 ppm group was also reduced by 5% ($p > 0.05$). Fertility indices (the ratio of the number of pregnancies to the number of copulations) were comparable among the four treatment groups (85-100% of control) for the F_{1a}, F_{1b}, and F_{2a} mating trials. However, in the F_{2b} mating trial, the fertility indices decreased from 90.9 of the controls to 66.7% ($p=0.056$) and 68.2% ($p=0.066$) in the 30 and 90 ppm groups.

At birth, the pup body weights of the treated groups were either higher or not significantly different from controls; the only exception was the lowered body weight of the F_{2a} 90 ppm group (Table 13). During lactation, the F_{1a} and F_{1b} pups of the 30 and 90 ppm groups showed significantly reduced body weights on lactation days 14 to 28. The F_{2a} 90 ppm pup body weights were lower at birth than the controls and remained reduced throughout lactation. The F_{2a} 30 ppm pup body weights showed significant reduction on lactation days 14 to 28. The F_{2b} 30 and 90 ppm pup body weights were decreased, starting as early as 4 days after birth. The reduction in body weight was greater in the F_{2a} and F_{2b} progeny (reduction of 9 to 21% at 90 ppm) compared respectively to the F_{1a} and F_{1b} pups (reduction of 5 to 11% at 90 ppm). Since the pups were not exposed to methyl bromide during the lactation period, except perhaps via the maternal milk, the finding of reduced body weights suggested that growth retardation might be an effect due to the 14 to 15 days of *in utero* exposure.

For the female F_{2b} progeny from the 90 ppm group, the absolute weights of the brain, heart, kidneys, and liver were reduced significantly ($p \leq 0.05$) by 7%, 15%, 18%, and 23% when compared to control values, respectively. The absolute weights of the kidneys, liver, and testes

of the corresponding male progeny were also reduced, though to a lesser degree. The organ to body weight ratios were generally not significantly different from control values.

Histomorphometric data showed a decrease in the width of the cerebral cortex (parameters IIIh and IVb in the sectioning scheme of Rodier and Gramann, 1979) of the F₁ 90 ppm groups (both sexes) (Table 13). Measurements for other parameters also were decreased in the F₁ 90 ppm females (IIh, IVb) or F₁ males (IIIa and IIIc). Since the mid- and low-dose F₁ groups were not examined, no NOAEL was established for these effects. However, the reduced brain (fixed) weights for the F₁ 30 ppm females suggested that the LOAEL for the reduced cerebral-cortex width was 30 ppm. Histomorphometric parameters were not affected in the F₀ 90 ppm adults which suggested that the F₁ effects were the result of the *in utero* exposure of the F₁ animals or via feeding in the milk.

The parental NOAEL was 3 ppm based on reduced fertility. The progeny NOEL was 3 ppm based on the decreased pup body weights and organ weights, reduced F₁ adult brain weight, and reduced cerebral cortex width assumed at 30 ppm. This study was considered marginally acceptable to DPR according to FIFRA guidelines. In the evaluation of this study, U.S. EPA established the NOEL for maternal toxicity at 30 ppm based on reduced body weights. Reduced fertility was considered a treatment-related finding and an indicator of reproductive toxicity (U.S. EPA, 1992a).

III.F.2. Oral - Rat

In a published study to investigate the effects of methyl-bromide fumigated feeds, Crj:CD (SD) rats (24/sex/group) were given either basal diet or fumigated feed (80 ppm, 200 ppm, or 500 ppm total bromine) for 18 weeks for each generation (Kaneda *et al.*, 1993). The methyl bromide concentration was reported as < 20 ppb but no details were given. Using the average consumption rates provided in the report, the exposure in terms of methyl bromide was approximately 200 ng/kg/day using average body weights of 0.35 kg and 0.25 kg for males and females, respectively, and consumption rates of 25 g/week and 20 g/week for males and females, respectively. The only significant effects were reduced food consumption in the 500 ppm total bromide F1 parental females during the weeks 9 and 10 of the pre-mating period and on days 0 to 21 of lactation (87-93% of controls), and lowered body weights throughout the lactation period of 500 ppm total bromide F2 female pups (91-95% of controls). Since the actual methyl bromide concentration in each dose is not known, it is not possible to determine whether the effects were due to bromine or methyl bromide.

Table 13. Body weight changes in rats after inhalation exposure to methyl bromide in a 2-generation reproductive toxicity study^a.

Mean Body Weights (grams) ^b								
Lactation Days	F _{1a} litter				F _{1b} litter			
	0	3	30	90 ppm	0	3	30	90 ppm
0	6.0	6.2**	6.2**	6.0	6.2	6.4**	6.2	6.5**
4	9.5	9.4	9.3	9.3	9.3	9.9**	9.5	9.7*
7	13.5	13.7	13.0*	13.1	13.7	14.9**	14.1	14.3
14	23.2	22.9	21.5**	21.6**	24.1	24.2	22.5*	22.5*
21	37.8	37.7	34.3**	33.8**	39.3	39.4	36.0**	36.4**
28	68.4	66.9	62.1**	61.8**	70.1	69.3	64.1**	66.4

Lactation Days	F _{2a} litter				F _{2b} litter			
	0	3	30	90 ppm	0	3	30	90 ppm
0	5.6	6.1**	5.5	5.4**	6.4	6.7**	6.2	6.2
4	8.1	8.4	7.8	7.4**	10.1	9.9	9.2**	9.2**
7	11.6	12.2*	11.6	10.6**	14.3	14.7	13.4*	13.3*
14	21.9	22.6	20.4**	18.6**	24.1	23.7	19.8**	19.6**
21	35.4	36.2	31.4**	29.1**	40.3	39.8	32.4**	32.0**
28	64.3	64.2	58.6**	53.8**	71.6	70.6	58.4**	58.2**

Mean Absolute Brain (wet) Weight (grams) ^c						
ppm	F0 Adults		F1 Adults		F2 Weanlings	
	males	females	males	females	males	females
0	2.26	2.11	2.16	2.05	1.57	1.53
3	2.25	2.09	2.15	2.04	1.54	1.48
30	2.20	2.11	2.14	1.95	1.51	1.45
90	2.14*	2.07	2.02*	1.93*	1.49	1.42**

Mean Cerebral Cortex Width (mm, mean ± standard deviation) ^d for F1 Adults				
ppm	Males		Females	
	IIIh	IVb	IIIh	IVb
0	1.41±0.130	1.47±0.149	1.38±0.163	1.43±0.092
90	1.30±0.143*	1.42±0.121	1.28±0.115*	1.37±0.086*

a/ Data from American Biogenics Corp., 1986. Fetuses were exposed *in utero* to methyl bromide for 5 days/week during gestation days 0 to 19. Offspring were not placed in the inhalation chambers during the lactation period.

b/ Values were mean body weights for both sexes. Statistical significance levels were * at p # 0.05, and ** at p # 0.01 levels using ANOVA and Scheffe's Multiple comparisons reported by the investigators.

c/ Statistical significance levels were * at p # 0.05, and **at p # 0.01 levels.

d/ Data from Busey, 1993. IIIh and IVb are sections of the brain based on Rodier and Gramann (1979).

III.G. DEVELOPMENTAL TOXICITY

Summary: Methyl bromide caused developmental effects in both rats and rabbits after inhalation exposure. The findings in the fetuses included delayed skull ossification in rats and fused sternbrae, gall bladder agenesis, and other effects in rabbits. Methyl bromide did not cause any significant developmental effects in rats and rabbits after oral exposure.

III.G.1. Inhalation - Rat

Pregnant Wistar rats were exposed to methyl bromide (99.5% pure; nominal concentrations of 0, 20, or 70 ppm) for 7 hours per day from days 1 to 19 of gestation (Sikov *et al.*, 1981). Additional groups received a pre-gestational exposure to methyl bromide (20 and 70 ppm) for 5 days per week for the three weeks before mating. There was no significant maternal toxicity, with and without pre-mating exposure, and the NOEL was greater than 70 ppm. The only developmental effect was an increased incidence of delayed skull ossification in the fetuses of both 70 ppm groups, with the developmental NOEL at 20 ppm. This study was considered acceptable to DPR according to FIFRA guidelines.

III.G.2. Inhalation - Rabbit

Pregnant New Zealand white rabbits (26/group) were exposed to methyl bromide (99.5% pure; nominal concentrations of 0, 20, or 70 ppm) 7 hours per day from days 1 to 24 of gestation (Sikov *et al.*, 1981). Maternal toxicity was observed only in the 70 ppm group. After 3 days of exposure and throughout the study, food consumption of the 70 ppm group was lower than the other groups. After approximately 1 week of exposure, the 70 ppm does showed a decrease in body weight and signs of neurotoxicity (convulsive movements, severe to partial paresis of the hind limb). The first doe died on gestation day 9, and two more does died on gestation day 10. Even though dosing stopped on gestation day 15 for all treatment groups (Hardin *et al.*, 1981), all but one of the 70 ppm does were dead by gestation day 30. The NOEL for maternal neurotoxicity was 20 ppm. No developmental toxicity was observed in the fetuses of the 20 ppm group or those (8 fetuses/1 litter) of the one survivor from the 70 ppm group. Because of the loss of the 70 ppm group and the abbreviated duration of gestational exposure in the 20 ppm group, this study was not a valid developmental toxicity study according to FIFRA guidelines.

In a probe study to determine the maternal toxicity and embryo lethality of methyl bromide, pregnant New Zealand white rabbits (7/group) were exposed to methyl bromide (99.6% pure; nominal concentrations of 0, 10, 30, or 50 ppm in Part I; and 0, 50, 70, or 140 ppm in Part II) 6 hours per day by inhalation on days 7 to 19 of gestation (Breslin *et al.*, 1990a). All animals were sacrificed on gestation day 20 except for the 140 ppm group which was sacrificed on gestation day 17 due to their moribund state. No toxicity was observed in Part I.

In Part II, the does in the 140 ppm group showed early signs of toxicity (lethargy and decreased food consumption) after 8 exposures. With continued exposure, the treatment related effects were: decreased body weights and body weight gains; and neurotoxicity (lethargy, labored breathing, ataxia, right-sided head tilt, reduced sensations in the extremities, dilated pupils, lateral recumbency, loss of placing or righting reflex, and rear leg splay).

Histological examination of the brains from the 140 ppm group showed pathological lesions in all animals (multi-focal areas of inflammation of the meninges overlying most regions of the brain and/or bilaterally symmetrical necrosis or spongiosis of the midbrain dorsolateral to the pyramidal tracts). Fetal examinations were limited to counting the number of implantations and resorptions. In the 70 ppm group, there appeared to be a reduction in litter size in association with an increase in pre-implantation loss. However, the loss was not considered treatment-related since implantation occurred before treatment. No examination of the 140 ppm group was reported. This study was considered supplemental information to DPR.

In the definitive study by the same group of investigators, pregnant New Zealand white rabbits were exposed to methyl bromide (99.6% pure; nominal concentrations of 0, 20, 40, or 80 ppm in Part I; and 0 or 80 ppm in Part II) for 6 hours per day by inhalation from days 7 to 19 of gestation (Breslin *et al.*, 1990b). The Part II experiment was designed to determine if the gall bladder agenesis observed in Part I was associated with a particular male used for artificial insemination. Rabbits (in Part II) designated as naive controls were inseminated with sperm from the suspect male.

Maternal effects were observed only in the 80 ppm group and included: decreased body weight gain (Parts I and II), decreased feces, and neurotoxicity (3 of 26 rabbits in Part I only; lethargy, right-sided head tilt, slight ataxia, and slight lateral recumbency). Neurotoxicity was observed on gestation day 19, the last day of exposure. The body weight gain was reduced in both Part I and II 80 ppm groups, but only the reduction in Part II was statistically significant ($p \neq 0.05$). This reduction in maternal body weight gain in the 80 ppm group in Part II was seen in the presence of reduced fetal weight. The significance of the maternal body weight gain reduction is uncertain because body weight changes in rabbits during pregnancy are more variable than other species (U.S. EPA, 1991). The maternal NOEL was 40 ppm based on neurotoxicity.

Fetal effects were also observed primarily in the 80 ppm groups (Table 14). The fetal effects included omphalocele, hemorrhaging (with or without generalized edema), retro-esophageal right subclavian artery, gall bladder agenesis, fused sternbrae, and decreased fetal body weight (13% in Part II). In the 80 ppm group (Part I), the incidences of gall bladder agenesis and fused sternbrae were significantly ($p \neq 0.05$) different from the controls. The increased incidences of gall bladder agenesis and fused sternbrae were independent of maternal toxicity because these effects were observed in fetuses from both normally behaving and affected (with neurotoxicity) does. The finding of gall bladder agenesis was confirmed in Part II with approximately the same litter incidence (29%) as for Part I (26%). Additionally, gall bladder agenesis was not associated with a particular male since the malformation was not observed in the naive controls (in Part II) which had been inseminated only with sperm from the suspect male. The historical control incidences of gall bladder agenesis are in Appendix B. The distribution of affected fetuses with respect to neurotoxicity in the does is shown in the footnotes of Table 14. The developmental NOEL was 40 ppm based on omphalocele, hemorrhaging, retro-esophageal right subclavian artery, gallbladder agenesis, fused sternbrae and decreased fetal body weight at 80 ppm. This study was considered acceptable to DPR according to FIFRA guidelines.

Table 14. The incidence of fetal effects in rabbits after inhalation exposure to methyl bromide during gestation^a.

Effects ^b	Methyl bromide Concentrations						
	Part I				Part II		
	0	20	40	80ppm	0	0 ^c	80ppm
# Examined:							
fetuses	190	137	143	159	114	102	92
litters	21	15	19	19	16	13	14
<u>Fetal body weight (g)</u>	31.8	32.2	35.0	30.4	36.2	33.8	31.4*
<u>External Effects</u>							
omphalocele	0	0	0	2/2 (11%) ^d	0	0	0
hemorrhage (with or without edema)	0	0	0	2/2 (11%) ^d	0	0	1/1 (7%) ^d
<u>Soft Tissues</u>							
retro-esophageal right subclavian artery	0	0	0	2/2 (11%) ^d	0	0	0
gall bladder agenesis	2/1 (5%)	1/1 (7%)	1/1 (5%)	13/5* ^e (26%) ^d	1/1 (6%)	0	4/4 ^e (29%) ^d
<u>Skeletal Effects</u>							
fused sternebrae	0	0	3/2 (11%)	20/10* ^f (53%) ^d	NA ^g	NA ^g	NA ^g

- a/ Incidence data were expressed as the number of fetuses affected/number of litters affected. Data were from Breslin *et al.* (1990b) with does exposed to methyl bromide 6 hours/day on days 7 to 19 of gestation. Parts I and II were two separate experiments. Statistical significance in comparison to the controls, * (p # 0.05), is indicated after each incidence.
- b/ Omphalocele is the protrusion of intestines through a defect in the abdominal wall at the umbilicus. Hemorrhage is subdermal hematoma with either multiple petechiae or edema. Retro-esophageal right subclavian artery refers to the placement of the artery posterior to the esophagus. Fused sternebrae is the premature fusion of the sternum segments.
- c/ These rabbits were designated as naive controls and were inseminated with sperm from suspect male.
- d/ Percent of litters affected= (affected litters/total litters examined) x 100.
- e/ Of the 13 fetuses with missing gall bladder in Part I, 6 were from 3 does without neurotoxicity and 7 were from 2 does with neurotoxicity. In part II, all 4 affected fetuses were from 4 does without neurotoxicity.
- f/ Of the 20 fetuses with fused sternebrae, 19 were from 9 does without neurotoxicity, and 1 from 1 doe with neurotoxicity.
- g/ NA=skeletal examination was not performed.

III.G.3. Oral - Rat

Pregnant rats (Crj:CD (SD), 23-24 rats/dose) were given methyl bromide (purity 99.5%; 0, 3, 10, or 30 mg/kg) dissolved in corn oil by gavage on gestation days 6-15 and sacrificed on day 20 (Kaneda *et al.*, 1998). No clinical signs were observed. Food consumption and weight gain were reduced in the dams of the 30 mg/kg group. Food consumption was also reduced in the control group given corn oil; this suggested that the effect may be related to large volume of corn oil used (10 mL/kg) or the method of administration. At necropsy, all dams in the 30 mg/kg group showed erosion and thickening of the wall in the nonglandular part of the stomach and adhesions between the stomach and the spleen, liver or diaphragm. In the fetuses from the 30 mg/kg dams, there were increased incidences of microphthalmia in 2 fetuses (two litters, 8% incidence), and having 25 (not 26) presacral vertebrae count in 5 fetuses (two litters, 8% incidence). While neither effect was statistically significant, no cases were observed in the control group. This study was considered supplemental information by DPR.

III.G.4. Oral - Rabbit

Pregnant rabbits (Kbl:JW, 15-18 rabbits/dose) were given methyl bromide (purity 99.5%; 0, 1, 3, or 10 mg/kg) dissolved in corn oil by gavage on gestation days 6-18 and sacrificed on day 27 (Kaneda *et al.*, 1998). No clinical signs were observed. Food consumption and weight gain were reduced in the dam of the 10 mg/kg group. In the fetuses, total litter resorption occurred in 2 does of the 10 mg/kg groups and one control doe; the number of resorptions involved were not indicated. Skeletal malformations involving 2-3 litters in at least one methyl bromide-treated group included: splitting of the nasal/frontal/parietal bones; hemivertebra; fusion of the ribs/sternae; and absence of the metacarpal and phalangeal bones. While the number of fetuses with malformation were higher in the treated groups than the control groups; the increase was neither statistically significant at the litter level nor related to the dose. This study was considered supplemental information by DPR.

III.H. NEUROTOXICITY

Methyl bromide is a known neurotoxicant to both humans and experimental animals. The neurotoxicity of methyl bromide in laboratory animals was described **II.A. CHEMICAL IDENTIFICATION** and in previous sections of the **III. TOXICOLOGY PROFILE**. Delayed neurotoxicity study under the FIFRA guidelines is not currently required.

For neurotoxicity in humans, information was available primarily from accidental poisonings. The following selected reports from residential and occupational exposures were those where methyl bromide levels were available or effects adequately described.

III.H.1. Occupational Exposure

One of the earliest reports on the effects of methyl bromide from occupational exposure was those from a 2-week observation in a repackaging plant (Watrous, 1942). The workers manually filled, sealed, inspected, recovered, and packaged methyl bromide from a tank into glass ampules. It is not known whether these workers had previous exposure to methyl bromide. During the first week of the study, the workers had potentially high exposures to methyl bromide due to handling of ampules by hand, spillage during filling, inadequate hood exhaust, and two reported accidents (broken ampules from a packaged box and broken pipe from the methyl bromide tank). While gross methyl bromide leaks were detected by the flame detector, actual measurements were not recorded. In the second week, the work environment was better controlled in that ampules were held by clamps, spillage was vented, and a larger blower was installed for the hoods. Methyl bromide air concentrations were measured hourly at the breathing level of the workers by the flame detector (detection limit of 35 ppm). The recorded methyl bromide air concentration was generally less than 35 ppm. During this 2-week period, twenty-two cases of skin lesions were reported. These were attributed to direct contact of methyl bromide on the skin (i.e. from spillage) and resulted in blisters and/or dermatitis. Seventeen of these 22 cases also complained of systemic symptoms. Overall, 31 of the 90 workers in the filling and sealing rooms developed systemic effects over the 2-week period. The onset of the symptoms as to time after exposure was noted as "variable...sometimes occurring at work after a few hours exposure, and sometimes being delayed until several hours after..the shift." The most common symptoms were anorexia and nausea with variable onset and duration. Headaches were reported to occur only during exposure and were often elicited only after specific questioning of the workers. Vertigo and faintness were reported following "known exposure." The relationship between methyl bromide exposure and other symptoms such as sensations in the eyes, sleepiness, spigastric or substernal pain, and muscular pain were considered questionable.

An investigation was initiated in a date packing plant when 15-20 more workers were absent within a period of 2 to 10 days (Johnstone, 1945). Neurotoxicity was reported in 34 workers exposed to methyl bromide between 100 to 500 ppm for an unknown amount of time while working in the plant. Visual disturbance was present in every case and other symptoms were speech disturbance (23/34), numbness of the extremities (18/34), mental confusion (15/34), hallucination (7/34), melancholia (2/34), coma (4/34), convulsion (1/34), apathy (1/34), tremor (4/34), neurosis (3/34), and fainting attacks (3/34).

Neurobehavioral functions were evaluated in 128 California structural and soil fumigators who had been working with methyl bromide, sulfuryl fluoride, or the combination for at least one year (Anger *et al.*, 1986). The following summary of the report describes only the results for the methyl bromide group. Exposure data for the various work activities are summarized below:

<u>Occupation (number of workers)</u>	<u>Estimated Work hours/day</u>	<u>Exposure Mean (range) ppm</u>	<u>Source of Data</u>
soil fumigator (4)	8	2.3 (0-6.2)	NIOSH
soil fumigator (30)	not given	2.6 (0-7.4)	DPR
tarp remover (4)	variable	4.5 (0-8.6)	NIOSH
shoveler (11)	variable	0.8 (0-2.3)	DPR
structural fumigator (10)	1.5	0.8 (0-2.2)	NIOSH

There was a higher prevalence of muscle aching and fatigue, increased threshold for the two-point test for finger sensitivity, and a lower number of facts recalled in the Wechsler Memory Scale for the methyl bromide group. This group also consistently showed lower performance on neurobehavioral test measures. Mild neurologic dysfunctions were observed in some subjects; they included increased tremors, unsteadiness on standing with eyes closed, ataxia, and poor grip strength. The authors offered the following caveats for the interpretation of the results: (1) lack of information on participation rates and bias; (2) group differences in age, educational level, race, alcohol consumption, use of prescription drugs, and use of "illegal drugs"; and, (3) the possibility of over-reporting of the symptoms.

In an acute exposure, four nursery workers were exposed to methyl bromide while untarping a field (6 acres) which had been treated with methyl bromide (350 lbs/acre) and covered for 10 days (Herzstein and Cullen, 1990). None of the workers had a history of exposure to methyl bromide or had used pesticides during the 6-week period before the incident. During the days of untarping, the workers experienced fatigue and light-headedness. After work, they developed severe coughing, chest tightness, nausea, vomiting, frontal headaches, tremulousness, ataxia, and tremor. The symptoms were less severe over the next 2 to 3 weeks. By 3 weeks, two of the workers reported upper- and lower-extremity paresthesia and reduced hand dexterity. Clinical tests conducted after exposure and follow-up visits were within normal limits. No long-term adverse health effects were reported after 18 months.

In the Netherlands, 9 greenhouse workers were exposed to methyl bromide from an adjacent fumigated area via poor seals around a door and open pipes (Hustinx *et al.*, 1993). Methyl bromide used (200 g/m²) was 5 times the legal use. Some workers were previously exposed to methyl bromide and had experienced symptoms (nausea, vomiting, dizziness, and poor memory). On the first day of fumigation, the methyl bromide level was 25 ppm in the non-fumigated side. On the second day when the workers were poisoned, the methyl bromide levels ranged from 150 to 200 ppm. All, except one worker, experienced extreme nausea, repeated vomiting, and dizziness. The other one felt only a burning sensation in the throat. Two workers later developed seizures. Others complained of headache, nausea, ataxia, slurred speech, and a sensation described as "floating." The serum bromide levels ranged from 51 to 363 mg/L and were higher than the general population. Most of the levels (5-119 mg/L)

remained elevated 19 days after exposure. The severity of the symptoms did not correlate with the bromide levels, but was associated with known previous exposures to methyl bromide.

Two workers were exposed to a high concentration (4400 ppm) of methyl bromide for 45 minutes when aerating a fumigated mill (Deschamps and Turpin, 1996; Garnier *et al.*, 1996). Both experienced nausea, vomiting, headache and dizziness after exposure. Two hours later, one (44 years old) of them had severe myoclonic seizures. This worker was hospitalized and still had ataxia, debilitating action/intention myoclonus, bilateral cortical deafness, and mental deterioration upon discharge after 52 days in the hospital. One year later, the worker remained affected and was confined to a wheel chair. The second worker (39 years old) was less affected and did not develop seizures or myoclonus. One year after the accident, he showed only a mild deficit in verbal memory. Analysis of the blood collected 7 weeks after the accident showed that the difference in response between these two workers may be due to a difference in glutathione transferase level. The more severe neurotoxicity experienced by the older worker was attributed to two factors. First, glutathione transferase activity was higher in this worker (Garnier *et al.*, 1996). The formation of S-methyl-glutathione was hypothesized to be involved in the neurotoxicity observed. On the other hand, the second worker had higher levels of S-methylcysteine adducts in the erythrocyte proteins. Second, the filter mask of this individual was considered "inefficient" which resulted in a 3-fold higher blood bromide level than that for the younger worker.

In a cross-sectional study conducted in 1992 and 1993 by the National Institute for Occupational Safety and Health (NIOSH) and the University of Miami, potential chronic health effects of 123 structural fumigation workers in South Florida were evaluated (Calvert *et al.*, 1998b). A majority of the workers (112) were exposed to methyl bromide and sulfuryl fluoride, with the remaining workers exposed to sulfuryl fluoride only. The median year for employment in the structural fumigation was 4 years (range 0.5 to 32 years). The medians for years worked with methyl bromide and sulfuryl fluoride were 1.2 years (range 0-22.1 years) and 2.85 years (0.11 to 20.5 years), respectively. The tests for neurological function included those for: nerve conduction, vibration, neurobehavioral tests from the Neurobehavioral Evaluation System, vocabulary, Santa Ana Dexterity, posture, contrast sensitivity, color vision, and olfactory function. In addition, urine analysis, lung function, and physical examination were conducted. Pattern memory and olfactory function were the only tests which showed significant difference between workers with high sulfuryl fluoride exposures and the referents. Reduced performance in the Santa Ana Dexterity test by the workers was attributed to the physical damage due to the use of heavy-duty spring clamps to fasten tarps. The authors found few health effects associated with methyl bromide but noted that the study had limited power to assess the exposure.

III.H.2. Residential Exposure

A woman was exposed to methyl bromide after returning to her home which had been fumigated and cleared for reentry (Reidy *et al.*, 1994). Initially, she had trouble breathing and developed headaches, diarrhea, continued nausea, and rashes over uncovered portions of the body for 2 days. The headache continued for several months after the initial exposure. Air sampling done 9 weeks after reentry showed the highest measurements were 3 ppm in the air above the bathroom carpet pad, and 2 ppm in the air above the front door. It was determined

that an excessive amount of methyl bromide was used during fumigation. The actual peak exposure on the first day of reentry was estimated to be 400 to 1500 ppm. The woman continued to live in the home for 14 weeks after the fumigation. She showed impairments in concentration, information processing, learning, and memory as well as emotional stress in a comprehensive neuro-psychological evaluation administered a few days after she moved out of the home.

In another fumigation, a family (a couple with a 3-1/2 month old infant) was exposed to methyl bromide from a neighboring house through emptied sewage pipes (Langard *et al.*, 1996). The estimated air concentration was 6430 ppm in the fumigated house. After about 2.5 hours of exposure, the infant cried vigorously, vomited, and had severe diarrhea. The symptoms persisted until the infant died on the next day. The autopsy showed inflammation in the lungs, blood vessels, heart, and brain. The parents were also affected as they sensed burning in their eyes, throat, and mouth and vomited. However, they recovered with no apparent neurological deficits when tested about 1 week later. The bromide levels were 170 mg/L (36 hours after exposure) for the infant, and 110-130 mg/L (39 hours) for the parents.

Another fatal case of exposure through an open connection was reported in California where a fumigated room was connected to an unattached guest house via 8 uncapped electrical conduits (Swenson *et al.*, 1997). The occupant of the guest house initially complained of flu-like symptoms and was found unconscious and convulsing on the next day. Blood and urine samples collected at the emergency room showed serum bromide levels of 27 mg/dL and 6.2 mg/dL, respectively. Air analysis of one adjoining tube on the 6th day after fumigation showed 15 ppm methyl bromide.

IV. RISK ASSESSMENT

IV.A. HAZARD IDENTIFICATION

The most appropriate data for the hazard identification of methyl bromide are those from human studies. However, human case reports (**III.H. NEUROTOXICITY**) did not provide sufficient detail on the dose-response relationship. In the absence of human data, results from animal studies were extrapolated to humans assuming that the effects observed in laboratory animals would also be observed in humans. Toxicity endpoints and critical NOELs for risk characterization are discussed in this section (details of the studies are in **III. TOXICOLOGY PROFILE**). The no-effect levels may be expressed in terms of NOAELs or NOELs. Only those endpoints considered of toxicological significance were used for risk characterization.

The study to evaluate the chronic inhalation toxicity of methyl bromide in a nonrodent species (*i.e.* dogs), as part of the SB 950 data requirement, has not been submitted to DPR. While a chronic inhalation study in the dog would further characterize the neurotoxicity of methyl bromide, the requirement was waived by DPR based on the evaluation of short-term studies in the dog (Newton, 1994 a and b) which showed that a chronic study would have to be conducted at relatively low dose levels (Oshima, 1995; Gee, 1995). These levels would be in the range of the estimated NOELs from the rat (Reuzel *et al.*, 1987 and 1991) and mouse oncogenicity studies (NTP, 1992; Eustis, 1992). The chronic inhalation study would be required by DPR if the basis for the waiver is changed.

IV.A.1. Selection of Toxicity Endpoints

The selection of a toxicity endpoint and its associated NOEL involved the review of the published literature and registrant-submitted studies for the most appropriate endpoint to assess human exposure.

IV.A.1.a. Developmental Toxicity

One endpoint DPR considered for risk assessment is the development toxicity observed in experimental animals after methyl bromide exposure. In a developmental toxicity study, the pregnant animals are exposed continuously to methyl bromide during a specified period of gestation (when organ formation occurs). Any adverse effect observed in the fetus is considered an acute effect under the current assumption that only a single exposure at a critical time is necessary for the induction of developmental adverse effects according to the U.S. Environmental Protection Agency Guidelines for Developmental Toxicity Risk Assessment. Since this endpoint is the result of exposure during pregnancy, it is only used for the assessment of exposure by women of childbearing age.

The critical NOEL for developmental toxicity was 40 ppm from a study with rabbits. The result from this study was the basis for the emergency regulation and permit conditions currently used in California. U.S. EPA also considered these endpoints of concern and has used the same study in a Section 18 evaluation on the use of methyl bromide on imported fruits at ports of entry. In this rabbit developmental toxicity study, fetuses exposed to 80 ppm *in utero*

showed gall bladder agenesis (no gall bladders), fused sternbrae (early fusion of the sternbrae), and lowered body weights. The missing gallbladder finding was seen in Part I of the experiment, which by itself is a complete study and fulfilled FIFRA guidelines for an acceptable study. The investigator was concerned with the finding as it was rarely observed in the negative-control litters in the conducting laboratory as well as in other laboratories using the same rabbit strain. When the experiment (Part II) was repeated three months later, missing gall bladders were again observed in fetuses exposed to methyl bromide *in utero*. The fused sternbrae found in Part I was not confirmed since skeletal examination was not performed in Part II.

The developmental toxicity effects observed in fetuses should not be discounted because of maternal toxicity (body weight changes and neurotoxicity) reported at the same dose level. Consideration must be given to when the effects were observed. First, the decrease in the body weight gain of the 80 ppm group does was not a consistent finding. Statistically significant decreases were reported for gestation days 13-16 in Part I and gestation days 7-20 and 10-13 periods in Part II. The reduced weight gain in the does of Part II occurred concomitantly with a reduction in the mean fetal body weight. Second, there was no significant difference in the strictly maternal parameter calculated as the terminal body weight minus gravid uterine weight. Third, body weight changes in pregnant rabbits are known to be more variable than rodents. As a result, body weight changes in rabbits often do not carry as much weight as an indicator for maternal toxicity as for rodents as discussed in the U.S. EPA Developmental Toxicity Risk Assessment guidelines. Fourth, maternal neurotoxicity was characterized by clinical signs; including lethargy, head tilt, slight ataxia and slight lateral recumbency. These signs were observed in only 3 of 43 does (7%) dosed at 80 ppm and they did not appear until gestation days 19-20 (the last days of the 13-day exposure period). Based on the description and comparison with observations reported in other studies, DPR did not consider these signs as indicators of excessive toxicity.

Furthermore, the failure of gall bladders to form in some fetuses was independent of maternal neurotoxicity. In Part I, 6 of the fetuses with missing gallbladders were from 3 does without neurotoxicity while the remaining 7 affected fetuses were from 2 does with neurotoxicity. In Part II, none of the does showed neurotoxicity while 4 fetuses (from 4 does) had missing gallbladders. In addition, the development of the gall bladder in rabbits can be considered an acute event since it takes place in one to two days after its onset on gestation day 11.5 (Hoar and Monie, 1981). The maternal neurotoxicity reported on gestation days 19-20 would have occurred too late to have been a factor in the agenesis of the gall bladder.

Similar findings have not been reported in the rat developmental toxicity studies. While it is worth noting that rats do not have gall bladders, the absence of these findings in another species should not negate their significance as indicators of the potential for methyl bromide to cause developmental toxicity in humans. Species specificity in developmental effects has been demonstrated for some chemicals. Developmental toxicity testing under the FIFRA guidelines requires two species to be tested, a rodent and a non-rodent species, typically the rabbit, for the purpose of identifying species susceptibility. The need to test non-rodent species arose from the findings of thalidomide where it was demonstrated that this human teratogen did not exhibit significant teratological effects in rats but caused at least some significant effects in rabbits (Schardein, 1985). As stated in the U.S. EPA Developmental Toxicity Risk Assessment

guidelines, developmental effects may not be evident in more than one species. The findings from the most sensitive species are appropriate to use to estimate human risk.

The significance of the developmental toxicity findings was discussed in a 1994 Proposition 65 meeting to determine whether methyl bromide should be listed for all uses. The emergency regulation in 1992 resulted in methyl bromide being listed as a chemical known to the state to be a reproductive toxicant for structural fumigation use only. The Developmental and Reproductive Toxicity Identification (DART) Committee was presented with results from animal developmental toxicity (absence of gall bladders and fused sternbrae) and reproductive toxicity (decreased pup body weight) studies. After much discussion, the Committee voted not to expand the listing of methyl bromide from structural fumigation to all uses because there was not enough evidence to support the "clearly shown" criteria as mandated by the Proposition. However, the members expressed several concerns: the need for more experimental studies to clarify the findings, potential for exposure to methyl bromide via the milk during lactation, and the lack of information on human exposure especially during pregnancy.

After this meeting, DPR received additional data to support the consideration of reproductive or developmental toxicity as a pertinent endpoint for risk assessment and regulatory actions. First, supplemental data on the rat reproductive toxicity study showed that methyl bromide caused a reduction in the width of a certain part of the brain (cerebral cortex) in the F₁ adults exposed to methyl bromide *in utero* (American Biogenics Corp., 1986). Second, a study received by DPR in 1998 showed that methyl bromide caused a breakage of DNA in the testicular cells isolated from rats after inhalation exposure (Bentley, 1994). It is not known whether the effect was due to methyl bromide or a metabolite.

IV.A.1.b. Neurotoxicity

Methyl bromide caused neurotoxicity in all animal species studied. There are four aspects to the methyl bromide-induced neurotoxicity: (1) species sensitivity, (2) exposure duration dependency, (3) persistency of effects, and (4) cumulative toxicity.

Of the laboratory animals studied, there was a species sensitivity to the neurotoxicity of methyl bromide after short-term exposure. Based on the comparisons of the lowest-observed effect level (LOEL) for neurotoxicity, the dog and rabbit showed greater sensitivity than the guinea pig, mouse and rat. For example, dogs exposed to 156 ppm (human equivalent level of 68 ppm) showed severe neurological effects in 2 to 7 days of exposure while rats exposed to the same concentration in terms of human equivalent level (65 ppm; 70 ppm actual air concentration) for the same exposure duration did not show any neurotoxicity. In pregnant animals, the rabbit was more sensitive to methyl bromide than the rat. For pregnant rabbits, severe neurotoxicity was observed at the LOEL of 70 ppm (Sikov *et al.*, 1981; Breslin *et al.*, 1990) while no neurotoxicity was reported in the pregnant rats at the same level (Sikov *et al.*, 1981).

Although the dog inhalation toxicity studies were not designed to be one of the FIFRA guideline study types, they were conducted under Good Laboratory Practices and DPR considered the results valid for hazard identification. These same data were used by the MBIP to support their position that a chronic inhalation toxicity study in the dog should not be required

(CMA, 1994). The selection of results from the most sensitive species based on the review of the database, in this case the dog, is consistent with the U.S. EPA Neurotoxicity Risk Assessment guidelines (U.S. EPA, 1998a).

Methyl bromide-induced neurotoxicity was dependent on the dose and duration of exposure. As shown in Tables 1 and 3, there was a relatively steep dose-response relationship. The data which best illustrate this point are the results from the dog studies (Newton, 1994 a and b). No effects were observed at 103 ppm (NOEL) for up to 5 days of exposure. However, lacrimation was noted on the first day when the concentration was increased to 156 ppm (1.5 times the NOEL), and the dogs had to be sacrificed after day 6 due to neurotoxic effects. At 314 ppm (3 times the NOEL), tremors and hunched appearance were observed in both dogs after only 7 hours of exposure.

Another aspect of the methyl bromide-induced neurotoxicity is the persistence of the effect after the termination of exposure. In the NTP study, mice were treated with methyl bromide at 100 ppm (NTP, 1992). The exposure was stopped after 20 weeks because of neurotoxicity (tremors, paralysis, and other signs) and mortality. However, neurotoxicity continued to be observed in the survivors for the remainder of the 2-year study. In some cases, the first signs of neurotoxicity appeared one month or more after the last exposure to methyl bromide. Also, at the terminal sacrifice (84 weeks after the last exposure), brain lesions were found in some 100 ppm survivors suggesting that these lesions, which may have been induced during exposure, were not repaired. In rabbits, the neurotoxicity observed was shown to be associated with lesions in the midbrain and meninges (Breslin *et al.*, 1990a). Another example of persistence of effect is the neurotoxicity observed in dogs (Newton, 1994b) and in rabbits after the last exposure to methyl bromide (Sikov *et al.*, 1981).

The persistence of effect may be due to cumulative toxicity after repeated exposure to low doses. In dogs, a comparison of the clinical signs showed that the 158 ppm group was more affected than the 156 ppm by methyl bromide exposure (Newton, 1994 a and b). Decreased activity was noted in the second day (<14 total hours; hourly observations not available) of exposure to 158 ppm compared to the third day (17 total hours) for 156 ppm. The 158 ppm group was previously exposed to 11 ppm for 24 days and appeared normal during the exposure. In a study of California structural and soil fumigators working at air concentration of less than 5 ppm methyl bromide for at least one year, the workers showed lower performance on neurobehavioral test measures and some also showed mild neurological dysfunctions (Anger *et al.*, 1986).

IV.A.1.c. Brain Monoamines and Enzyme Activity

Methyl bromide was shown to decrease tyrosine hydroxylase activity in the rat brain after acute exposure (Honma *et al.*, 1991). This decreased activity was hypothesized by the authors to be due to methyl bromide-induced structural changes to the enzyme. This endpoint and a NOEL of 16 ppm were used by the ATSDR to establish the minimum risk levels for methyl bromide (ATSDR, 1996; ATSDR, 1992). The intent of the MRL was to raise concerns for additional studies rather than for regulatory action.

DPR determined that the LOELs were 16 ppm and 63 ppm based on *in vitro* and *in vivo* assay methods, respectively, for reduced tyrosine hydroxylase activity in the brain segments. It was not possible to determine which LOEL was valid since the publication was incomplete in explaining how important parts of the study were conducted.

Furthermore, the proposed hypothesis was not well substantiated by the investigators. First, there were other plausible explanations for the decreased tyrosine hydroxylase activity in the assays besides direct effects on the enzyme. Second, a decrease in tyrosine hydroxylase activity should result in a decrease in dopamine, the next metabolite in the pathway after DOPA. However, studies done earlier by Dr. Honma's group (Honma *et al.*, 1982 and 1987) either indicated that brain dopamine was not affected or that it was decreased only after levels of its catabolite, homovanillic acid, had increased. Third, the subsequent research in Dr. Honma's group (Honma *et al.*, 1994) did not corroborate or extend the findings of the 1991 paper (Honma *et al.*, 1991). More studies are needed that show the effects on tyrosine hydroxylase and brain dopamine are reproducible. Therefore, the effect of methyl bromide on tyrosine hydroxylase and catecholamines was not selected by DPR as the critical endpoint at this time. While the effect of methyl bromide on tyrosine hydroxylase, by itself, is not considered an adverse effect, the effect on catecholamines in more than one region of the brain (Table 2) was a significant finding for the consideration of critical NOELs for risk characterization.

IV.A.1.d. Nasal Cavity Toxicity

Another important toxicity endpoint is the methyl bromide-induced damage to the olfactory epithelium of rats and mice after inhalation exposure (Reuzel *et al.*, 1987 and 1991; NTP, 1992). This endpoint is used by the U.S. EPA in the determination of the chronic reference inhalation concentration (RfC) (U.S. EPA 1992a) and by the Office of Environmental Health and Hazards Assessment (OEHHA) for chronic toxicity reference exposure levels (RELs) (OEHHA, 1996). With acute exposure to 200 ppm, the damage to the rat olfactory epithelium included epithelial disruption, fragmentation, and exfoliation (Hurtt *et al.*, 1988). Repair of the epithelium included replacement by a squamous epithelium, loss of sensory cells, and respiratory metaplasia (conversion of the olfactory epithelium to a ciliated respiratory type). In other short-term studies, the damage to the nasal epithelium was described as necrosis and degeneration (Eustis *et al.*, 1988) and dysplasia (NTP, 1992; Eustis, 1992). In the chronic inhalation toxicity study, basal cell hyperplasia and degeneration in the olfactory epithelium were observed in the rat (Reuzel *et al.*, 1987 and 1991).

While the effect on the nasal cavity may generally be considered a finding limited to the rat due to anatomical considerations, it is not the case with methyl bromide. Dogs exposed to 156 ppm methyl bromide for only 6 days showed moderate to moderately severe olfactory degeneration (Newton, 1994b). Boorman *et al.* (1990) suggested that the specificity for the toxicity of methyl bromide to this region was due to an abundance of endoplasmic reticula with high metabolic (biotransformation) activity. The greater susceptibility of the olfactory epithelium to pyridine-induced lesions has also been attributed to metabolic activation at this site (Nikula and Lewis, 1994). Air flow to this area, which amounts to 8 to 12% of inspired air (Morris *et al.*, 1993; Kimbell *et al.*, 1993), was considered too slow to be a factor as a target site.

Epithelial hyperplasia of the basal cell layer observed in rats and mice may be a regenerative response or an early indication of neoplasia (Boorman *et al.*, 1990). The olfactory epithelium has remarkable regenerative capacity with a turnover time of 28 days. The basal cells are the stem cells for olfactory neurons. If the basal cells are destroyed, then olfactory epithelium cannot be reconstituted and olfactory function is impaired or lost. The regenerative ability of the basal cells does decline with age (Hastings, 1990). As part of the reparative process, prolonged injury may result in squamous metaplasia and respiratory metaplasia. Squamous metaplasia is characterized by multiple layers of epithelial cells with eosinophilic cytoplasm (Boorman *et al.*, 1990; Haschek and Witschi, 1991). Squamous cell neoplasms have been shown to develop from areas of squamous metaplasia in the olfactory epithelium (Boorman *et al.*, 1990). Respiratory metaplasia, the conversion of olfactory epithelium to a ciliated respiratory type, after exposure to methyl bromide indicates a permanent change in the cell type (Hurt *et al.*, 1988; NTP, 1992; Eustis, 1992). Olfactory epithelial reconstitution after methyl bromide exposure has been used as the model to study the mechanism of recovery of the olfactory system (Schwob *et al.*, 1995).

The nasal effects represented the most sensitive endpoint for chronic exposure. In the rat, lesions in the heart, and decreased brain weights were observed at higher concentrations than that for nasal effects (Table 7). In the mice, the no-effect levels were the same for nasal cavity, brain, sternum, and heart lesions (Table 8).

IV.A.2. Selection of Critical No-Observed-Effect Level

Humans are exposed to methyl bromide by inhalation (occupational, residential, and ambient air) and by oral (dietary) routes. In the evaluation of the potential effects after exposure to methyl bromide, route-specific critical NOELs were derived because the toxicity and pharmacokinetics are different between inhalation and oral exposures. After oral administration, liver and kidneys were the major organs of deposition, and there was reabsorption of biliary metabolites and/or reaction products from the gut. Toxicity was limited to the stomach and forestomach in the rat. For inhalation, the effects were systemic and involved several organs including the nasal cavity, brain, and heart. The primary routes of excretion were via the exhaled air (50% of the dose) as ¹⁴CO₂ for inhalation and intraperitoneal routes, and the urine (43% of the dose) for oral routes of administration.

IV.A.2.a. Acute Toxicity- Inhalation

Studies with animals showed that there is a relatively steep dose-response relationship in which the difference between 100% survival and 100% mortality was only a 2- to 10-fold increase in concentration. The sublethal effects of methyl bromide included neurotoxicity, biochemical alterations, and tissue degeneration (Table 4).

The critical NOEL for acute exposure was selected from neurotoxicity and developmental toxicity studies. For the adult population, especially for women of childbearing age, the lowest NOEL was from the rabbit developmental study by Breslin *et al.* (1990b) since the dosage was lower than that used in the rat study (Sikov *et al.*, 1981). Even though in the developmental study the exposure was repeated during the gestation period, the primary assumption is that only a single exposure at a critical time is necessary for the induction of

developmental adverse effects according to the U.S. EPA Guidelines for Developmental Toxicity Risk Assessment (U.S. EPA, 1991).

In the rabbit developmental study, the NOEL was 40 ppm (human/adult equivalent of 21 ppm) (Breslin *et al.*, 1990b). Fetal malformations (omphalocele, retro-esophageal right subclavian artery, and gall bladder agenesis), variations (fused sternbrae), and decreased fetal body weights were observed in the fetuses of does exposed to methyl bromide at 80 ppm. Since these malformations are rarely seen in litters of untreated rabbits, they were considered treatment-induced in this study. The occurrence of these effects in a single group by chance was unlikely given the historical control data for the conducting laboratory (for more discussion, see Appendix B). The increased incidences of gall bladder agenesis and fused sternbrae were statistically significant ($p \# 0.05$) (Table 14). Furthermore, gall bladder agenesis was confirmed when the experiment was repeated. Skeletal examinations were not done in the second experiment. Both adverse effects were observed in fetuses from does with and without neurotoxicity, indicating that maternal toxicity was unlikely to be the determining factor in the development of the defects.

Developmental effects were also observed in the rat, though to a lesser extent compared to the rabbit. In rats, the fetuses of the two groups exposed to 70 ppm methyl bromide showed increased delayed skull ossification, and the NOEL was 20 ppm (human/adult equivalent NOEL of 22 ppm) (Sikov *et al.*, 1981). No maternal toxicity was observed in this study.

For other population groups, including infants and children, the most appropriate endpoint for risk assessment is neurotoxicity. The lowest NOEL was <16 ppm (human/child equivalent of <11 ppm) for a decrease in tyrosine hydroxylase activity in the rat hypothalamus (Honma *et al.*, 1987 and 1991). This NOEL from the *in vitro* assay was not selected for risk assessment because of the inconsistencies in the findings and interpretation of the results (**IV.A.1.b. Brain Monoamines and Enzyme Activity**). However, the NOEL (31 ppm or human/child equivalent of 22 ppm) for altered catecholamines in more than one region of the brain was considered as a support for the critical NOEL derived from the dog studies.

The critical acute NOELs for neurotoxicity were considered from clinical observations in dogs (Table 3; Newton, 1994 a and b). The selection of the 103 ppm dose (human/child equivalent of 25 ppm) as the acute NOEL considered three major factors: subjectiveness of the observations, severity of the neurotoxicity at higher concentrations, and possibility of delayed neurotoxicity. The NOEL of 103 ppm in the dogs was based on gross observations. Neurotoxicity may be present but not detected unless more refined methods such as the Functional Observation Battery were used. It is possible that the actual NOEL may be lower than 103 ppm. Furthermore, severe neurotoxicity was observed at higher doses (1.5 times the NOEL) with a few additional days of exposure. At 156 ppm, one of two dogs showed lacrimation (tearing) on the first day. This finding by itself may arguably be considered less significant with respect to adversity. However, there were only two dogs in this group. With 2-3 days of additional exposure, both dogs showed significant toxicity manifested as difficulty in breathing and decreased activity. In another study, all dogs (8 in the group) exposed to 158 ppm showed decreased activity before the end of the second exposure day (the first dose was on a Friday and the second dose was on the following Monday). With 5 additional days of

exposure, all showed severe neurotoxicity and brain lesions. The selection of 103 ppm as the acute NOEL also addresses, indirectly, the possibility of delayed neurotoxicity which has been reported in humans after accidental poisonings. Since no effects were observed in the dogs at 103 ppm for 7 days of continuous exposure, it is unlikely that there would be delayed neurotoxicity within one week after a single exposure to the same level. The human equivalent NOEL (25 ppm) from the dog study is only two-fold or less than those for the acute effects observed in the rats and guinea pigs (Table 6).

From the two critical acute NOELs selected to address human population groups, the critical NOEL of 40 ppm (human/adult equivalent of 21 ppm) for developmental toxicity in the rabbit was most appropriate for workers and residents since women of child bearing age are in both groups. For children, the critical NOEL was 103 ppm (human/child equivalent of 25 ppm) for neurotoxicity in the dog (Newton, 1994a). Since the developmental toxicity NOEL was lower than that for neurotoxicity, this NOEL was selected for risk characterization to address occupational and residential exposures and would also be protective of the children population.

IV.A.2.b. Subchronic Toxicity- Inhalation

From occupational and residential uses of methyl bromide, there is a potential for subchronic inhalation toxicity from short-term exposure periods due to a single application, and from multiple applications during a season. One example of potential short-term exposure is from the release of methyl bromide from wall space of fumigated homes. Current label allows residents to reenter treated homes when the methyl bromide concentration reached #1 ppm in the wall space. Residents living next to fumigation chambers or fields with consecutive application may also experience short-term exposures.

For continuous exposure of 1 week, methyl bromide effects on the brain and clinical signs of neurotoxicity were the endpoints used to select the critical NOEL (Table 6). For this duration of exposure, the lowest and the critical NOEL was 20 ppm (human/child equivalent of 7 ppm) for neurotoxicity and death in pregnant rabbits (Sikov *et al.*, 1981). After approximately 1 week (exact number of days not reported), the 70 ppm does showed a decrease in body weight and signs of neurotoxicity (convulsive movements, severe to partial paresis of the hind limb). One of 26 does died on gestation day 9, and two more does died on gestation day 10. Though dosing stopped on gestation day 15, all but one of the 70 ppm does were dead by gestation day 30. Neurotoxicity was also observed in pregnant rabbits in two other studies (Breslin *et al.*, 1990 a and b). The NOELs (70 ppm and 40 ppm) were higher in these latter studies compared to that from Sikov *et al.* (1981). The lower NOEL from the Sikov *et al.* (1981) study may be due to longer exposure as the rabbits were exposed earlier (gestation day 1 compared to gestation day 7) and longer each day (7 hours instead of 6 hours).

The NOEL of 26 ppm (human equivalent/child of 5 ppm) for decreased activity in the dog after 14 days of exposure to 53 ppm was also considered for the short-term critical NOEL. The decreased activity was considered an early sign of neurotoxicity since decreased activity and more severe signs of neurotoxicity were observed in the 103 ppm group during the 1-2 weeks of exposure. However, the time to effect (14 days) was much longer than the duration (7 days) for the scenarios of concern.

For subchronic exposures of longer duration (> 1 month, seasonal), the critical NOEL was an estimated NOEL (ENEL) of 0.5 ppm (human/child equivalent of 0.1 ppm) based on a LOAEL of 5 ppm for decreased responsiveness in two of eight dogs during a neurological examination after about 7 weeks of exposure (34 exposure days). A default uncertainty factor of 10 was used to calculate the NOEL from a LOAEL (Dourson and Stara, 1983). While the duration is shorter than the 13-week generally considered for subchronic exposure, it was chosen because of the endpoint (neurotoxicity) and species sensitivity (the dog is a more sensitive species than the rat to methyl bromide). It is possible that the NOEL might be lower if the dogs were exposed to methyl bromide for 13 weeks.

This ENEL was lower than the NOEL (3 ppm, human/adult equivalent of 2 ppm) for lowered body weights of rat pups from dams exposed to methyl bromide before mating and during part of the pregnancy in the reproductive toxicity study (American Biogenics Corp., 1986) (Table 13). The reduction of pup body weight was both dose- and time- dependent with respect to the adult exposure, with the highest reduction occurring in the F_{2b} 30 ppm and 90 ppm litters. There was also decreased brain weight in the F₁ adults at 30 ppm. This study and the endpoint are appropriate to evaluate the potential effects in humans exposed to methyl bromide during pregnancy. The potential post-natal toxicity was, therefore, accounted for by the use of the lower ENEL for neurotoxicity in the dog. Another study also showed an estimated NOEL of 3 ppm (human/child equivalent of 1.1 ppm) based on a significant decrease in the absolute brain weight in female rats exposed to 30 ppm and higher concentrations of methyl bromide for 13 weeks (Norris *et al.*, 1993 a and b). This effect on the brain weight was considered biologically significant since the brain is a target organ of methyl bromide. The absence of neurotoxicity by Functional Observation Battery testing at the same dose (30 ppm) does not diminish the importance of the brain weight finding since the etiology of the two effects are not necessarily related.

IV.A.2.c. Chronic Toxicity- Inhalation

For chronic inhalation exposure, all chronic studies conducted with rodents (rats and mice), the reproductive toxicity study, and the subchronic dog inhalation toxicity study were considered in the determination of the chronic critical NOEL. While the exposure duration in chronic toxicity studies generally lasted for the life-time for rodents, the actual duration of the testing was two years. Since humans may be exposed to methyl bromide on a yearly basis, not just one or two years in the lifetime, the NOEL from the chronic toxicity study after two years of exposure was, therefore, appropriate for use. This NOEL may underestimate the risk of repeated yearly exposure as there is evidence of cumulative toxicity for this endpoint, as well as for neurotoxicity.

After chronic inhalation exposure, tissue damage was noted in the nasal cavity, brain, and heart of rodents. The most significant finding after chronic exposure was the dose-dependent increase in the incidences of nasal olfactory epithelial hyperplasia and respiration metaplasia found in rodents (Reuzel *et al.*, 1987 and 1991; NTP, 1991; Eustis, 1992; Gotoh *et al.*, 1994). In the Reuzel *et al.* study (1987 and 1991), the incidences of hyperplasia were dependent on the exposure duration as there was an increase in the number of rats affected from 1 year to 2.5 years of exposure. The LOAELs were >90 ppm, 30 ppm, and 3 ppm for exposures lasting 12 months, 12-24 months, and 24-29 months, respectively (Table 7). At 30

ppm, there was also a reduction of absolute brain weight with a NOEL of 3 ppm. Using a standard default uncertainty factor of 10 for the calculation of an estimated NOEL from the LOEL (Dourson and Stara, 1983) for the 24-29 months of exposure, the critical ENEL for basal cell hyperplasia/ degeneration in the rat olfactory epithelium was 0.3 ppm (human/child equivalent of 0.1 ppm).

The LOAEL of 3 ppm from the rat inhalation study by Reuzel *et al.* (1987 and 1991) was supported by the study of Gotoh *et al.* (1994). In the Gotoh *et al.* study, the estimated NOEL was 4 ppm (human/child equivalent of 0.1 ppm) for nasal cavity inflammation (4 ppm males) and respiratory metaplasia (4 ppm females) in rats exposed to methyl bromide by inhalation for 104 weeks. Other long-term studies showed higher NOELs or ENELs than the critical ENEL. In the mouse chronic toxicity study, the LOEL was 10 ppm for neurobehavioral effects and sternal dysplasia (NTP, 1992; Eustis, 1992). Using a default uncertainty factor of 10 for the calculation of a NOEL from a LOEL, the ENEL would be 1 ppm (human/adult equivalent of 0.7 ppm). In the reproductive toxicity study, pregnant rats showed a reduction in fertility indices (American Biogenics Corp., 1986). However, the NOEL of 3 ppm (human equivalent of 2 ppm) was higher than the 0.3 ppm ENEL for olfactory epithelial hyperplasia observed in the rat oncogenicity study (Reuzel *et al.*, 1987 and 1991).

Another comparison of the NOELs was made with the results from the subchronic neurotoxicity study. The ENEL of 0.3 ppm for nasal cavity effects when expressed as human/child equivalent level (0.1 ppm) was the same as the human equivalent level for neurotoxicity after subchronic exposure (ENEL of 0.5 ppm). This implied that the actual NOEL for chronic exposure if based on neurotoxicity could be lower than that based on the effects in the nasal cavity. However, it is not possible to extrapolate such a NOEL at this time because the subchronic ENEL was already an estimated NOEL based on a LOEL which was reduced by a 10-fold uncertainty factor. The use of two 10-fold uncertainty factors to the subchronic LOEL may result in a no-effect level well below the threshold for neurotoxicity after chronic exposure.

IV.A.2.d. Oncogenicity

Positive findings in genotoxicity studies indicate that methyl bromide is potentially oncogenic. Methyl bromide is a direct-acting mutagen. It has been shown to alkylate DNA in different organs in *in vivo* studies (Djalali-Behzad *et al.*, 1981; Gansewendt *et al.*, 1991) and was positive for genotoxicity in several *in vitro* and *in vivo* assays (Table 12). Also, a recent report showed genotoxicity in workers exposed to methyl bromide (Calvert *et al.*, 1998).

However, there was no clear evidence of oncogenicity under the experimental conditions used in the oncogenicity and chronic toxicity studies. First, dose-related responses were found only for non-neoplastic lesions and included nasal epithelial hyperplasia in Wistar rats (Reuzel *et al.*, 1987 and 1991) and B6C3F1 mice (NTP, 1992; Eustis, 1992) after chronic inhalation exposure, and forestomach hyperplasia in Wistar rats after subchronic gavage exposure (Danse *et al.*, 1984; Boorman *et al.*, 1986; Hubbs, 1986). The enlarged spleens observed in Sprague-Dawley rats fed methyl bromide in microcapsules mixed in the feed (Mertens, 1997) can not be considered lymphomas without actual histological data. Second, tumors reported in some studies did not provide sufficient evidence for oncogenicity. The incidences were low and were observed either in the low dose females only (adrenal gland pheochromocytoma in

Fischer rats and lymphoma in BDF1 mice, Gotoh *et al.*, 1994) or were not statistically significant from the control (for example, glioma and granular cell myoblastoma in Wistar rats, Reuzel *et al.*, 1987 and 1991). Therefore, oncogenic risk for methyl bromide was not considered based on the currently available data.

A summary of the critical NOELs for risk characterization is presented in Table 15.

Table 15. The critical no-observed-effects levels (NOELs) for the risk characterization of inhalation exposures to methyl bromide.

Scenarios	Experimental NOEL	Human Equivalent NOEL ^a		Reference Concentration ^d	Effects in Animal Studies	Ref ^e
		Adult ^b	Child ^c			
Acute	40 ppm	21 ppm	na	210 ppb	Developmental toxicity (pregnant rabbit)	1*
	103 ppm ^f	45 ppm	25 ppm		Neurotoxicity (dog)	2
Subchronic 1 week	20 ppm	12 ppm	7 ppm	120 ppb(adult) 70 ppb (child)	Neurotoxicity (pregnant rabbit)	3
	6 weeks	0.5 ppm (ENEL)	0.2 ppm			0.1 ppm
Chronic	0.3 ppm (ENEL)	0.2 ppm	0.1 ppm	2 ppb (adult) 1 ppb (child)	Nasal epithelial hyperplasia/ degeneration (rat)	4*

a/ Experimental NOELs were converted to human equivalents using equations in Appendix G. na= child equivalent NOEL were not calculated because the effects were observed in pregnant animals.

b/ The adult equivalent NOELs are appropriate to address worker exposures. They are also used for residential exposures when child equivalent NOELs were not calculated.

c/ The child equivalent NOELs are appropriate to address resident exposures (see footnote b).

d/ The reference concentration is the ratio of the human equivalent NOEL and a default uncertainty factor of 100 since the NOEL was derived from experimental animal studies.

e/ * indicates study was acceptable to DPR according to FIFRA guidelines. References: 1. Breslin *et al.*, 1990b; 2. Newton, 1994b; 3. Sikov *et al.*, 1981; 4. Reuzel *et al.*, 1987 and 1991.

f/ The NOEL and human equivalents are presented in this Table for comparison purposes only. They are not used for risk characterization.

IV.B. EXPOSURE ASSESSMENT

Human exposure assessments were conducted for occupational and residential inhalation exposures to methyl bromide (Appendices F, H, and I). The exposure levels were expressed as methyl bromide concentration in the air (ppb). The duration of exposure considered included: acute (daily exposure), short-term (referred to as 7 day or subacute in Appendix F), seasonal (90 days), and chronic (annual) exposures. For acute exposure, the highest detected (when available) and the mean values were presented. Non-acute exposure levels (short-term, seasonal, and chronic) were calculated based on acute exposures. Workers in pre-plant soil fumigation and residents in structural (home) fumigation were considered not to have chronic exposures to methyl bromide. Data variability expressed as standard deviation of the mean was calculated only when there were sufficient data points. All the exposure estimates were adjusted for 50% recovery (discussed in Appendices F and K).

Certain scenarios were not included in the Exposure Assessment. The inhalation exposures for workers in structural fumigation were not determined because they are required to wear self-contained breathing apparatus. Published studies showed that dermal exposure was significant at high methyl bromide air concentration. Skin lesions were observed in workers exposed to high concentrations of methyl bromide (9,000 ppm) (Zwaveling *et al.*, 1987; Hezemans-Boer, 1988). However, dermal exposure to methyl bromide was also not assessed because of lack of dermal absorption information for methyl bromide (discussed in Appendix F).

Because of the number of scenarios determined, only a general discussion with a focus on the acute exposure is provided in this section. Also, the location of the type of application discussed in the Tables is indicated in italics. Detailed descriptions are provided in Appendices F, H, and I.

IV.B.1. Occupational Exposures

Occupational exposure levels were determined for the use of methyl bromide in soil fumigation (Table 16) and commodity fumigation (Tables 17 and 18). For each scenario, there was a wide range of exposure levels depending on the job task or the location monitored. For most workers (except for walnut processing and brewery), exposures were measured by personal monitoring devices. A summary of the data is in Table 20.

For the workers involved in the use of methyl bromide for preplant soil fumigation, the range of acute exposures were 0.6 ppb (*16h*. shovelman of shallow shank tarp-bed method) to 1659 ppb (*16a*. tarp remover) (Table 16). Applicators, co-pilots, cutters, pullers, cultipackers, and other workers had exposures between these two levels. While the shovelman of shallow shank non-tarped-bed method had the lowest acute exposure (0.6 ppb), the shovelman for the shallow-shank tarp method (*16a*) had much higher exposures (high of 515 ppb and a mean of 147 ppb). Tarp removers of the latter method (*16a*) also had the highest exposures for all durations: 1659 ppb (mean of 835 ppb) for acute, 596 ppb for short-term, and 510 ppb for seasonal exposures.

For nursery potting soil workers, the use of the hot-gas method (*16l*) resulted in high exposure levels (Table 16). The highest exposure was 724 ppb measured for the applicators of

this method compared to 26-31 ppb for applicators not using such a method (16k). The range of acute exposures were 1 (tarp remover of hot gas method) to 724 ppb (applicators of hot gas method). No seasonal or chronic exposures were determined.

For workers involved in the fumigation of grain products, the range of acute exposures was 8 ppb (forklift driver of chamber) to 8458 ppb (aerator of sea container/trailer) (Table 17a). These aerators also had the highest exposures for short-term (4314 ppb), seasonal (3271 ppb), and chronic (2482 ppb) exposures. However in comparison, aerators of tarp fumigation had much lower (less than one-tenth) exposures for all durations. Workers in the rice warehouse had relatively low exposures at 10 ppb and 7 ppb for acute and short-term exposures, respectively.

For workers involved in the fumigation of dried fruit and tree nut products, the range of acute exposures was 0.35 ppb (17b2. forklift operator of dried prunes in chamber) to 1463 ppb (17b4. worker involved in clear chamber for raisins) (Table 17). For dried fruits, the fumigators and aerators had higher exposures than other workers. Around non-certified chambers for walnuts, workers not directly involved in the application of methyl bromide had higher exposures than that (225 ppb) for the fumigator (17c). The acute exposures of these workers ranged from 770 ppb (sorting worker) to 1789 ppb (cracking worker).

For workers involved in the fumigation of cherries for export, the range of acute exposures was 47 ppb (forklift driver) to 877 ppb (fumigator, start-up) (Table 17d). As with dried fruit fumigation, the fumigator had the highest exposures.

For workers in a walnut processing plant (17e), potential exposures were based on air sampling in work stations rather than specific tasks (Appendix F). Cleaning plant (448 ppb), vacuum chamber (636 ppb), and Dock 5 (500 ppb) had the highest methyl bromide levels for acute exposure compared to other areas (Table 17). The lowest mean level (39 ppb) was measured in the sorting area.

For workers in a brewery, exposures were also based on air concentrations measured at various locations (Table 17f). The exposure levels ranged from 28.9 to 173 ppb.

For a rice warehouse, the highest ambient air concentration was 55 ppm (mean of 32 ppb) for acute exposure (Table 18a). For other commodity fumigation facilities, high methyl bromide levels were generally detected near the chambers (Table 18). For example, 13281 ppb were detected in a leak check in a dried prune chamber (18c), 2175 ppb in raisins aeration sheds (18d), 1789 ppb in walnut vacuum chamber area (18f), and 6800 ppb by the injection site in a walnut processing facility (18i2). Other areas in these facilities had various exposure levels which probably depended on the distance from the source of methyl bromide, ventilation, and other factors. Each facility had areas with < 10 ppb methyl bromide.

IV.B.2. Residential Exposures

Residential exposures were determined for several exposure scenarios (Table 19). The ranges of acute exposure levels were: 406 to 2990 ppb for residents near houses during fumigation (19a), 74 to 1064 ppb for residents downwind from a fumigated house during

aeration (19b), 210 ppb for residents after reentry into treated homes (19c), and 210 ppb for residents living near commodity fumigation facilities (19d). The highest exposure level was 2990 ppb measured for outdoor air 20 feet from a house during fumigation (19a). Measurable levels of methyl bromide were detected in the indoor air of neighboring houses: 406 ppb (during fumigation) and 168 ppb (during aeration). Lower levels were detected during the aeration process and declined with distance from the fumigated house. The levels were 1064 ppb at 10 feet, 208 ppb at 50 feet, and 74 ppb at 100 feet. The acute exposure for residents reentering fumigated homes and those living near commodity fumigation facilities was 210 ppb, the current regulatory limit in the DPR permit conditions.

The exposure level for residents living near fumigated fields was 210 ppb (Table 19e) which is the regulatory level used to determine the buffer zone distance in the DPR permit conditions (Appendices H and I). While there may be a potential for longer duration of exposure, a quantitative determination of the exposure cannot be made at this time (Appendix J).

Table 16. Estimates of occupational exposures to methyl bromide in preplant soil injection^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
a. Shallow-shank tarp method (Appendix F Tables 13 to 17)					
applicator	303	111 ± 98	95 ± 84	49 ± 44	n/a
co-pilot	518	224 ± 152	192 ± 130	100 ± 68	n/a
shovelman	515	147 ± 135	63 ± 58	n/a	n/a
tarp remover with PCO	1659	835 ± 596	596 ± 426	510 ± 364	n/a
tarp remover with grower	553	278 ± 199	79 ± 57	n/a	n/a
b. Deep shank injection fumigation (Appendix F Table 18)					
applicators	181	154	132	68	n/a
copilot	n/a	49	42	22	n/a
cultipacker	n/a	99	85	n/a	n/a
c. Improved deep shank injection fumigation (Appendix F Table 18)					
applicator	n/a	57	49	25	n/a
cultipacker	n/a	70	60	n/a	n/a
d. Deep shank injection fumigation (Appendix F Table 19)					
1. basic, second tractor					
applicator	n/a	88	75	39	n/a
disc driver	n/a	512	439	228	n/a
2. basic, cultipacker					
applicator	165	94	81	42	n/a
supervisor	n/a	67	57	30	n/a
cultipacker	58	34	29	n/a	n/a
e. Deep shank injection fumigation (Appendix F Table 20)					
applicator	n/a	7	6	3	n/a
cultipacker	n/a	7	6	n/a	n/a

Table 16. Estimates of occupational exposures to methyl bromide in preplant soil injection (continued)^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
f. Shallow shank tarped-bed fumigation (Appendix F Table 21)					
1. raised platform applicator	n/a	80	69	36	n/a
co-pilot	111	104	89	46	n/a
2. closing shoes applicator	n/a	44	38	20	n/a
co-pilot	209	167	143	74	n/a
g. Shallow shank tarped-bed fumigation (Appendix F Table 22)					
1. tractor fumigation driver	n/a	28	24	12	n/a
applicator	n/a	45	39	20	n/a
drip tape layer	n/a	65	28	n/a	n/a
2. tractor with tarp driver	n/a	4	3	1.8	n/a
co-pilot	65	34	29	15	n/a
h. Shallow shank tarped-bed fumigation (Appendix F Table 23)					
applicator	n/a	3	3	1	n/a
co-pilot	31	31	27	14	n/a
shovelman	0.6	0.6	0.3	n/a	n/a
pipe layer	2	2 ±0	1	n/a	n/a
pipe tractor driver	918	411 ±477	176±204	n/a	n/a
i. Tarp shallow with Noble plow shanks (Appendix F Table 24)					
cutter (grower's)	237	82±134	23±38	n/a	n/a
cutter (company)	237	82±134	59±96	27±45	n/a
puller (grower's)	324	33± 92	9 ±26	n/a	n/a
puller (company)	324	33± 92	24±66	11±31	n/a

Table 16. Estimates of occupational exposures to methyl bromide in preplant soil injection (continued)^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
j. Tarp shallow with Noble plow shanks (Appendix F Table 25)					
1. PCO's					
cutter	n/a	78	56	26	n/a
remover:tractor driver	n/a	343	245	114	n/a
remover:basketman	n/a	325	232	108	n/a
remover:end puller	n/a	7	5	2	n/a
2. grower's					
cutter	n/a	78	56	n/a	n/a
remover:tractor driver	n/a	343	245	n/a	n/a
remover:basketman	n/a	325	232	n/a	n/a
remover:end puller	n/a	7	5	n/a	n/a
k. Nursery potting soil fumigation (Appendix F Table 26)					
applicator	26	21	3	n/a	n/a
applicator assistant	31	16	2	n/a	n/a
tarp remover	133	94±38	13 ± 5	n/a	n/a
tractor driver	31	16	2	n/a	n/a
truck driver	8	3±4	0.4±0.6	n/a	n/a
potter	95	32±43	5 ±6	n/a	n/a
l. Greenhouse potting soil fumigation-hot gas method (Appendix F Table 27)					
applicator	724	562	80	n/a	n/a
tarp venter	333	93±160	13±23	n/a	n/a
tarp remover	2	1±1	0.1±0.1	n/a	n/a

^{a/} Detailed description of the exposure scenarios are provided in Appendix F. n/a= not applicable or no exposure data available.

Table 17. Estimates of occupational exposures to methyl bromide in commodity fumigation facilities^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
a. Fumigation of grain products (Appendix F Table 28)					
applicator	3340	1652±1462	1180±1044	895±792	697±601
aerator(sea container/trailer)	8458	6039±4062	4314±2901	3271±2201	2482±1669
aerator (tarp)	726	251±411	179±294	136±223	103±169
forklift driver(container/trailer)	43	16±24	11±17	4±6	4±6
forklift driver (chamber)	8	6±2	4±1	2±1	1±0.5
workers (rice warehouse)	10	10±0	7±0	n/a	n/a
b. Fumigation of dried fruit and tree nut products (Appendix F Table 29)					
1. sea van					
fumigator	n/a	33	9	n/a	n/a
fumigator observer	n/a	9	3	n/a	n/a
2. chamber (dried prunes)					
forklift operator	n/a	0.35	0.2	n/a	n/a
fumigator	n/a	938	402	n/a	n/a
3. big chamber (raisins)					
primary fumigator	n/a	800	571	445	311
secondary fumigator	n/a	118	84	65	46
aerator	925	490	350	272	190
forklift driver	69	49±19	35±14	26±10	18±7
catchall operator	n/a	203	145	135	95
hopper operator	227	160	114	107	75
capper dumper	n/a	127	91	85	59
inspector	n/a	15	11	10	7
moisture checker	n/a	9	6	6	4
forklift to side hopper	n/a	8	6	4	3
stem picker	28	22±6	16±4	15±4	10±3
packer	n/a	19	14	13	9
4. chamber (raisins)					
fumigator	107	63	54	44	17
aerator	64	47	40	33	13
clear chamber	1463	1434	1229	1004	393
stem picker	30	28	24	20	12
forklift driver	n/a	3	3	2	0.4
hopper operator	n/a	19	16	13	8

Table 17. Estimates of occupational exposures to methyl bromide in commodity fumigation facilities (continued)^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
c. Fumigation of two non-certified chambers (walnut) (Appendix F Table 29)					
cracking worker	1789	1141±269	978±231	887±209	n/a
sorting worker	770	593±116	508± 99	461± 90	n/a
fumigator	1123	559±446	479±382	316±252	129±103
cleaning-fumigator	n/a	80	69	45	18
cleaning-sort	1089	889±185	762±159	691±144	n/a
bulk casing worker	n/a	856	734	666	n/a
in-shell packer	1011	972	833	756	n/a
hopper operator	n/a	778	667	605	n/a
fumigator	n/a	225	193	127	52
d. Fumigation of cherries for export (Appendix F Table 30)					
fumigator, start-up	877	267±306	191±219	n/a	n/a
fumigator, closing-up	435	327	234	n/a	n/a
fumigator, opening-up	124	58± 46	41±33	n/a	n/a
forklift driver	47	11± 14	8±10	n/a	n/a
sorter	337	123± 72	88±51	n/a	n/a
e. Walnut processing facility (Appendix F Table 31)					
bulk packaging	74	39±28	33±24	33±23	n/a
cleaning plant	448	233±165	200±141	194±138	n/a
Dock 5	n/a	500	429	417	n/a
fumigatorium	106	45±50	39±43	27±31	10±11
packaging	80	62	53	52	n/a
vacuum chamber	636	283±216	243±185	236±180	n/a
sorting	70	39±17	33±15	33±14	n/a
special cracking	170	62±56	53±48	52±47	n/a

Table 17. Estimates of occupational exposures to methyl bromide in commodity fumigation facilities (continued)^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
f. Fumigation and aeration at a brewery facility (Appendix F Table 32)					
1. application					
entry to open canister	n/a	28.9	8.3	n/a	n/a
area sample	n/a	42	12	n/a	n/a
2. aeration					
aerator	25	25±24	7±7	n/a	n/a
area sample (entrance)	n/a	173	49	n/a	n/a
area sample (truck)	n/a	100	29	n/a	n/a

^{a/} Detailed description of the exposure scenarios are provided in Appendix F. n/a= not applicable or no exposure data available.

Table 18. Ambient and area sampling air concentrations of methyl bromide in or near commodity fumigation facilities^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
a. Fumigated rice warehouse (Appendix F Table 26)					
Ambient air	55	32	23	16	16
b. Sea van (Appendix F Table 29)					
15 feet downwind	4	3	0.9	n/a	n/a
c. Chamber (dried prunes) (Appendix F Table 29)					
1 m from door	n/a	0.78	0.3	n/a	n/a
2 m from chamber	n/a	141	60	n/a	n/a
15 m from chamber	n/a	5	2	n/a	n/a
leak check, side seal	n/a	13281	5692	n/a	n/a
d. Big Chamber (raisins) (Appendix F Table 29)					
shed-green forklift	n/a	117	84	62	43
shed-blue tractor	n/a	173	124	92	64
aeration sheds	2175	1292±780	923±557	717±433	501±303
capper area	n/a	280	200	187	130
hopper area	n/a	61	44	41	28
catchoff area	n/a	127	91	85	59
side hopper area	n/a	23	16	15	11
stem picker area	n/a	10	7	7	5
filer area	n/a	19	14	13	9
e. Chambers (raisins) (Appendix F Table 29)					
chamber	n/a	88	75	62	24
cage	n/a	54	46	38	15
leak checker	6	4	n/a	n/a	n/a
aeration	186	116	99	81	32
clearing	66	46	39	32	13
hopper area	13	8	7	6	3
stem picker area	30	27±3	23±0	19±2	11±1

Table 18. Ambient and area sampling air concentrations of methyl bromide in or near commodity fumigation facilities (continued)^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
f. Fumigation chambers(walnuts) (Appendix F Table 29)					
fumigatorium	n/a	75	64	42	17
sorting area	529	323	277	251	n/a
cracking area	n/a	1089	933	847	n/a
vac. chamber area	n/a	1789	1533	1391	n/a
clearing building	n/a	802	687	453	185
g. Sea van aeration (Appendix F Table 29)					
upwind of van	n/a	9	8	7	n/a
downwind- center	32	18±13	15±11	14±10	n/a
downwind- left and right	41	17±12	15±10	13±9	n/a
h. Fumigation of cherries (Appendix F Table 30)					
control room start-up	366	171±147	122±105	n/a	n/a
control room left overnight	26	11±10	8±7	n/a	n/a
dump station	27	18±7	13±5	n/a	n/a
i. Walnut processing facility (Appendix F Table 31)					
1. area samples					
meats pool	126	71±53	20±15	n/a	n/a
warehouse	66	53±18	15±5	n/a	n/a
sorting line	86	57±46	16±13	n/a	n/a
west alleyway	n/a	28	8	n/a	n/a
west cage door	n/a	20	6	n/a	n/a
east alleyway	n/a	6	2	n/a	n/a
east cage door	n/a	6	2	n/a	n/a
2. on-site ambient air					
vicinity of chambers	435	171±174	49±50	n/a	n/a
chamber near injection	6800	4100	1171	n/a	n/a
fence/ gate area (12/16/93)	5	2±2	0.6±0.6	n/a	n/a
chambers (12/20/93)	1933	576±907	165±259	n/a	n/a
fence /gate area (12/20/93)	5	2±1	0.6±0.3	n/a	n/a

Table 18. Ambient and area sampling air concentrations of methyl bromide in or near commodity fumigation facilities (continued)^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
i. Walnut processing facility (Appendix F Table 31)					
2. on-site ambient air					
chambers (12/20/93)	13	5±4	1±1	n/a	n/a
fence/gate area (12/20/93)	4	3±0.7	0.9±0.2	n/a	n/a
chamber (3/11/94)	1467	548±796	157±227	n/a	n/a
chamber (3/17/94)	47	17±18.5	5±5	n/a	n/a
fence	35	14±18.6	4±5	n/a	n/a
lamp post	6.3	5	1	n/a	n/a
north fence	21.9	14	4	n/a	n/a
north gate	7.9	6	2	n/a	n/a
pallets	64	36	10	n/a	n/a
polygon fence	27.6	13	4	n/a	n/a
polygon ramp	72	60	17	n/a	n/a
rooftop	13.2	7	2	n/a	n/a
scale house	24.9	19	5	n/a	n/a
south center&west fences	45.7	20±18	6±5	n/a	n/a
storage area	4.3	4	1	n/a	n/a
warehouse dock	198	111	32	n/a	n/a
west fence	12.2	10	3	n/a	n/a
3. compliance monitoring					
sorting line	343	318±28	273±24	265±23	n/a
cello packaging	375	355±26	304±22	296±22	n/a
bulk packaging	245	243	208	203	n/a
foreman's desk top	n/a	369	316	308	182
foreman's desk	n/a	28	24	23	14
fence between chambers	n/a	479	137	n/a	n/a

^{a/} Detailed description of the exposure scenarios are provided in Appendix F. n/a= not applicable or no exposure data available.

Table 19. Residential exposures to methyl bromide in fumigated homes or from living near commodity fumigation facilities^a.

Type of Application	Acute		Short-term	Seasonal	Chronic
	high (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)	mean±sd (ppb)
a. Outdoor and indoor air concentration during fumigation (Appendix F Table 34)					
outdoor air-20 feet	2990	522	522	n/a	n/a
indoor air-next to house	406	24	24	n/a	n/a
b. Outdoor and indoor air concentrations of neighboring houses during aeration (Appendix F Table 34)					
1. outdoor air					
10 feet	1064	296	n/a	n/a	n/a
50 feet	208	80	n/a	n/a	n/a
100 feet	74	40	n/a	n/a	n/a
2. Indoor air	168	60	n/a	n/a	n/a
c. Exposure of residents to methyl bromide in fumigated house (Appendix F Table 37)					
Southern California	n/a	210	172±146	n/a	n/a
Northern California	n/a	210	344±294	n/a	n/a
d. Exposure of residents to methyl bromide from living near commodity fumigation facilities (Appendix F Table 38)					
low range of exposure days	n/a	210	90	70	86
high range of exposure days	n/a	210	180	175	106
e. Exposure of residents to methyl bromide from living near fumigated fields (Appendices F and H)					
at or beyond the buffer zone	n/a	210	n/a	n/a	n/a

^{a/} Detailed description of the exposure scenarios are provided in Appendices F, H, and I. n/a= not applicable or no exposure data available.

Table 20. Summary of occupational and residential exposures to methyl bromide^a.

Scenarios	Workers at specific tasks (ppb)				Workers around the use of methyl bromide (ppb)				Residential exposures (ppb)			
	acute	short-term	seasonal	chronic	acute	short-term	seasonal	chronic	acute	short-term	seasonal	chronic
Preplant soil injection	0.6-835 (0.6-1659)	0.3-596	1.8-510	n/a	n/a	n/a	n/a	n/a	210	n/a	n/a	n/a
Nurseries and greenhouses	1-562 (2-724)	0.1-80	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Grain products	6-6039 (8-8458)	4-4314	2-3271	1-2482	32 (55)	23	16	16	210	90-180	70-175	86-105
Dried fruits and tree nuts	0.35-1434 (30-1463)	0.2-1229	2->1000	0.4-393	0.78-13281 (4-2175)	0.9-1533	6-1391	3-501				
Cherries for export	11-327 (47-877)	8-234	n/a	n/a	11-171 (27-366)	8-122	n/a	n/a				
Walnut processing	39-500 (70-636)	33-429	33-417	10	2-4100 (4-6800)	0.6-1171	23-308	14-182	n/a	n/a	n/a	n/a
Brewery facility	25-173 (25)	7-49	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
House during fumigation	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	24-522 (406-2990)	24-522	n/a	n/a
House during aeration	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	40-296 (74-1064)	n/a	n/a	n/a
Reentry into home	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	210	172-344	n/a	n/a

^{a/} Data were presented in Tables 16 to 19 of this document. n/a=not applicable or no exposure data were available. Ranges presented are those for the mean values. Ranges for the high end values are in parentheses.

IV.C. RISK CHARACTERIZATION

The potential health hazard associated with the use of methyl bromide was considered for occupational and residential exposures. Non-oncogenic effects were characterized in terms of a margin of exposure (MOE), defined as the ratio of the critical human equivalent NOEL to the estimated human exposure levels. For acute exposures, the high end values, or the mean if not available, were used to calculate the margins of exposure (MOEs). MOEs based on the high end values are indicated in bold in the Tables 21 to 24. For other exposure duration, the mean values were used to determine the MOEs. The oncogenic risk of exposure to methyl bromide was not evaluated since methyl bromide has not been shown to be oncogenic based on the current database. The critical NOELs used to address the various exposure scenarios for humans are listed in Table 15. The location of the type of application discussed in this section is indicated in italics. A summary of the margins of exposures is presented in Table 25.

IV.C.1. Occupational Exposure

The MOEs varied from <1 to greater than 1000 for workers using methyl bromide (Tables 21 and 22) and for those working in commodity fumigation facilities (Table 23). For the workers involved in the use of methyl bromide for preplant soil fumigation, the range of acute MOEs were 13 (*21a. tarp remover in shallow-shank tarp*) to 35000 (*21h. shovelman of shallow shank tarp method*) (Table 21). Applicators, co-pilots, cutters, pullers, cultipackers, and other workers had exposures between these two levels. The methods with acute MOEs of less than 100 were: shallow-shank tarp (*21a. all workers*), deep shank injection (*21d1. disc driver*), shallow shank tarped-bed (*21h. irrigation pipe tractor driver*), and tarp shallow with Noble plow (*21 i and j. cutter, puller, and some removers*). The short-term MOEs for most of these workers were also less than 100. The MOEs for seasonal exposures were all less than 100, except for tractor driver for shallow shank tarped-bed fumigations (*21g*), applicators of shallow shank tarped-bed fumigations (*21h*), and end puller of tarp shallow with Noble plow shank fumigations (*21j*).

For nursery potting soil workers, the use of the hot-gas method resulted in low acute MOEs of 29 and 63 for applicators and tarp venters, respectively (Table 21 *l*). The MOEs were 158 to 2625 for workers not using such a method (*21k*). However, the MOEs were greater than 100 for all workers after short-term exposures.

The MOEs were 2100 (acute) and 1714 (short-term) for workers in the rice warehouse with relatively low exposures (*22a*). For workers involved in the fumigation of grain products, the range of acute MOEs was 2 (aerator of sea container/trailer) to 2625 (forklift driver of chamber) (Table 22a). These aerators also had the lowest MOEs for short-term, seasonal, and chronic exposures. Applicators and aerator of tarp fumigation (*22a*) also had MOEs of less than 100 for all durations. The MOEs for forklift driver of container/trailer were 50 for seasonal and chronic exposures.

For workers involved in the fumigation of dried fruit and tree nut products, the range of acute MOEs was 14 (*22b4. worker involved in clear chamber for raisins*) to 60000 (*22b2. forklift operator of dried prunes in chamber*) (Table 22). For dried fruits, the fumigators and aerators had higher exposures than other workers; however, the acute and short-term MOEs were less

than 100 only for these workers in dried prunes (22b2), raisins (22b3), and walnut fumigations (22c). The seasonal MOEs were less than 100 for almost all workers. For walnuts in non-certified chambers (22c), the acute MOEs ranged from 12 (cracking worker) to 263 (cleaning-fumigator). The MOEs for most of the workers, except a few, were less than 100 for all exposure durations.

For workers involved in the fumigation of cherries for export, the range of acute MOEs was 24 (fumigator, start-up) to 447 (forklift driver) (Table 22d). The fumigators of start-up and closing-up had MOEs of 63 and 51, respectively, for short-term exposures. Short-term MOEs for other workers were greater than 100.

For workers in a walnut processing plant, the range of acute MOEs was 33 (vacuum chamber) to 300 (sorting area) (Table 22e). The MOEs were less than 100 for workers in the cleaning plant, Dock 5, and vacuum chamber areas for all duration of exposures.

For workers in a brewery, the acute MOEs ranged from 121 (area sample at entrance) to 840 (aerator) (Table 22f). The MOEs were equal to or greater than 245 for short-term exposures.

In the rice warehouse, the acute and short-term MOEs were 382 and 522, respectively, for ambient methyl bromide concentrations (23a). However, the MOE was only 13 for subchronic and chronic exposures. For exposure from working in commodity fumigation facilities (Table 23), the acute MOEs near the chambers were 2 for leak check in a dried prune chamber (23c), 10 in raisins aeration sheds (23d), 12 in walnut vacuum chamber area (23f), and 3 by the injection site of a chamber in a walnut processing facility (23i). The acute and short-term MOEs were generally much greater than 100 for other areas in these facilities, except for walnut fumigation chambers where the MOEs were less than 100 for most areas (23f).

IV.C.2. Residential Exposure

The MOEs for residents living next to a house during fumigation were 7 (outdoor air) and 52 (indoor air) for acute, 13 (outdoor air) and 292 (indoor air) for short-term exposures (Table 24 a to c). During aeration, only the MOE of 20 for outdoor air 10 feet from a fumigated house was less than 100 (Table 24).

The acute MOE for residents reentering fumigated homes and those living near commodity fumigation facilities or fumigated fields was 100 because the exposure was 210 ppb, the current regulatory limit (Table 24 c, d, and e). However, the MOEs were 20-78, 1, and 1 for short-term, seasonal, and chronic exposures, respectively; these were less than 100 for the potential exposures of some of these residents.

Table 21. Margins of exposure for occupational exposures to methyl bromide in preplant soil injection^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
a. Shallow-shank tarp method				
applicator	69	126	4	n/a
co-pilot	41	63	2	n/a
shovelman	41	190	n/a	n/a
tarp remover with PCO	13	20	<1	n/a
tarp remover with grower	38	152	n/a	n/a
b. Deep shank injection fumigation				
applicators	116	91	3	n/a
copilot	429	286	9	n/a
cultipacker	212	141	n/a	n/a
c. Improved deep shank injection fumigation				
applicator	368	245	8	n/a
cultipacker	300	200	n/a	n/a
d. Deep shank injection fumigation				
1. basic, second tractor				
applicator	239	160	5	n/a
disc driver	41	27	1	n/a
2. basic, cultipacker				
applicator	127	148	5	n/a
supervisor	313	211	7	n/a
cultipacker	362	414	n/a	n/a
e. Deep shank injection fumigation				
applicator	3000	2000	67	n/a
cultipacker	3000	2000	n/a	n/a
f. Shallow shank tarped-bed fumigation				
1. raised platform				
applicator	263	174	6	n/a
co-pilot	189	135	4	n/a
2. closing shoes				
applicator	477	316	10	n/a
co-pilot	100	84	3	n/a

Table 21. Margins of exposure for occupational exposures to methyl bromide in preplant soil injection (continued)^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
g. Shallow shank tarped-bed fumigation				
1. tractor with fumigation				
driver	750	500	17	n/a
applicator	467	308	10	n/a
drip tape layer	323	429	n/a	n/a
2. tractor with tarp				
driver	5250	4000	111	n/a
co-pilot	323	414	13	n/a
h. Shallow shank tarped-bed fumigation				
applicator	7000	4000	200	n/a
co-pilot	677	400	14	n/a
shovelman	35000	40000	n/a	n/a
pipe layer	10500	12000	n/a	n/a
irrigation pipe tractor driver	23	68	n/a	n/a
i. Tarp shallow with Noble plow shanks				
cutter (grower's)	89	522	n/a	n/a
cutter (company)	89	203	7	n/a
puller (grower's)	65	1333	n/a	n/a
puller (company)	65	500	18	n/a
j. Tarp shallow with Noble plow shanks				
1. PCO's				
cutter	269	214	8	n/a
remover:tractor driver	61	49	2	n/a
remover:basketman	65	52	2	n/a
remover:end puller	3000	2400	100	n/a
2. grower's				
cutter	269	214	n/a	n/a
remover:tractor driver	61	49	n/a	n/a
remover:basketman	65	52	n/a	n/a
remover:end puller	3000	2400	n/a	n/a

Table 21. Margins of exposure for occupational exposures to methyl bromide in preplant soil injection (continued)^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
k. Nursery potting soil fumigation				
applicator	808	4000	n/a	n/a
applicator assistant	677	6000	n/a	n/a
tarp remover	158	923	n/a	n/a
tractor driver	677	6000	n/a	n/a
truck driver	2625	30000	n/a	n/a
potter	221	2400	n/a	n/a
l. Greenhouse potting soil fumigation-hot gas method				
applicator	29	150	n/a	n/a
tarp venter	63	923	n/a	n/a
tarp remover	10500	120000	n/a	n/a

^{a/} Margins of exposure were based on exposure levels in Table 16 and human equivalents of the critical NOELs in Table 15 for each of the durations. The critical NOELs were: acute, 40 ppm (adult equivalent of 21 ppm) for developmental toxicity in rabbits; short-term, 20 ppm (adult equivalent of 12 ppm) for neurotoxicity in rabbits; seasonal, 0.5 ppm (adult equivalent of 200 ppb) for neurotoxicity in dogs; and chronic, 0.3 ppm (adult equivalent of 200 ppb) for nasal epithelial hyperplasia/degeneration in rats.

^{b/} Bolded margins of exposure were those based on the highest exposure values; other MOEs were based on the mean exposures.

Table 22. Margins of exposure for occupational exposures to methyl bromide in commodity fumigation facilities^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
a. Fumigation of grain products				
applicator	6	10	<1	<1
aerator (sea container/trailer)	2	3	<1	<1
aerator (tarp)	29	67	1	2
forklift driver (container/trailer)	488	1091	50	50
forklift driver (chamber)	2625	3000	100	200
workers (rice warehouse)	2100	1714	n/a	n/a
b. Fumigation of dried fruit and tree nut products				
1. sea van				
fumigator	636	1333	n/a	n/a
fumigator observer	2333	4000	n/a	n/a
2. chamber (dried prunes)				
forklift operator	60000	60000	n/a	n/a
fumigator	22	30	n/a	n/a
3. big chamber (raisins)				
primary fumigator	26	21	<1	1
secondary fumigator	178	143	3	4
aerator	23	34	1	1
forklift driver	304	343	8	11
catchall operator	103	83	1	2
hopper operator	93	105	2	3
capper dumper	165	132	2	3
inspector	1400	1091	20	29
moisture checker	2333	2000	33	50
forklift to side hopper	2625	2000	50	67
stem picker	750	750	13	20
packer	1105	857	15	22
4. chamber (raisins)				
fumigator	196	222	5	12
aerator	328	300	6	15
clear chamber	14	10	<1	1
stem picker	700	500	10	17
forklift driver	7000	4000	100	500
hopper operator	1105	750	15	25

Table 22. Margins of exposure for occupational exposures to methyl bromide in commodity fumigation facilities (continued)^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
c. Fumigation of two non- certified chambers (walnut)				
cracking worker	12	12	<1	n/a
sorting worker	27	24	<1	n/a
fumigator	19	25	1	2
cleaning-fumigator	263	174	4	11
cleaning-sort	19	16	<1	n/a
bulk casing worker	25	16	<1	n/a
in-shell packer	21	14	<1	n/a
hopper operator	27	18	<1	n/a
fumigator	93	62	2	4
d. Fumigation of cherries for export				
fumigator, start-up	24	63	n/a	n/a
fumigator, closing-up	48	51	n/a	n/a
fumigator, opening-up	169	293	n/a	n/a
forklift driver	447	1500	n/a	n/a
sorter	62	136	n/a	n/a
e. Walnut processing facility				
bulk packaging	284	364	6	n/a
cleaning plant	47	60	1	n/a
Dock 5	42	28	<1	n/a
fumigatorium	198	308	7	20
packaging	263	226	4	n/a
vacuum chamber	33	49	1	n/a
sorting	300	364	6	n/a
special cracking	124	226	4	n/a
f. Fumigation and aeration at a brewery facility				
1. application				
entry to open canisters	727	1446	n/a	n/a
area sample	500	1000	n/a	n/a
2. aeration				
aerator	840	1714	n/a	n/a
area sample	121-210	245-414	n/a	n/a

a/ Margins of exposure were based on exposure levels in Table 17 and human equivalents of the critical NOELs in Table 15 for each of the durations. The critical NOELs were: acute, 40 ppm (adult equivalent of 21 ppm) for developmental toxicity in rabbits; short-term, 20 ppm (adult equivalent of 12 ppm) for neurotoxicity in rabbits; seasonal, 0.5 ppm (adult equivalent of 200 ppb) for neurotoxicity in dogs; and chronic, 0.3 ppm (adult equivalent of 200 ppb) for nasal epithelial hyperplasia/degeneration in rats.

b/ Bolded margins of exposure were those based on the highest exposure values; other MOEs were based on the mean exposures.

Table 23. Margins of exposure for exposure to ambient and area sampling air concentrations of methyl bromide in or near commodity fumigation facilities^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
a. Fumigated rice warehouse	382	522	13	13
b. Sea van-15 feet downwind	5250	13333	n/a	n/a
c. Chamber (dried prunes)				
1 m from door	26923	40000	n/a	n/a
2 m from chamber	149	200	n/a	n/a
15 m from chamber	4200	6000	n/a	n/a
leak check, side seal	2	2	n/a	n/a
d. Big Chamber (raisins)				
shed-green forklift	179	143	3	5
shed-blue tractor	121	97	2	3
aeration	10	13	<1	<1
capper area	75	60	1	2
hopper area	344	273	5	7
catchoff area	165	132	2	3
side hopper area	913	750	13	18
stem picker area	2100	1714	29	40
filer area	1105	857	15	22
e. Chambers (raisins)				
chamber	239	160	3	8
age	389	261	5	13
leak checker	3500	n/a	n/a	n/a
aeration	113	121	2	6
clearing	318	308	6	15
hopper area	1615	1714	33	67
stem picker area	700	522	11	18
f. Fumigation chambers(walnuts)				
fumigatorium	280	188	5	12
sorting area	40	43	1	n/a
cracking area	19	13	<1	n/a
vac. chamber area	12	8	<1	n/a
clearing building	26	17	<1	1

Table 23. Margins of exposure for exposure to ambient and area sampling air concentrations of methyl bromide in or near commodity fumigation facilities (continued)^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
g. Sea van aeration				
upwind of van	2333	1500	29	n/a
downwind- center	656	800	14	n/a
downwind- left and right	512	800	15	n/a
h. Fumigation of cherries				
control room start-up	57	98	n/a	n/a
control room left overnight	808	1500	n/a	n/a
dump station	778	923	n/a	n/a
i. Walnut processing facility				
1. area samples				
meats pool	167	600	n/a	n/a
warehouse	318	800	n/a	n/a
sorting line	244	750	n/a	n/a
west alleyway	750	1500	n/a	n/a
west cage door	1050	2000	n/a	n/a
east alleyway	3500	6000	n/a	n/a
east cage door	3500	6000	n/a	n/a
2. on-site ambient air				
vicinity of chambers	48	245	n/a	n/a
chamber by injection	3	10	n/a	n/a
fence/ gate area (12/16/93)	4200	20000	n/a	n/a
chambers (12/20/93)	11	73	n/a	n/a
fence /gate area (12/20/93)	4200	20000	n/a	n/a
chambers (12/20/93)	1615	12000	n/a	n/a
fence/gate area (12/20/93)	5250	13333	n/a	n/a
chamber (3/11/94)	14	76	n/a	n/a
chamber (3/17/94)	447	2400	n/a	n/a
fence	600	3000	n/a	n/a
lamp post	3333	12000	n/a	n/a
north fence	959	3000	n/a	n/a
north gate	2658	6000	n/a	n/a
pallets	328	1200	n/a	n/a
polygon fence	761	3000	n/a	n/a
polygon ramp	292	706	n/a	n/a

Table 23. Margins of exposure for exposure to ambient and area sampling air concentrations of methyl bromide in or near commodity fumigation facilities (continued)^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
i. Walnut processing facility				
2. on-site ambient air				
rooftop	1591	6000	n/a	n/a
scale house	843	2400	n/a	n/a
south center&west fences	460	2000	n/a	n/a
storage area	4884	12000	n/a	n/a
warehouse dock	106	375	n/a	n/a
west fence	1721	4000	n/a	n/a
3. compliance monitoring				
sorting line	61	44	1	n/a
cello packaging	56	39	1	n/a
bulk packaging	86	58	1	n/a
foreman's desk top	57	38	1	1
foreman's desk	750	500	9	14
fence between chambers	44	88	n/a	n/a

^{a/} Margins of exposure were based on exposure levels in Table 18 and human equivalents of the critical NOELs in Table 15 for each of the durations. The critical NOELs were: acute, 40 ppm (adult equivalent of 21 ppm) for developmental toxicity in rabbits; short-term, 20 ppm (adult equivalent of 12 ppm) for neurotoxicity in rabbits; seasonal, 0.5 ppm (adult equivalent of 200 ppb) for neurotoxicity in dogs; and chronic, 0.3 ppm (adult equivalent of 200 ppb) for nasal epithelial hyperplasia/degeneration in rats.

^{b/} Bolded margins of exposure were those based on the highest exposure values; other MOEs were based on the mean exposures.

Table 24. Margins of exposures for residential exposures to methyl bromide in fumigated homes or from living near commodity fumigation facilities^a.

Type of Application	Acute ^b	Short-term	Seasonal	Chronic
a. Outdoor and indoor air concentration during house fumigation				
outdoor air-20 feet	7	13	n/a	n/a
indoor air-next to house	52	292	n/a	n/a
b. Outdoor and indoor air concentrations of neighboring houses during aeration				
1. outdoor air				
10 feet	20	n/a	n/a	n/a
50 feet	101	n/a	n/a	n/a
100 feet	284	n/a	n/a	n/a
2. Indoor air	125	n/a	n/a	n/a
c. Exposure of residents to methyl bromide in fumigated house (reentry)				
Southern California	100	41	n/a	n/a
Northern California	100	20	n/a	n/a
d. Exposure of residents to methyl bromide from living near commodity fumigation facilities				
low range of exposure days	100	78	1	1
high range of exposure days	100	39	1	1
e. Exposure of residents to methyl bromide from living near fumigated fields.				
at or beyond the buffer zones	100	n/a	n/a	n/a

^{a/} Margins of exposure were based on exposure levels in Table 19 and human equivalents of the critical NOELs in Table 15 for each of the durations. The critical NOELs were: acute, 40 ppm (adult equivalent of 21 ppm) for developmental toxicity in rabbits; short-term, 20 ppm (child equivalent of 7 ppm) for neurotoxicity in rabbits; seasonal, 0.5 ppm (child equivalent of 100 ppb) for neurotoxicity in dogs; and chronic, 0.3 ppm (child equivalent of 100 ppb) for nasal epithelial hyperplasia/degeneration in rats.

^{b/} Bolded margins of exposure were those based on the highest exposure values; other MOEs were based on the mean exposures.

Table 25. Summary of margins of exposure for occupational and residential exposures to methyl bromide^a.

Scenarios	Workers at specific tasks				Workers around the use of methyl bromide				Residential exposures			
	acute ^b	short-term	sea-sonal	chronic	acute ^b	short-term	sea-sonal	chronic	acute ^b	short-term	sea-sonal	chronic
Preplant soil injection	13->1000	20->1000	1-200	n/a	n/a	n/a	n/a	n/a	100	n/a	n/a	n/a
Nurseries in greenhouses	29->1000	150->1000	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Grain products	2->1000	3->1000	<1-100	<1-200	382	522	13	13	100	39-78	1	1
Dried fruits and tree nuts	14->1000	10->1000	<1-100	1-500	2->1000	2->1000	<1-33	<1-40				
Cherries for export	24-447	51->1000	n/a	n/a	57-808	98->1000	n/a	n/a				
Walnut processing	33-300	28-364	<1-7	20	3->1000	10->1000	1-9	1-14	n/a	n/a	n/a	n/a
Brewery facility	121-840	245->1000	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
House during fumigation	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	7-52	13-292	n/a	n/a
House during aeration	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	20-284	n/a	n/a	n/a
Reentry into home	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	100	20-41	n/a	n/a

a/ Data were presented in Tables 21 to 24 of this document. n/a=not applicable or exposure data were not available.

b/ Range of MOEs based on the highest or the mean exposure levels, if the highest was not available. ">1000" was used as the upper limit for this Table, actual MOEs are indicated in Tables 21-24.

V. RISK APPRAISAL

V.A. INTRODUCTION

The human health risk assessment of methyl bromide was conducted for occupational and residential exposures. Risk assessment is the process used to evaluate the potential for human exposure and the likelihood that the adverse effects of a substance will occur in humans under specific exposure conditions. Every risk assessment has inherent limitations on the application of existing data to estimate the potential risk to human health. Therefore, certain assumptions and extrapolations are incorporated into the hazard identification, dose-response assessment, and exposure assessment processes. This, in turn, results in uncertainty in the risk characterization which integrates all the information from the previous three processes. Qualitatively, risk assessments for all chemicals have similar uncertainties. However, the degree or magnitude of the uncertainty can vary depending on the availability and quality of the data, and the types of exposure scenarios being assessed. Specific areas of uncertainty associated with this risk assessment of methyl bromide are delineated in the following discussion.

V.B. HAZARD IDENTIFICATION

The uncertainties associated with the selection of the endpoints and the NOELs have already been discussed in some details (**IV.A. HAZARD IDENTIFICATION**).

V.B.1. Acute Toxicity

For acute inhalation exposure to methyl bromide, one of the critical NOELs was based on developmental effects observed in rabbits. The assumption was that methyl bromide will also cause developmental toxicity in humans. Epidemiological data were not available to support or refute this assumption for methyl bromide. Comparative studies of other agents showed that laboratory animal data were generally predictive of adverse developmental effects in humans (U.S. EPA, 1991). There are agents which caused developmental toxicity in laboratory animals but have not been clearly shown to be human developmental toxicants. Available human data were too limited to establish cause-effect relationships.

The risk for adults, based on developmental toxicity, may be overestimated because of the approach used to determine the human equivalent NOEL to calculate the margins of exposures. Consistent with the U.S. EPA Guidelines for Developmental Toxicity Risk Assessment (U.S. EPA, 1991), the developmental effects (gall bladder agenesis and fused sternebrae) observed at the end of the gestation (after 12 days of exposure) were assumed to be due a single day's exposure. The actual NOEL for a single day's exposure may be higher than that for 12-days of exposure. Furthermore, DPR determined a 24-hour time-weighted average dose by amortizing the NOEL from 6 hours to 24 hours of exposure using Haber's Rule ($C^n \times T = \text{effect}$ where $n=1$). This calculation resulted in an amortized dose that is one-fourth of the 6-hour NOEL and assumed that the dose would remain above the threshold for developmental effects. In an interim toxicology assessment for methyl bromide, the U.S. EPA also used amortization to determine the dose for the same endpoint. The calculated dosage (14 mg/kg/day) was different than the DPR dosage (21 mg/kg/day) because rabbit respiration rates

used by DPR (0.54 m³/kg/day) and U.S. EPA (0.37 m³/kg/day) were different (Hansen, 1993; Lewis, 1995).

On the other hand, a comparison of NOELs for neurotoxicity in the dog and human showed that the extrapolated level may not be an overestimation of the risk. As discussed in the Hazard Identification section (**IV.A. HAZARD IDENTIFICATION**), the human equivalent NOEL for neurotoxicity observed in the dog was 25 ppm for children. This level was comparable to the 21 ppm based on developmental toxicity in the rabbits. For comparison with neurotoxicity in humans, a report of worker exposure showed a LOEL of 35 ppm (Watrous, 1942) (**III.H.1. Occupational Exposure**). In that study, workers experienced anorexia, nausea, and headaches when exposed to an overall methyl bromide concentration of <35 ppm during a 2 week period (Watrous, 1942). If the actual no effect in one day is assumed to be 1/10 of the 35 ppm, then the NOEL would be 3.5 ppm. Using a default factor of 1/10 for intraspecies extrapolation, the regulatory level would be 0.35 ppm (or 350 ppb), a level only 1.5-fold higher than the 210 ppb (21 ppm x 1/10 for interspecies x 1/10 for intraspecies variations) for developmental toxicity used by DPR for regulation. However, there are uncertainties in determining whether the symptoms were associated with acute exposure in the Watrous (1942) study. First, the report did not specify whether the workers had previous exposures to methyl bromide. It is unlikely that inexperienced workers were used since all tasks were manually performed. Second, the report was not specific as to the relationship between actual exposure concentration and duration and symptoms observed. Higher exposures were likely to have occurred in the first week compared to the second week when control measures were instituted. It would have been useful to have individual data on the onset of symptoms as well as measured exposure concentrations for both weeks. Workers with both dermal and systemic effects due to methyl bromide spillage and other accidents, especially during the first week, were likely to be exposed to levels higher than 35 ppm.

V.B.2. Subchronic Toxicity

The endpoints for the critical short-term and seasonal exposures were based on neurotoxicity in the pregnant rabbit (Sikov *et al.*, 1981) and dogs (Newton, 1994b), respectively. Since methyl bromide is a known neurotoxicant, these endpoints are appropriate to use for hazard identification and risk characterization. The severity of the effects (convulsion and paresis) observed in the pregnant rabbits at 70 ppm and death after treatment had stopped suggested that a margin of exposure greater than the conventional benchmark of 100 might have to be considered.

It has been suggested that the decreased responsiveness observed in the dog after 34 exposures to 5 ppm should not be considered a treatment-related effect and is thus not appropriate for use as the critical NOEL for seasonal exposure. This suggestion was based on the assumption that effects after repeated exposure were only related to total dosage. The calculated total dosage for "decreased responsiveness" was less than those for "decreased activity" which was considered a treatment-related effect. DPR considers the analysis an oversimplification of the biological processes involved in the manifestation of the observed effects since the mechanism is unknown. The most important consideration is that the decreased responsiveness was specifically reported by an animal neurologist who had examined all 48 dogs on test previously on two occasions.

V.B.3. Chronic Toxicity

There were uncertainties associated with the use of hyperplasia/degeneration to the nasal cavity as the endpoint to evaluate chronic inhalation toxicity. One uncertainty was the interspecies variability in the nasal cavity between rodents and humans. Compared to humans, the increased complexity of the nasal turbinates, the straighter nasopharyngeal region, and the lower regional flow rates of rats might alter the toxicity at given exposure concentrations (Schreider, 1986). Additional information on the deposition, reactivity, solubility, absorption, metabolism, and clearance of methyl bromide in the nasal cavity epithelium of animals and humans would permit additional consideration of interspecies dosimetric adjustment (Jarabek *et al.*, 1990; U.S. EPA, 1990). Such information would also help define the relative contributions of regional versus systemically absorbed methyl bromide in the etiology of nasal damage.

As already discussed in the **IV.A. HAZARD IDENTIFICATION 2.c. Chronic Toxicity-Inhalation**, the use of the chronic NOEL based on effects in the nasal cavity may underestimate the risk since this NOEL expressed as human equivalents (0.1 ppm) is the same as that for the NOEL based on neurotoxicity after subchronic exposure.

V.B.4. Extrapolation of Estimated No-Effect Level from the Lowest-Effect Level

In this RCD, both the subchronic (seasonal) and chronic NOELs were estimated from the LOEL, the lowest dose tested. It is the DPR policy to use the 10-fold uncertainty factor (UF) to estimate the NOEL from a LOEL. Therefore, the estimated subchronic NOEL was 0.5 ppm based on neurotoxicity observed in two of eight dogs exposed to 5 ppm for 34 exposures. For chronic exposures, the estimated NOEL was 0.3 ppm based on a LOEL of 3 ppm for nasal epithelial hyperplasia and degeneration in the rat.

Instead of using a default 10-fold UF, two approaches have generally been discussed. One approach is to use the NOEL/LOEL ratios from the database. The problem with this approach is that these ratios reflect largely the dose selection of the studies, instead of toxicity. Two examples are provided.

(1) Methyl bromide is lethal to all species and shows a steep dose-response relationship in most cases. In one study, all rats exposed to 220 ppm methyl bromide died (100% mortality) but none died at 100 ppm (Irish, 1940). This implied that the UF could be 2-fold. However, the data also showed that the rats at 100 ppm suffered severe neurotoxicity and some were moribund. Obviously, neither the dose with no mortality can be used as the NOEL for risk assessment, nor can the factor of 2 be used as the UF for this study or other studies.

(2) With the developmental toxicity study (Breslin *et al.*, 1990), the NOEL and LOEL were 40 ppm and 80 ppm, respectively, for increased incidences of gall bladder agenesis and other effects. This also implied a 2-fold UF for extrapolation. On the other hand, the NOEL and LOEL ratio was 10-fold for parental (decreased fertility) and progeny (decreased pup body weights) effects in the reproductive toxicity study (American Biogenics Corp., 1986).

A more appropriate approach is the consideration of the magnitude of the response at the LOEL (Dourson *et al.*, 1996). The 10-fold UF may be excessive if the response rate at the

LOEL is marginal or low. On the other hand, the 10-fold UF may not be sufficient when the response at the LOEL is considered severe. If data are sufficient for determining the slope of the dose-response, a benchmark dose approach might be considered for estimating the NOEL.

In the case with the dog study used to derive the critical subchronic (seasonal) NOEL, there was a qualitative dose-response relationship between the exposure duration and concentration and the severity of neurotoxicity (Table 5). However, the magnitude of the response at the lowest dose (5 ppm) can not be compared with those at the higher doses since those experiments were not conducted for the same duration. Therefore, it is appropriate to use the 10-fold default factor as the UF for this study.

For the critical chronic NOEL, there was also a dose-response relationship between the exposure duration and concentration and the severity of the nasal cavity lesions in the rat (Table 7). At the LOEL (the lowest dose tested), the incidence was significantly elevated (p# 0.05), for both sexes, when compared to that in the controls. However, the lesion was described as very slight. The mildness of the lesion suggested that an UF of less than 10 might be sufficient to estimate the NOEL from the LOEL. While the default factor of 10-fold for the extrapolation was used as per DPR policy, a benchmark dose approach may be considered when guidelines for the use of the approach are established.

V.C. EXPOSURE ASSESSMENT

The uncertainties involved in occupational (soil or commodity fumigation) and residential (structural or commodity fumigation) exposures were addressed in the exposure assessment (**APPENDIX F. under EXPOSURE APPRAISAL**). In summary, the uncertainties were associated with the following factors: (1) conduct of some studies not in compliance with Good Laboratory Practices, in particular, absence of field recovery studies, (2) use of default factors when data were not available; for example, frequency and duration of exposure data, (3) exposures based on specific work task with no consideration of additional exposure occurred from working in other work task(s) for the remainder of the workday, and (4) no consideration for overtime work during peak use season. Therefore, the actual acute exposure was likely to be underestimated or overestimated because of these uncertainties.

There were numerous uncertainties involved in estimating residential exposures from field fumigations, both in the monitoring portions as well as the modeling portions of the exposure estimates (Appendices F and H). The most significant uncertainties included the following: (1) the precision and accuracy of the sampling and analytical methods, (2) influence of environmental factors on air concentrations, such as season and soil type, (3) application variability, such as the variation in application methodology and application rate, (4) use of default weather conditions in calculating buffer zone sizes, and (5) use of default assumptions in estimating air concentrations associated with overlapping applications, both in time and location. Actual exposure may be underestimated or overestimated because of these uncertainties.

V.D. RISK CHARACTERIZATION

The MOEs for potential acute, seasonal, and chronic exposures were based on NOELs for toxicity observed in laboratory animals. When the NOEL for non-oncogenic effects is based on animal data, a MOE of 100 is generally considered adequate for protection against potential acute or chronic toxicity of a chemical. This benchmark of 100 includes an uncertainty factor of 10 for interspecies extrapolation and a factor of 10 for intraspecies variability. These uncertainty factors assume that the average human is 10 times more sensitive to the effects of a chemical than the most sensitive laboratory animal, and that a sensitive individual is 10 times more susceptible than an average individual (Davidson *et al.*, 1986; Dourson and Stara, 1983).

V.D.1. Interspecies Extrapolation

The sensitivity of humans and laboratory animals to methyl bromide toxicity was difficult to compare because of inadequate exposure information in human case reports. For endpoints such as developmental toxicity and nasal hyperplasia/degeneration, there were no data for these effects in humans. For other endpoints such as death, human exposure levels were generally very high (> 1000 ppm) and the exposure durations were determined by the circumstance of cases and were limited to one individual in most cases. In addition, it is difficult to estimate the exposure since methyl bromide at high concentration tends not to be evenly distributed (Holling and Clarke, 1944). As a result, human case reports can not be used for comparison to the LC100 data from animal testing since the latter were experimentally determined and involved a large number of animals (Table 1). At lower concentrations of methyl bromide, a limited comparison was made with neurotoxicity observed in humans and dogs. In the following example, these data showed that symptoms in humans may be more severe than those in the dogs under similar exposure concentrations and durations (Table 26). However, a quantitative determination can not be made because of the scarcity of human data for comparison.

The current DPR default for interspecies extrapolation is a factor of 10-fold with respect to the dose. Accordingly, methyl bromide air concentrations were converted to the "dose" by taking into account the exposure concentration and duration, as well as the intake rate (respiration rate) of the exposed population. This approach is similar to that generally used for dietary exposure studies. The no-effect level is expressed as the dose after taking into account the consumption rate, instead of the concentrations in the diet. Thus, the net interspecies adjustment included the interspecies ratio of the intake rate, duration, and the 10-fold uncertainty factor.

Table 26: Comparison of neurotoxicity in humans and dogs.

Dose and duration	Human	Dog
156 ppm for 5 hours		lacrimation (Newton, 1994a)
400 ppm for 24 hours	difficulty in breathing, headaches, nausea, skin rash (Reidy <i>et al.</i> , 1994)	
150 ppm for 8 hours	nausea, vomiting, dizziness; later seizures, headaches, nausea, ataxia, and others (Hustinx <i>et al.</i> , 1993)	
103 ppm for 9 days		decreased activity and emesis (Newton, 1994b)
53 ppm for 14 days		decreased activity (Newton, 1994b)
overall <35 ppm over a 2 week period; onset and duration not specified	anorexia, nausea, headaches, vertigo, and other effects (Watrous, 1942)	

V.D.2. Intraspecies Extrapolation

For intraspecies variation in the response to methyl bromide, the default uncertainty factor of 10 was used because human illness/poisoning reports did not provide sufficient information to derive another factor. In these reports (discussed in **III.H. Neurotoxicity**), some individuals showed more severe symptoms than others. However, this difference in response was not quantified, and may only be quantified in well-conducted experimental studies.

Studies on the role of glutathione-S-transferase (GST) in methyl bromide metabolism and toxicity also provided evidence for variations in human response to methyl bromide. The glutathione-S-transferases are a multi-gene family of enzymes involved in the metabolism (activation and detoxification) of a wide variety of chemicals (Eaton and Bammler, 1999). They catalyze the general reaction: $GSH + R-X \rightarrow GSR + HX$. The mammalian soluble GSTs are divided into 4 main classes, alpha (A), mu (M), pi (P), and theta (T). The role of these enzymes in individual susceptibility to chemical exposure and toxicity is difficult to determine because of the large number of isozymes in the body. Second, GST expression varies among tissues. Not all isoforms are found in every tissue or all species. One important example, with respect to methyl bromide, is GSTT which is found in human erythrocytes but not in rodent erythrocytes. Third, GSTs have been found to be polymorphic in the human population. There are individuals who do not have the gene for certain GSTs. It has been determined that 50% of the Caucasian population do not have GSTM1, and 16% do not have GSTM3. GSTT has been found to be variable among different ethnic groups. The percentages of the population without the GSTT gene ranged from 9.7 % (Mexican-Americans) to 64.6 % (Chinese-Americans), as cited by

Garnier *et al.* (1996). With GSTP, the different variants of the enzyme are due to the transition mutation of a codon (s) such that other amino acids are substituted.

Methyl bromide has been identified as a substrate for GSTT (Eaton and Bammler, 1999). In 1990, Hallier *et al.* (1990) showed that when human erythrocyte cytoplasm was incubated with methyl bromide, there was a loss of methyl bromide in the gas phase with the formation of S-methylglutathione via an enzymatic reaction. Individuals which showed this activity were designated as “conjugators” while those with activity level comparable to boiled erythrocyte cytoplasm were “non-conjugators”.



This interaction of methyl bromide with sulfhydryl groups has been used in the study of methyl bromide workers. Iwasaki *et al.* (1989) and Goergens *et al.*, (1994) showed methylcysteine levels in hemoglobin proteins were higher in some methyl bromide workers compared to controls (Details of these studies are in **III.E.4.**). However, a quantitative relationship between adduct level and exposure was not determined.

There is evidence which shows conjugators, individuals with GSTT, may be more protected than non-conjugators from the genotoxicity of methyl bromide. In the study by Hallier *et al.*, (1993), methyl bromide was incubated with whole blood from conjugators and non-conjugators. Lymphocytes from conjugators had lower number (range of 6.51 to 7.95) of sister chromatid exchanges per cell than non-conjugators (range of 10.19 to 13.97). The lower level of SCEs in the conjugators, compared to the non-conjugators, was attributed to the reduced amount of methyl bromide available to interact with lymphocyte DNA because of reaction with erythrocyte proteins mediated by GSTT.

On the other hand, there is evidence that shows methyl bromide reaction with GST may be involved in the manifestation of neurotoxicity. In Davenport *et al.*, (1992), GST activity was inhibited in the brain of rats exposed to methyl bromide (details in **III.B.1.**). The GST activity was protected when the rats were either pre- or post-treated with an inhibitor of monohalomethane toxicity. In a report of poisoning of two workers, the non-conjugator had fewer neurotoxic effects when compared to the conjugator (Garnier *et al.*, 1996) (details in **III.H.1. Occupational Exposure**). The formation of S-methylglutathione, via conjugation of methyl bromide with GSH, in the brain of the conjugator was hypothesized to be involved in the neurotoxicity. The non-conjugator also had 2-fold higher concentrations of S-methylcysteine adduct in the erythrocytes; but the reaction was considered non-enzymatic. However, in another report of the same study, the worker designated as a conjugator may have been exposed to higher levels of methyl bromide because of an “inefficient” filter mask (Deschamps and Turpin, 1996).

In conclusion, the data show that the interaction of methyl bromide and GST is complex. While the polymorphism of GSTT in the human population is important to consider, it is not possible to quantitatively determine whether or not the variation is sufficiently addressed by the 10-fold default intra-individual uncertainty factor.

V.E. ISSUES RELATED TO THE FOOD QUALITY PROTECTION ACT

The Food Quality Protection Act of 1996 mandated U.S. EPA to “upgrade its risk assessment process as part of the tolerance setting procedures” (U.S. EPA, 1997b and c). The improvements to risk assessment were based on the recommendations from the 1993 National Academy of Sciences report, “Pesticides in the Diets of Infants and Children” (NRC, 1993). The Act required an explicit finding that tolerances are safe for children. U.S. EPA was required to use an extra 10-fold safety factor to take into account potential pre- and post-natal developmental toxicity and the completeness of the data unless U.S. EPA determined, based on reliable data, that a different margin would be safe. In addition, U.S. EPA must consider available information on: 1) aggregate exposure from all non-occupational sources; 2) effects of cumulative exposure to the pesticide and other substances with common mechanisms of toxicity; 3) the effects of *in utero* exposure; and 4) the potential for endocrine disrupting effects.

In recent documents, U.S. EPA used 3 criteria for the determination of the factor: (1) completeness and reliability of the toxicology database, (2) completeness and reliability of the exposure database, and (3) potential pre- and post-natal toxicity (U.S. EPA, 1999 a, b, and c). The latter criterion requires the consideration of evidence for developmental and reproductive toxicity, and developmental neurotoxicity.

V.E.1. Pre- and Post-natal Sensitivity

In recent years, issues have been raised on the adequacy of the risk assessment process to address the potential increased sensitivity of infants and children to pesticides. The basis of this concern was discussed in detail in the 1993 National Research Council report: *Pesticides in the Diets of Infants and Children* (NRC, 1993). In the 1994 DPR report: *A Joint Review of Existing Federal and State Pesticide Registration and Food Safety Programs* (also known as the PECC report), the PECC recommendation was to address the susceptibility issue on a case-by-case basis when specific data become available (DPR, 1994). A review of recent time-limited tolerances for pesticides showed that U.S. EPA generally applied an additional uncertainty factor of 10 for an incomplete database where one or more FIFRA required reproductive or developmental toxicity studies were not available. An additional uncertainty factor of 3 (MOE of 300) was used when the NOEL for developmental or reproductive effects was lower than that for maternal toxicity, or when there was no developmental neurotoxicity study. The criteria for the developmental toxicity testing included findings of teratogenicity to the central nervous system, neuropathology and neurotoxicity, hormone-like activity, and developmental toxicity (for effects other than structural abnormalities of the CNS) (Levine and Butcher, 1990). USEPA recently proposed that a development neurotoxicity study be included in the core toxicology data set (U.S. EPA, 1998c).

There was some evidence for increased sensitivity to the prenatal and post-natal toxicity of methyl bromide when NOELs for developmental or reproductive toxicity were compared with those for maternal toxicity. In the rabbit developmental toxicity study, the NOEL was 20 ppm for both effects in the fetus and neurotoxicity in the dam (Breslin et al., 1990b). However, it should be noted that the neurotoxicity was observed toward the end of the experiment while effects on the fetus were likely to have occurred earlier during organogenesis. In the developmental study conducted with rats, increased incidence of delayed skull ossification was reported in rat fetuses

with a NOEL of 20 ppm compared to a NOEL of >70 ppm for maternal toxicity (Sikov *et al.*, 1981). In the rat reproductive toxicity study, the NOEL was 3 ppm for reduced fertility in the F2b mated rats and for effects in the pups (decreased pup body weights and organs weights) (American Biogenics Corp., 1986). It is not known whether the reduced fertility in these F2b rats was due to exposure as adults or a manifestation of effects from previous *in utero* exposure (as fetuses from the F1 mating).

There may be a potential for increased sensitivity of infants and children to the neurotoxicity of methyl bromide based on consideration of the maturity of the central nervous system. In the young, the central nervous system is not fully developed at birth. Neuron proliferation and migration, blood-brain barrier, synaptic connections, as well as receptor and transmitter systems development continue after birth and into childhood (Rodier, 1995). Furthermore, there is a difference in the maturity of the brain development between laboratory animals and humans (Hoar and Monie, 1981; Dobbing and Sands, 1973). At birth, the rat brain is considered to be more developed than the human brain.

The potential impact of methyl bromide on the developing nervous system has not been evaluated. The toxicity studies for methyl bromide have only been conducted with adult animals and a developmental neurotoxicity study is not available. There is one case report in humans which showed that methyl bromide is more toxic to the young. In an acute exposure to methyl bromide in a home, an infant died while the parents recovered without apparent neurological deficits (Langard *et al.*, 1996). It is not known whether the severity of the effects on the infant was from increased exposure due to physiological differences (*i.e.* increased respiration rates) or increased sensitivity. Another concern is the potential cumulative toxicity from low level exposure to methyl bromide. As discussed in **IV.A.1.b. Neurotoxicity**, the dogs with prior exposure to relatively nontoxic level of methyl bromide (11 ppm) showed decreased activity earlier than those with no previous exposure when both groups were exposed to similar methyl bromide levels (156-158 ppm).

In this risk assessment for methyl bromide, inter-individual differences were accounted for by an uncertainty factor of 10. Given that methyl bromide is a potent neurotoxicant and there are inadequate toxicity information for infants and children, it may be prudent to consider an additional uncertainty factor to address the potential increased sensitivity for these population subgroups.

V.E.2. Aggregate Exposure

There could be a potential for aggregate exposure from occupation or residential exposures and dietary exposures. This potential scenario will be addressed in the subsequent Risk Characterization Document.

V.E.3. Cumulative Toxicity

Since the mechanism of methyl bromide toxicity is possibly due to alkylation of reactive groups, there is potential cumulative toxicity between methyl bromide and other chemicals with such a general mechanism of toxicity. The approach to address the cumulative risk of chemicals

is being discussed by the U.S. EPA Scientific Advisory Panel. The main focus of the discussion at this time is the toxicity of organophosphate pesticides.

V.E.4. Endocrine Effects

Based on the studies reviewed, methyl bromide has not been shown to cause endocrine disruption effects.

VI. CONCLUSIONS

The human health risk from potential inhalation exposure (occupational and residential) to methyl bromide was evaluated in this Risk Characterization Document. The critical toxicity endpoints were derived from experimental animals: developmental toxicity for acute exposure, neurotoxicity for short-term and seasonal exposures, and tissue damage to the nasal cavity for chronic exposures. For acute and chronic exposure endpoints, neurotoxicity was also considered in the determination of the critical NOELs.

The risks, expressed as the margins of exposure, were calculated for more than 180 exposure levels for workers involved in soil and commodity fumigations, and residents in the vicinity of structural and commodity fumigations. The acute exposure was based on the highest detected level when available. For longer exposure durations, the mean exposure level was used. The benchmark MOE traditionally considered as adequate for the protection of human health is a MOE of 100 based on an animal NOEL. Because of potential increased sensitivity of infants and children to the methyl bromide, an additional uncertainty factor was recommended for children exposure.

For exposure to methyl bromide, the MOEs varied from <1 to greater than 60,000. In general, the acute MOEs for workers close to the source of methyl bromide were less than 100. These workers included: applicators, copilots, shovelmen, removers, puller, tarp venters, fumigators, and aerators under certain work practices. In the commodity fumigation facilities, there was a wide range of MOEs for the monitored sites. Since these sites were specific to the particular facility, and the distance of the site to the source of methyl bromide is not known in most cases, it was not possible to make a conclusion about the magnitude of the MOEs.

For residential exposure from structural fumigation, the MOEs were less than 100 for residents living near (<50 feet) to a house being fumigated. For residents reentering fumigated homes and those living close to commodity fumigation facilities or fumigated fields, the acute MOE was 100 based on the permit conditions. However, the MOEs for short-term (1 week) exposures were less than 100.

The MOEs presented in this risk characterization document pertain only to the inhalation exposure from the air and does not include the concomitant exposures through dietary intake. It is essential that the significance of the MOEs be viewed in the context of the limitations and uncertainties discussed.

VII. REFERENCES

- Abdalla, N., D.J. Raski, B. Lear, and R.V. Schmitt, 1974. Distribution of methyl bromide in soils treated for nematode control in replant vineyards. *Pesticide Science* 5:259-269. (in DPR Vol.123-079).
- ACGIH, 1998. *Threshold Limit Values for Chemical Substances and Physical Agents, Biological Exposure Indices*. American Conference of Governmental Industrial Hygienists, Cincinnati, OH.
- ACGIH, 1997. *American Conference of Governmental Industrial Hygienists: "Chloropicrin," Documentation of the Threshold Limit Values, Sixth Edition*, Cincinnati, OH.
- Air Toxics Limited. 1993. *Report on Methyl Bromide Tarpaulin Fumigation Monitoring at San Pedro Dock Berth 54*. Report prepared for the Fresh Fruit Group in care of the California Grape and Tree Fruit League.
- Alexeeff, G.V. and W.W. Kilgore, 1983. Methyl bromide. *Residue Reviews* 88:101-153.
- Alexeeff, G.V. and W.W. Kilgore, 1985. Determination of acute toxic effects in mice following exposure to methyl bromide. *Journal of Toxicology and Environmental Health* 15:109-123 (in DPR Vol.123-135 #89716).
- AmeriBrom, Inc., 1983. *Toxicology*. DPR Vol.123-048.
- American Biogenics Corp., 1986. Two-generation reproduction study via inhalation in albino rats using methyl bromide. Study 450-1525. DPR Vol.123-82 #58196.
- Anger, W.K., J.V. Setzer, J.M. Russo, W.S. Brightwell, R.G. Wait, and B.L. Johnson, 1981. Neurobehavioral effects of methyl bromide inhalation exposures. *Scandinavian Journal of Work and Environmental Health* 7: Suppl 4:40-47.
- Anger, W.K., L. Moody, J. Burg, W.S. Brightwell, B.J. Taylor, J.M. Russo, N. Dickerson, J.V. Setzer, B.L. Johnson, and K. Hicks, 1986. Neurobehavioral evaluation of soil and structural fumigators using methyl bromide and sulfuryl fluoride. *NeuroToxicology* 7:137-156.
- ATSDR, 1992. *Toxicological profile for bromomethane*. TP-91/06. Agency for Toxic Substances and Disease Registry, Public Health Service, U.S. Department of Health & Human Services, Washington, D.C.
- ATSDR, 1996. *Minimal risk levels for priority substances and guidance for derivation*. Federal Register 61(101):25873-25882.
- Auletta, C.S., 1998. A chronic (12-month) toxicity study of methyl bromide fumigated feed in the dog. *Huntingdon Life Sciences*. Methyl Bromide Task Force, Inc. DPR Vol.123-209 #165490.

- Bentley, K.S., 1994. Detection of single strand breaks in rat testicular DNA by alkaline elution following in vivo inhalation exposure to methyl bromide. Haskell Laboratory Report No. 54-94. Chemical Manufacturers Association. DPR Vol.123-188 #162362.
- Blumberg, A., and W.B. Neli, 1967. Total body bromide excretion in a case of prolonged bromide intoxication. *Helvetica Medica Acta* 33: 330-333 (in DPR Vol.123-077 #4105).
- Bolt, H.M., 1996. Human GSH-transferase in risk assessment. In Biological Reactive Intermediates V. Basic Mechanistic Research in Toxicology and Human Risk Assessment (R. Snyder, I. G. Sipes, D. J. Jollow, T. J. Monks, J. J. Kocsis, G. F. Kalf, H. Greim, and C. M. Witmer, Ed.), pp. 405-409. Plenum Press, New York.
- Bond, J.A., J.S. Dutcher, M.A. Medinsky, R.F. Henderson, and L.S. Birnbaum, 1985. Disposition of ¹⁴C methyl bromide in rats after inhalation. *Toxicology and Applied Pharmacology* 78:259-267 (in DPR Vol.123-135 #89719).
- Boorman, G.A., 1984. Letter from Boorman to J.G. Vos, May 9, 1984. DPR Vol. 123-103.
- Boorman, G.A., H.L. Hong, C.W. Jameson, K. Yoshitomi, and R.R. Maronpot, 1986. Regression of methyl bromide-induced forestomach lesions in the rat. *Toxicology and Applied Pharmacology* 86:131-139.
- Boorman, G.A., K.T. Morgan, and L.C. Uriah, 1990. Nose, larynx, and trachea. In Pathology of the Fischer Rat (edited by Boorman, G.A., S.L. Eustis, M.R. Elwell, C.A. Montgomery, and W.F. MacKenzie, Ed.), pp. 315-337. Academic Press, Inc. New York, NY.
- Braun, A.L., and D.M. Supkoff, 1994. Options to methyl bromide for the control of soil-borne diseases and pests in California with reference to the Netherlands. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Breslin, W.J., C.L. Zablony, G.J. Bradley, K.D. Nitschke, and L.G. Lomax, 1990a. Methyl bromide inhalation teratology probe study in New Zealand white rabbits. The Toxicology Research Lab. Methyl Bromide Industry Panel. DPR Vol.123-138 #111266.
- Breslin, W.J., C.L. Zablony, G.J. Bradley, and L.G. Lomax, 1990b. Methyl bromide inhalation teratology study in New Zealand white rabbits. The Toxicology Research Lab. Methyl Bromide Industry Panel. DPR Vol.123-127 #95930.
- Bridges, R.G., 1955. The fate of labelled insecticide residues in food products. III. N-Methylation as a result of fumigating wheat with methyl bromide. *Journal of the Science of Food and Agriculture* 6:261-268.
- Brown, B.D., and D.E. Rolston, 1980. Transport and transformation of methyl bromide in soils. *Soil Science* 130:68-75. (in DPR Vol. 123-076 #39057).

- Burleigh-Flayer, H.D., W.J. Kintigh, and C.L. Benson, 1995. Chloropicrin: Vapor inhalation oncogenicity study in CD-1 mice. Laboratory Project ID#: 92N1105. Bushy Run Research Center. DPR Vol. 199-058 #136552.
- Busey, W.M., 1993. Neuropathological evaluation of brains from F0 and F1 rats in a two-generation reproduction study with methyl bromide-pathology report. EPL project number 303-007. DPR Vol.123-153 #125516.
- Butler, E.C.B., K.M.A. Perry, and J.R.F. Williams, 1945. Methyl bromide burns. British Journal of Industrial Medicine 2:30-31. DPR Vol.123-091 #59577.
- Calvert, G.M., G. Talaska, C.A. Mueller, M.M. Ammenheuser, W.W. Au, J.M. Fajen, L.E. Fleming, T. Briggie, and E. Ward, 1998a. Genotoxicity in workers exposed to methyl bromide. Mutation Research 417:115-128.
- Calvert, G.M., C.A. Mueller, J.M. Fajen, D.W. Chrislip, J. Russo, T. Briggie, L.E. Fleming, A.J. Suruda, and K. Steenland, 1998b. Health effects associated with sulfuryl fluoride and methyl bromide exposure among structural fumigation workers. American Journal of Public Health 88(12):1774-1780.
- Castro, C.E., and N.O. Belser, 1981. Photohydrolysis of methyl bromide and chloropicrin. Journal of Agricultural and Food Chemistry 29:1005-1008. (in DPR Vol.123-076 #39056, same as 123-100 #64925, 123-079 #52425).
- CDFA, 1995. Alternatives to methyl bromide: Research needs for California. Report of the Methyl Bromide Research Task Force. Pesticide Consultation and Analysis Unit, California Department of Food and Agriculture, Sacramento, CA.
- Clarke, C.A., C.G. Roworth, and H.E. Holling, 1945. Methyl bromide poisoning. An account of four recent cases met with in one of H.M. ships. British Journal of Industrial Medicine 2:17-23.
- California Code of Regulations, 1998. Standards Presentation to California Occupational Safety and Health Standards Board. Table AC-1 Permissible Exposure Limits for Chemical Contaminants. Title 8 Section 5155.
- CMA (Chemical Manufacturers Association), 1984. Methyl bromide residues in wheat and flour during fumigation, storage, cleaning, and milling in a commercial milling facility. Grocery Manufacturers of America, Inc. DPR Vol.123-089 #59570.
- CMA Methyl Bromide Industry Panel, 1998. Response to: California Department of Pesticide Regulation (CDPR) supplemental information request for study record 143945, dated September 5, 1997. Pharmaco-LSR study no. 94-3186: "A chronic (12-month) toxicity study of methyl bromide fumigated feed in the dog." Chemical Manufacturers Association. DPR Vol.123-208 #165489.
- CFR, 1989. Code of Federal Regulations 29 CFR Section 1910.1000.

- Curren, R.D., 1990. Unscheduled DNA synthesis in rat primary hepatocytes with a confirmatory assay. Study # T9152.380009. Microbiological Associates, Inc. DPR Vol.99-046 #088718.
- Danse, L.H.J.C., F.L. van Velsen, and C.A. van der Heijden, 1984. Methyl bromide: Carcinogenic effects in the rat forestomach. *Toxicology and Applied Pharmacology* 72:262-271. (in DPR Vol.123-043 #913094).
- Davenport, C.J., S.F. Ali, F.J. Miller, G.W. Lipe, K.T. Morgan, and M.S. Bonnefoi, 1992. Effect of methyl bromide on regional brain glutathione, glutathione-S-transferase, monoamines, and amino acids in F344 rats. *Toxicology and Applied Pharmacology* 112:120-127.
- Davidson, I.W.F., J.C. Parker, and R.P. Beliles, 1986. Biological basis for extrapolation across mammalian species. *Regulatory Toxicology and Pharmacology* 6:211-237.
- De Haro, L., J. Gastaut, J. Jouglard, and E. Renacco, 1997. Central and peripheral neurotoxic effects of chronic methyl bromide intoxication. *Clinical Toxicology* 35(1):29-34.
- Deschamps, F.J., and J.C. Turpin, 1996. Methyl bromide intoxication during grain store fumigation. *Occupational Medicine* 46(1):89-90.
- DFA, 1985. Organic methyl bromide residues of fumigated commodities. DPR Vol.123-128.
- Djalali-Behzad, G., S. Hussain, S. Osterman-Golkar, and D. Segerback, 1981. Estimation of genetic risks of alkylating agents. VI. Exposure of mice and bacteria to methyl bromide. *Mutation Research* 84:1-9. (in DPR Vol.123-109 #87809).
- Dobbing, J., and J. Sands, 1973. Quantitative growth and development of human brain. *Archives of Disease in Childhood* 48:757-767.
- Dourson, M.L., and J.F. Stara, 1983. Regulatory history and experimental support of uncertainty (safety) factors. *Regulatory Toxicology Pharmacology* 3:224-238.
- DPR, 1992. Toxicology review prompts changes in structural fumigations. Release no. 92-07, Department of Pesticide Regulation, California Environmental Protection Agency.
- DPR, 1993-1997. Pesticide Use Report-Annual. Information Services Branch, Department of Pesticide Regulation, California Environmental Protection Agency.
- DPR, 1994. A Joint Review of Existing Federal and State Pesticide Registration and Food Safety Programs. Department of Pesticide Regulation, California Environmental Protection Agency.
- Drawneek, W., M.J. O'Brien, H.J. Goldsmith, and R.E. Bourdillon, 1964. Industrial methyl bromide poisoning in fumigators. *The Lancet* 855-856. (in DPR Vol.123-051 #3694, same as 123-073 #24602, 123-076 #39060).

- Drew, R.T., 1983. A ten-day exposure inhalation toxicity study of methyl bromide in mice. The National Toxicology Program. Interagency Agreement Number 222-Y01-ES-20087. Brookhaven National Laboratory.
- Driscoll, C.D., and J.M. Hurley, 1993. Methyl bromide: single exposure vapor inhalation neurotoxicity study in rats. Bushy Run Research Center. Methyl Bromide Industry Panel. DPR Vol.123-158 #131609.
- Driscoll, C.D., S.J. Hermansky, and J.M. Hurley, 1994. Functional observational battery (FOB): Validation of procedures and personnel - 1992. Bushy Run Research Center. Methyl Bromide Industry Panel. DPR Vol.123-205 #161564.
- Eaton, D.L., and T.K. Bammler, 1999. Concise review of the glutathione S-transferases and their significance to toxicology. *Toxicological Sciences* 49:156-164.
- Extoxnet, 1999. Chloropicrin. Extoxnet, Extension Toxicology Network. Oregon State University. ([ace.orst.edu/cgi/bin/mfs/ol/pips/chloropi.htm?](http://ace.orst.edu/cgi/bin/mfs/ol/pips/chloropi.htm))
- Eustis, S.L., S.B. Haber, R.T. Drew, and R.S.H. Yang, 1988. Toxicology and pathology of methyl bromide in F344 rats and B6C3F1 mice following repeated inhalation exposure. *Fundamental and Applied Pharmacology* 11:594-610.
- Eustis, S.L., 1992. Toxicology and carcinogenesis studies of methyl bromide in B6C3F1 mice. NTP TR 385, U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health. DPR Vol.123-146 #116243.
- Fairall, R.J. and K.A. Scudamore, 1980. Determination of residual methyl bromide in fumigated commodities using derivative gas-liquid chromatography. *Analyst* 105:251-256.
- Farm Chemicals Handbook, 1998. Meister Publishing Company, Willoughby, OH.
- FDA, 1990-1991. Pesticide Residue Surveillance Monitoring Program, 1990-1991. Contaminants Division, U.S. Food and Drug Administration, Washington, D.C.
- Federal Register, 1985. Toxic Substances Control Act: Test Guidelines (Final Rule). Code of Federal Regulations. 40. part 798, subpart F. Office of the Register, National Archives and Records Administration. U.S. Government Printing Office, Washington, D.C.
- Federal Register, 1987. Revision of the TSCA Test Guidelines. *Federal Register* 52(97):19056-19082.
- Federal Register, 1991a. Code of Federal Regulations. 40. parts 180.3 (c)(2). Office of the Federal Register, National Archives and Records Administration. U.S. Government Printing Office, Washington, D.C.

Federal Register, 1991b. Code of Federal Regulations. 40. parts 156.10 and 158.34. Office of the Federal Register, National Archives and Records Administration. U.S. Government Printing Office, Washington, D.C.

Fieser, J.A., and B.A. Conrath, 1993. Magnitude and persistence of methyl bromide residues following fumigation of various food commodities. ABC/Del Monte Laboratories, Inc. DPR Vol.123-167 #137781.

Ford, J.H., M.G. Legendre, D.L. Ladner, J.A. Dawson, and C. Raymond, 1992. Automated closed-system headspace determination of methyl bromide in a variety of raw and processed nuts. J. AOAC International 75(2):328-333.

Friedman, M. and M.R. Gumbmann, 1979. Biological availability of e-N-methyl-L-lysine, 1-N-methyl-L-Histidine, and 3-N-methyl-L-histidine in mice. Nutrition Reports International 19(4):437-443.

Gansewendt, B., U. Foest, D. Xu, E. Hallier, H.M. Bolt, and H. Peter, 1991. Formation of DNA adducts in F-344 rats after oral administration or inhalation of [¹⁴C] methyl bromide. Food and Chemical Toxicology 29:557-563.

GAO, 1996. The phaseout of methyl bromide in the United States. Report to the Ranking Minority Member, Committee on Commerce, House of Representatives. GAO/RCED-96-16. U.S. General Accounting Office, Washington, D.C.

Garnier, R., M. Rambourg-Schepens, A. Muller, and E. Hallier, 1996. Glutathione transferase activity and formation of macromolecular adducts in two cases of acute methyl bromide poisoning. Occupational and Environmental Medicine 53: 211-215.

Gee, J., 1995. Memorandum from J. Gee to G. Patterson regarding methyl bromide and dog inhalation study. January 10, 1995. Dept. Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.

Gehring, P.J., R.J. Nolan, P.G. Watanabe, and A.M. Schumann, 1991. Chapter 14: Solvents, fumigants, and related compounds. In Handbook of Pesticide Toxicology, Vol. 2 Classes of Pesticides (W.J. Hayes, Jr., and E.R. Laws, Jr., Ed.), pp. 668-730. Academic Press, Inc., New York, N.Y.

Gennari, M., A. Cignetti, I. A. Gentile, and D. Vindrola, 1995. Effect of UV irradiation on the degradation of methyl bromide in air. Role of moisture and oxygen concentration. Pesticide Science 45:221-226.

Gibbons, D.B., H.R. Fong, R. Segawa, S. Powell, and J. Ross, 1996a. Methyl bromide concentrations in air downwind during aeration of fumigated single-family houses. HS-1713, March 20, 1996. Worker Health and Safety Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.

- Gibbons, D.B., H.R. Fong, R. Segawa, S. Powell, and J. Ross, 1996b. Methyl bromide concentrations in air near fumigated single-family houses. HS-1717, March 20, 1996. Worker Health and Safety Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Gill, M.W., 1989a. Single-dose functional observational battery validation study with chlorpromazine (CPZ) and amphetamine (AMP) in rats. DPR Vol.123-170 #143173.
- Gill, M.W., 1989b. Two-week repeated dose limb grip strength validation study with acrylamide (ACR) in rats. DPR Vol.123-170 #143175.
- Gill, M.W., 1989c. Single dose motor activity validation study with chlorpromazine (CPZ) and amphetamine (AMP) in rats. DPR Vol.123-171 #143176.
- Gill, M.W., 1989d. Two-week repeated dose functional observational battery validation study with acrylamide (ACR) and iminodipropionitrile (IDP) in rats. DPR Vol.123-171 #143178.
- Gillis, M. 1992. Verification of Chart I for Deep Tarped - Offsite. TRICAL, Inc. Study Number TC203.
- Gillis, M. 1998a. Results of First Monitoring Test. TRICAL, Inc. Study Number TC324.1.
- Gillis, M. 1998b. Methyl Bromide Soil Fumigation Monitoring Test #2 Results. TRICAL, Inc. Study Number TC324.2.
- Gillis, M. and J. Becker. 1993. New High Barrier Film Evaluation Studies. TRICAL, Inc. Study Number TC233.
- Goergens, H.W., E. Hallier, A. Muller, and H.M. Bolt, 1994. Macromolecular adducts in the use of methyl bromide as fumigant. Toxicology Letters 72:199-203.
- Golder Associates, Inc., 1985a. Florida methyl bromide sampling study. Methyl Bromide Industry Panel. DPR Vol. 123-064 #56726.
- Golder Associates, Inc., 1985b. California methyl bromide sampling study. Methyl Bromide Industry Panel. DPR Vol. 123-064 #56727.
- Gosselin, P.H., 1997a. Methyl bromide monitoring in Ventura County, first Lamplighter application (first 24 hour). Memorandum, August 26, 1997 from Gosselin to John S. Sanders. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Gosselin, P.H., 1997b. Methyl bromide monitoring in Ventura County, first Lamplighter application (second 24 hour). Memorandum, August 27, 1997 from Gosselin to John S. Sanders. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.

- Gosselin, R.E., H.C. Hodge, R.P. Smith, and M.N. Gleason, 1976. In Clinical Toxicology of Commercial Products, Acute Poisoning. Fourth edition. Williams and Wilkins, Baltimore, MD.
- Gotoh, K., T. Nishizawa, T. Yamaguchi, H. Kanou, T. Kasai, M. Ohsawa, H. Ohbayashi, S. Aiso, N. Ikawa, S. Yamamoto, T. Noguchi, K. Nagano, M. Enomoto, K. Nozaki, and H. Sakabe, 1994. Two-year toxicological and carcinogenesis studies of methyl bromide in F344 rats and BDF1 mice. Proceedings to Second Asia-Pacific Symposium on Environmental and Occupational Health (22-24 July, 1993 in Kobe). DPR Vol.123-174 #143944.
- Grant, W.M., 1974. In Toxicology of the Eye. Second edition. pp. 680-685. Charles C. Thomas Publisher, Springfield, IL.
- Greenberg, J.O., 1971. The neurological effects of methyl bromide poisoning. *Industrial Medicine* 40:27-29. (in DPR Vol. 123-048 #4107).
- Hallier, E., S. Deutschmann, C. Reichel, H.M. Bolt, and H. Peter, 1990. A comparative investigation of the metabolism of methyl bromide and methyl iodide in human erythrocytes. *International Archives of Occupational and Environmental Health* 62:221-225.
- Hallier, E., T. Langhof, D. Dannappel, M. Leutbecher, K. Schroder, J.W. Goergens, A. Muller, and H.M. Bolt, 1993. Polymorphism of glutathione conjugation of methyl bromide, ethylene oxide and dichloromethane in human blood: Influence on the induction of sister chromatid exchanges (SCE) in lymphocytes. *Archives of Toxicology* 67:173-178.
- Hansen, L.J., 1993. Methyl bromide. Toxicology assessment for applicator/residential exposure risk assessment. Memorandum from L.J. Hansen to L. Kutney, December 13, 1993. Office of Prevention, Pesticides and Toxic Substances. U.S. Environmental Protection Agency, Washington, D.C.
- Hansen, L.J., 1994. Methyl bromide. Evaluation of rat testicular DNA alkaline elution assay submitted as 6(a)(2) data. Memorandum from L.J. Hansen to L. Schnaubelt and B. O'Keefel, June 22, 1994. Office of Prevention, Pesticides and Toxic Substances. U.S. Environmental Protection Agency, Washington, D.C.
- Hardin, B.D., G.P. Bond, M.R. Sikov, F.D. Andrews, R.P. Beliles, and R.W. Niemeier, 1981. Testing of selected workplace chemicals for teratogenic potential. *Scandinavian Journal of Work and Environmental Health* 7 (Suppl. 4): 66-75.
- Hardisty, J.F., 1992. Histopathological evaluation of brains from rats-inhalation study of methyl bromide. EPL project number 303-007. DPR Vol. 123-142 #113606.
- Hardisty, J.F., 1997. Chronic (29-month) inhalation toxicity and carcinogenicity study of methyl bromide in rats. Reexamination of nasal cavity. Study number # B-91-8213/002. DPR Vol. 123-178 #156300.

Harsch, D.E. and R.A. Rasmussen, 1977. Identification of methyl bromide in urban air. *Analytical Letters* 10(13):1041-1047.

Hartsell, P.L., H.D. Nelson, and W.L. Stanley, 1983. A guide to methyl bromide fumigation of almonds and almond products, second edition. DPR Vol.123-047 #913087 and 913205, same as DPR Vol.123-089 #59532.

Hartsell, P.L., J.C. Tebbets, H.D. Nelson, and P.V. Vail, 1984. Large scale confirmatory studies: Residues. Walnut Research Reports, Walnut Marketing Board. DPR Vol.123-089 #59530.

Hartsell, P.L., H.D. Nelson, J.C. Tebbets, and P.V. Vail, 1986. Methyl bromide fumigation treatments for pistachio nuts to decrease residues and control navel orangeworm, *Amyelois transitella* (Lepidoptera: Pyralidae). *Journal of Economic Entomology* 79:1299-1302.

Hartsell, P.L., G.L. Obenauf, R.E. Gunnerson, and F.A. Mosebar, 1989. Methyl bromide residue protocols. (submitted to DPR in 1992 as part of data call in).

Harvey, S.C., 1985. Hypnotics and sedatives. In Goodman and Gilman's The Pharmacological Basis of Therapeutics (Gilman, A.G., L.S. Goodman, T.W. Rall, and F. Murad, Ed.), p. 339. Macmillan Publishing Co., New York, NY.

Harvey, J.M., C.M. Harris, and P.L. Hartsell, 1982. Commodity treatments: responses of nectarines, peaches, and plums to fumigation with methyl bromide. U.S. Department of Agriculture Marketing Research Report No. 1124. DPR Vol.123-089 #59573 and 123-144 #116225.

Haschek, W.M. and H.R. Witschi, 1991, Respiratory system. In Handbook of Toxicologic Pathology (Haschek, W.M. and C.G. Rousseaux, Ed.), pp.761-827. Academic Press, New York, NY.

Hastings, L., 1990. Sensory neurotoxicology: Use of the olfactory system in the assessment of toxicity. *Neurotoxicology and Teratology* 12:455-459.

Hazel, W.J., 1988. Methyl bromide Registration Standard. Protocol for almond residue studies (I.D. No. 53201-1; record No. 214966; RCB No., 3422), Office of Pesticides and Toxic Substances, U.S. Environmental Protection Agency. (Submitted to DPR as part of Data Call-in, received May 11, 1992).

Helliker, P.E., 1999. Status of methyl bromide permit conditions. Letter to Interested Parties, August 31, 1999. Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.

Herzstein, J., and M.R. Cullen, 1990. Methyl bromide intoxication in four field-workers during removal of soil fumigation sheets. *American Journal of Industrial Medicine* 17:321-326.

- Heuser, S.G., and K.A. Scudamore, 1970. Selective determination of ionized bromide and organic bromides in food-stuffs by gas-liquid chromatography with special reference to fumigant residues. *Pesticide Science* 1:244-249. (in DPR Vol. 123-060 #39042).
- Hezemans-Boer, M., J. Toonstra, J. Meulenbelt, J. Zwaveling, B. Sangster, and W. A. van Vloten, 1988. Skin lesions due to exposure to methyl bromide. *Archives of Dermatology* 124:917-921.
- Hicks, S.C. 1992a. Deep Shank Injection for Measurement of Methyl Bromide Drift Offsite. Siemer and Associates, Inc. Study Number 924096E2.
- Hicks, S.C. 1992b. Shallow Shank Non-Tarp Injection for Measurement of Methyl Bromide Drift Offsite. Siemer and Associates, Inc. Study Number 924096E1.
- Hicks, S.C. 1993a. Deep-Shank, Non-Tarped Fumigation - Mitigation of Methyl Bromide Worker Exposure and Offsite Drift. Siemer and Associates, Inc. Study Number 934104.2-1.
- Hicks, S.C. 1993b. Measurement of Onsite and Offsite Exposure From Methyl Bromide Fumigation Applied by the Deep Shank Non-Tarp Method. Siemer and Associates, Inc. Study Number 934100B1.1.
- Hicks, S.C. 1993c. Measurement of Onsite and Offsite Exposure From Methyl Bromide Fumigation Applied by the Tarped Bed Method. Siemer and Associates, Inc. Study Number 934110F1.
- Hicks, S.C. 1996. Methyl Bromide Deep Shank, Non-Tarp Soil Application. Siemer and Associates, Inc. Study Number 954110.1.
- Hoar, R.M., and I.W. Monie, 1981. Comparative development of specific organ systems. In Developmental Toxicology (C.A. Kimmel and J. Buelke-Sam, Ed.), pp.13-33. Target Organ Toxicology Series. Raven Press, New York.
- Holling, H.E., and C.A. Clarke, 1944. Methyl bromide intoxication. *Journal of the Royal Naval Medical Service* 30:218-224.
- Honma, T., A. Sudo, M. Miyagawa, M. Sato, and H. Hasegawa, 1982. Significant changes in monoamines in rat brain induced by exposure to methyl bromide. *Neurobehavioral Toxicology and Teratology* 4:521-524.
- Honma, T., M. Miyagawa, M. Sato, and H. Hasegawa, 1985. Neurotoxicity and metabolism of methyl bromide in rats. *Toxicology and Applied Pharmacology* 81:183-191.
- Honma, T., M. Miyagawa, and M. Sato, 1987. Methyl bromide alters catecholamine and metabolite concentrations in rat brain. *Neurotoxicology and Teratology* 9:369-375.
- Honma, T., M. Miyagawa, and M. Sato, 1991. Inhibition of tyrosine hydroxylase activity by methyl bromide exposure. *Neurotoxicology and Teratology* 13:1-4.

- Honma, T., M. Kanada, and M. Miyagawa, 1994. Behavioral evidence for modified receptor sensitivity in rat brain induced by methyl bromide exposure. *Industrial Health* 32:1-16.
- HSDB, 1994. Hazardous Substances Data Bank. National Library of Medicine, Bethesda, MD.
- Hubbs, A.F., 1986. The subchronic effects of oral methyl bromide administration in the rat. DPR Vol.123-083 #59183.
- Hurt, M.E., K.T. Morgan, and P.K. Working, 1987. Histopathology of acute toxic responses in selected tissues from rats exposed by inhalation to methyl bromide. *Fundamental and Applied Toxicology* 9:352-365. (in DPR Vol.123-109 #87805).
- Hurt, M.E., and P.K. Working, 1988. Evaluation of spermatogenesis and sperm quality in the rat following acute inhalation exposure to methyl bromide. *Fundamental and Applied Toxicology* 10:490-498. (in DPR Vol.123-109 #87804).
- Hurt, M.E., D.A. Thomas, P.K. Working, T.M. Monticell, and K.T. Morgan, 1988. Degeneration and regeneration of the olfactory epithelium following inhalation exposure to methyl bromide: Pathology, cell kinetics, and olfactory function. *Toxicology and Applied Pharmacology* 94:311-328.
- Hustinx, W.N.M., R.T.H. van de Laar, A.C. van Huffelen, J.C. Verwey, J. Meulenbelt, and T.J.F. Savelkoul, 1993. Systemic effects of inhalational methyl bromide poisoning: a study of nine cases occupationally exposed due to inadvertent spread during fumigation. *British Journal of Industrial Medicine* 50:155-159.
- Iwasaki, K., I. Ito, and J. Kagawa, 1989. Biological exposure monitoring of methyl bromide workers by determination of hemoglobin adducts. *Industrial Health* 27:181-183.
- I-R 4, 1982. Petition proposing a tolerance for residues of inorganic bromides in strawberries resulting from postharvest fumigation with methyl bromide. Interregional Research Project No. 4. Rutgers University. DPR Vol.123-060.
- I-R 4, 1983. Petition proposing a tolerance for methyl bromide for use in blueberry production. Interregional Research Project No. 4. Rutgers University. DPR Vol.123-060 #39047.
- Irish, D.D., E.M. Adams, H.C. Spencer, and V.K. Rowe, 1940. The response attending exposure of laboratory animals to vapors of methyl bromide. *Journal of Industrial Hygiene and Toxicology* 22:218-230.
- Jarabek, A., M. Menache, J. Overton, M. Dourson, and F. Miller, 1990. The U.S. Environmental Protection Agency's inhalation RFD methodology: Risk assessment for air toxics. *Toxicology and Industrial Health* 6:279-301.
- Jaskot, R.H., Grose, E.C., B.M. Most, M.G. Menache, T.B. Williams, and J.J. Roycroft, 1988. The distribution and toxicological effects of inhaled methyl bromide in the rat. *Journal of American College of Toxicology* 7:631-642. (in DPR Vol.123-135 #89717).

- Johnstone, R.T., 1945. Methyl bromide intoxication. *Industrial Medicine* 14(6):495-497.
- Jordi, A.U., 1953. Absorption of methyl bromide through the intact skin. *Aerospace Medicine* 24:536-539.
- Kane, L.E., C.S. Barrow, and Y. Alarie, 1979, A short-term test to predict acceptable levels of exposure to airborne sensory irritants. *American Industrial Hygiene Association Journal* 40:207-229.
- Kaneda, M., H. Hojo, S. Teramoto, and K. Maita, 1998. Oral teratogenicity studies of methyl bromide in rats and rabbits. *Food and Chemical Toxicology* 36:421-427.
- Kaneda, M., N. Hatakenaka, S. Teramoto, and K. Maita, 1993. A two-generation reproduction study in rats with methyl bromide-fumigated diets. *Food and Chemical Toxicology* 31(8):533-542.
- Kato, N., S. Morinobu, and S. Ishizu, 1986. Subacute inhalation experiment for methyl bromide in rats. *Industrial Health* 24:87-104.
- Kim, D. and R. Segawa. 1998a. Monitoring Results From a Tarped, Virtually Impermeable Film, Shallow Injection Methyl Bromide Application in Orange County Memorandum, September 4, 1998 from Kim and Segawa to Douglas Y. Okumura. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Kim, D. and R. Segawa. 1998b. Monitoring Results From Virtually Impermeable Film and High Density Polyethylene (High Barrier) Tarped, Shallow Injection Methyl Bromide Application in Orange County Memorandum, September 8, 1998 from Kim and Segawa to Douglas Y. Okumura. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Kimbell, J.S., E.A. Gross, D.R. Joyner, M.N. Godo, and K.T. Morgan, 1993. Application of computational fluid dynamics to regional dosimetry of inhaled chemicals in the upper respiratory tract of the rat. *Toxicology and Applied Pharmacology* 121:253-263.
- King, J.R., C.A. Benschoter, and A.K. Burditt, 1981. Residues of methyl bromide in fumigated grapefruit determined by a rapid, headspace assay. *Journal of Agricultural and Food Chemistry* 29:1003-1005. (in DPR Vol.123-116 #88773).
- Kiplinger, G.R., 1994. Acute oral toxicity comparison study of microencapsulated methyl bromide and liquid methyl bromide in albino rats. Study number WIL-49011. WIL Research Lab., Inc. DPR Vol.123-162 #132699.
- Kishi, R., I. Itoh, S. Ishizu, I. Harabuchi, and H. Miyake, 1991. Symptoms among workers with long-term exposure to methyl bromide. *Japan Journal of Industrial Health* 33:241-250.

- Kolbezen, M.J., D.E. Munnecke, W.D. Wilbur, L.H. Stolzy, F.J. Abu-El-Haj., and T.E. Szuszkiewicz, 1974. Factors that affect deep penetration of field soils by methyl bromide. *Hilgardia* 42: 465-492 (in DPR Vol.123-079, same as 123-022 #913084).
- Kramers, P.G.N., C.E. Voogd, A.G.A.C. Knaap, and C.A. van der Heijden, 1985. Mutagenicity of methyl bromide in a series of short-term test. *Mutation Research* 155:41-47. (in DPR Vol.123-109 #87802).
- Lambert, R.A., and L. Jackson, 1920. The pathology of chloropicrin poisoning. In Collected Studies on the Pathology of War Gas Poisoning (M.C. Winternitz, Ed.), pp. 69-90. New Haven: Yale University Press. (in DPR Vol.123-028 #913104)
- Langard, S., T. Rognum, O. Flotterod, and V. Skaug, 1996. Fatal accident resulting from methyl bromide poisoning after fumigation of a neighbouring house; leakage through sewage pipes. *Journal of Applied Toxicology* 16 (5):445-448.
- Le Goupil, M., 1932. Les proprietes insecticide du bromure de methyl. *Revue de Pathologie Vegetale et d'Entomologie Agricole de France* 19:169.
- Levine, T.E., and R.E. Butcher, 1990. Workshop on the qualitative and quantitative comparability of human and animal developmental neurotoxicity, Work Group IV Report: Triggers for developmental neurotoxicity testing. *Neurotoxicology and Teratology* 12:281-284.
- Lewis, C., 1995. Exposure Assessment- Section 18 use of methyl bromide by USDA/APHIS on imported fruits at ports of entry. Memorandum from C. Lewis to W. Dykstra. September 25, 1995. Office of Prevention, Pesticides and Toxic Substances. U.S. Environmental Protection Agency, Washington, D.C.
- Lewis, S.E., 1948. Inhibition of SH enzymes by methyl bromide. *Nature* 161:692-693.
- Lindsay, L.D, 1985. Memo to Frank Light regarding methyl bromide sample-State Food & Drug. May 24, 1985. Sun-Diamond Growers of California. Submitted as record 0065 as part of the 1992 DPR data call in.
- Liscombe, E.A., J. Sansone, and K. Leitner. 1995. Investigation of Mitigation Measures for Methyl Bromide Structural Fumigations. Soil Chemical Corporation.
- Liscombe, E.A., J. Sansone, and K. Leitner. 1996. Investigation of Mitigation Measures for Methyl Bromide Structural Fumigations Using the MAKR Process. Soil Chemical Corporation.
- Lobert, J.M., J.H. Butler, S.A. Montzka, L.S. Geller, R.C. Myers, and J.W. Elkins, 1995. A net sink for atmospheric CH₃Br in the East Pacific Ocean. *Science* 267:1002-1005.
- Lovelock, J.E., 1975. Natural halocarbons in the air and in the sea. *Nature* 256:193-194.

- Majewski, M.S., M.M. McChesney, J.E. Woodrow, J.H. Prueger, and J.N. Seiber, 1995. Aerodynamic measurements of methyl bromide volatilization from tarped and nontarped fields. *Journal of Environmental Quality* 24:742-752.
- Marraccini, J.V., G.E. Thomas, J.P. Ongley, C.D. Pfaffenberger, J.H. Davies, and L.R. Bednarczyk, 1983. Death and injury caused by methyl bromide, an insecticide fumigants. *Journal of Forensic Science* 28:601-607.
- Marty, M.A., B. Blaisdell, M. Hill, R. Broadwin, S. Hurley, and P. Jenkins, 1997. Breathing rate distribution for use in California's Air Toxics Hot Spots Program risk. *Toxicological Sciences* 42 (1-S):40.
- MBIP, 1984a. Fumigant survey. Flour and flour products. Oregon Department of Agriculture. DPR Vol.123-109 #87812.
- MBIP, 1984b. Determination of methyl bromide residues in strawberries after a commercial fumigation. DPR Vol.123-109 #87813, 123-089 #59535, and 123-144 #116230.
- MBIP, 1985a. Commodity fumigation trials. DPR Vol.123-089 Attachment 1 #59503, 123-144 #16228, and discussed in Hazel, 1988.
- MBIP, 1985b. Fumigation of Kerman pistachios with methyl bromide. Stored-Product Insects Research Laboratory. DPR Vol.123-089 #59534.
- MBIP, 1986. Methyl bromide field dissipation study. DPR Vol.123-081 #58907, same as 123-079 #52429.
- MBIP, 1988a. Methyl bromide residue study (Pre-plant). Bolsa Research Associates. DPR Vol. 123-109 #87810.
- MBIP, 1988b. Removal of residues from post-harvest treatments. DPR Vol.123-109 #87811.
- MBIP, 1992a. Methyl Bromide. Cover letter (July, 1, 1992) from Dee Kuhn to Wayne Stringer. DPR Vol.123-144.
- McGregor, D.B., 1981. Tier II mutagenic screening of 13 NIOSH priority compounds. National Institute for Occupational Safety and Health. Inveresk Research Int. Ltd. DPR Vol.123-103 #66718, #66719, # 66720, #66721, and #66722.
- McKenry, M.V., and S. Secara, 1990. Dissipation of methyl bromide following treatment of a dwelling for scale broom weed. DPR Vol.123-114 #91023.
- Medinsky, M.A., J.A. Bond, J.S. Dutcher, and L.S. Birnbaum, 1984. Disposition of [¹⁴C] methyl bromide in Fischer-344 rats after oral or intraperitoneal administration. *Toxicology* 32:187-196. (in DPR Vol.123-135 #89720).

- Medinsky, M.A., J.S. Dutcher, J.A. Bond, R.F. Henderson, J.L. Mauderly, M.B. Snipes, J.A. Mewhinney, Y.S. Cheng, and L.S. Birnbaum, 1985. Uptake and excretion of [¹⁴C] methyl bromide as influenced by exposure concentration. *Toxicology and Applied Pharmacology* 78:215-225. (in DPR Vol.123-135 #89718).
- Mehler, L., 1999. Summary report of cases reported to the California Pesticide Illness Surveillance Program 1982-1996. Worker Health and Safety Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Mertens, J.J.W.M., 1997. A 24-month chronic dietary study of methyl bromide in rats. Laboratory Study Number WIL-49014. WIL Research Laboratories. DPR Vol.123-179 #158746.
- Michalodimitrakis, M.N., A.M. Tsatsakis, M.G. Christakis-Hampsas, N. Trikilis, and P. Christodoulou, 1997. Death following intentional methyl bromide poisoning: Toxicological data and literature review. *Veterinarian and Human Toxicology* 39 (1):30-34.
- Miller, J.W., 1943. Fatal methyl bromide poisoning. *Archives of Pathology* 36:505-507.
- Mitsumori, K., K. Maita, T. Kosaka, T. Miyaoka, and Y. Shirasu, 1990. Two-year oral chronic toxicity and carcinogenicity study in rats of diets fumigated with methyl bromide. *Food and Chemical Toxicology* 28:109-119 (in DPR Vol.123-127 #095929).
- Miyagawa, M., 1982. Conditioned taste aversion induced by inhalation exposure to methyl bromide in rats. *Toxicology Letters* 10:411-416.
- Moosa, M.R., J. Jansen, and C.L. Edelstein, 1994. Treatment of methyl bromide poisoning with haemodialysis. *Postgraduate Medicine Journal* 70:733-735.
- Moriya, M., T. Ohta, K. Watanabe, T. Miyazawa, K. Kato, and Y. Shirasu, 1983. Further mutagenicity studies on pesticides in bacterial reversion assay systems. *Mutation Research* 116: 185-216. (in DPR Vol.123-109 #87808).
- Morris, J.B., D.N. Hassett, and K.T. Blanchard, 1993. A physiologically based pharmacokinetic model for nasal uptake and metabolism of nonreactive vapors. *Toxicology and Applied Pharmacology* 123:120-129.
- Mortelmans, K.E., and G.F. Shepherd, 1980. *In vitro* microbiological mitotic recombination assay of methyl bromide using *S. cerevisiae* D3. SRI International. DPR Vol.123-044 #913095.
- Munnecke, D.E., and Van Gundy, S.D., 1979. Movement of fumigants in soil, dosage responses, and differential effects. *Annual Review of Phytopathology* 17:405-429 (in DPR Vol.123-079).
- Naas, D.J., 1990. Acute oral toxicity study in beagle dogs with methyl bromide. WIL Research Lab. Inc. DPR Vol.123-124 #91578.

- Nelson, H.D., P.L. Hartsell, and P.V. Vail, 1984. Summary of research conducted to develop a quarantine treatment for the codling moth in walnuts for export to Japan. Walnut Research Reports, Walnut Marketing Board. DPR Vol.123-089 #59530.
- Newton, P.E., 1994a. An up-and-down acute inhalation toxicity study of methyl bromide in the dog. Study number 93-6067. Pharmaco LSR. DPR Vol.123-163 #132818.
- Newton, P.E., 1994b. A four week inhalation toxicity study of methyl bromide in the dog. Study number 93-6068. Pharmaco LSR. DPR Vol.123-164 #132821.
- Newton, P.E., 1996. A chronic (12-month) toxicity study of methyl bromide fumigated feed in the dog. Study no. 94-3186. Huntingdon Life Science. DPR Vol.123-175 #143945.
- Nikula, K.J. and J.L. Lewis, 1994. Olfactory mucosal lesions in F344 rats following inhalation exposure to pyridine at threshold limit value concentrations. *Fundamental and Applied Pharmacology* 23:510-517.
- NIOSH, 1997. Methyl bromide IDLH Documentation. (<http://www.cdc.gov/niosh/74839.html>).
- NIOSH, 1999. Chloropicrin, IDLH Documentation. (<http://www.cdc.gov/niosh/idlh/76062.html> and <http://www.cdc.gov/niosh/idlh/idlhintr.html>)
- NOAA/NASA/UNEP/WMO, 1995. Scientific Assessment of Ozone Depletion: 1994, Executive Summary. World Meteorological Organization Global Ozone Research and Monitoring Project - Report No. 37.
- Norris, J.C., C.D. Driscoll, and J.M. Hurley, 1993a. Methyl bromide: ninety-day vapor inhalation neurotoxicity study in CD rats. Bushy Run Research Center. Methyl Bromide Industry Panel. DPR Vol.123-159 #131619.
- Norris, J.C., C.D. Driscoll, and J.M. Hurley 1993b. Methyl bromide: ninety-day vapor inhalation neurotoxicity study in CD rats- Amendment 1. Bushy Run Research Center. Methyl Bromide Industry Panel. DPR Vol.123-160 #131621.
- NRC (National Research Council), 1993. Pesticides in the Diets of Infants and Children. National Academy Press, Washington, D.C.
- NTP, 1992. Toxicology and carcinogenesis studies of methyl bromide in B6C3F1 mice. NTP TR 385, U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health. DPR Vol.123-145 #76659.
- OEHHA, 1994. Proposition 65 Developmental and Reproductive Toxicant (DART) Identification Committee Public Meeting, May 5, 1994 at the Air Resources Board. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, CA.

- OEHHA, 1996. Draft: Technical Support Document: Exposure Assessment and Stochastic Analysis. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, CA.
- OEHHA, 1999. Chloropicrin, Acute Toxicity Summary. In Technical Support Document for the Determination of Acute Reference Exposure Levels for Airborne Toxicants. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, CA.
- Orlowski, M., and A. Karkowsky, 1976. Glutathione metabolism and some possible functions of glutathione in the nervous system. *International Review of Neurobiology* 19:75-121.
- Oshima, R., 1995. Memorandum from R. Oshima to W. Thomas regarding chronic toxicity study by the inhalation route. February 17, 1995. Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Penkett, S.A., B.M.R. Jones, M.J. Rycroft, and D.A. Simmons, 1985. An interhemispheric comparison of the concentrations of bromine compounds in the atmosphere. *Nature* 318:550-553.
- Pickrell, S., S. Dwinell, and D. Tterlikkis, 1985. chemical residues in freshwater resulting from soil fumigation by methyl bromide and chloropicrin. Department of Environmental Regulation, Tallahassee, Florida. DPR Vol.123-095 #60019.
- Prentiss, A., 1937. *Chemicals in war*. New York: McGraw-Hill. p.143
- Price, N.R., 1985. The mode of action of fumigants. *J. Stored Product Research* 21:157-164.
- Putman, D.L. and M.J. Morris, 1990. Chromosome aberrations in Chinese hamster ovary (CHO) cells with confirmatory assay. Study # T9152.337001. Microbiological Associates, Inc. DPR Vol.199-041 #086983.
- Putman, D.L., and M.J. Morris, 1991. Micronucleus cytogenetic assay in mice. Microbiological Assoc. Inc. DPR Vol.123-136 #99090.
- Raabe, O.G., 1986. Inhalation uptake of selected chemical vapors at trace levels. University of California, Davis, CA. Submitted to The Biological Effects Research Section, California Air Resources Board, Sacramento, CA.
- Raabe, O.G., 1988. Retention and metabolism of toxics. Inhalation uptake of xenobiotic vapors by people. University of California, Davis, CA. Submitted to The Biological Effects Research Section, California Air Resources Board, Sacramento, CA.
- Radian Corp., 1988. Final report for the environmental fate studies of methyl bromide. Methyl Bromide Industry Panel. DPR Vol.123-105 #70489 and 123-106.

- Radian Corporation. 1992. Determination of Methyl Bromide Exposure During Dried Fruit and Tree Nut Fumigation Practices. Volume 1 and 2. Report prepared for the Dried Fruit Association of California.
- Rathus, E.M. and P.J. Landy, 1961. Methyl bromide poisoning. *British Journal of Industrial Medicine* 18:53-57.
- Reidy, T.J., J.F. Bolter, and J.E. Cone, 1994. Neuropsychological sequelae of methyl bromide: a case study. *Brain Injury*: 8(1):83-93.
- Reuzel, P.G.J., C.F. Kuper, H.C. Dreef-van der Meulen, and V.M.H. Hollanders, 1987. Chronic (29-month) inhalation toxicity and carcinogenicity study of methyl bromide in rats. Civo Institutes TNO. DPR Vol. 123-084 #59184; 123-147 #116337; 123-148 #120402; 123-148 #120406; and 123-166 #133417.
- Reuzel, R.G.J., H.C. Dreef-van der Meulen, V.M.H. Hollanders, C.F. Kuper, V.J. Feron, and C.A. van der Heijden, 1991. Chronic inhalation toxicity and carcinogenicity study of methyl bromide in Wistar rats. *Food and Chemical Toxicology* 29:31-39.
- Rodier, P.M., 1995. Developing brain as a target of toxicity. *Environmental Health Perspectives* 103(Suppl 6):73-76.
- Rodier, P.M. and Gramann, W.J., 1979. Morphologic effects of interference with cell proliferation in the early fetal period. *Neurobehavioral Toxicology* 1:129-135.
- Rosenblum, I., A.A. Stein, and G. Eisinger, 1960. Chronic ingestion by dogs of methyl bromide-fumigated food. *Archives of Environmental Health* 1:316-323. (in DPR Vol. 123-048 #913093).
- Ross, L.J., B. Johnson, K.D. Kim, and J. Hsu. 1996. Prediction of methyl bromide flux from area sources using the ISCST model. *Journal of Environmental Quality* 25:885891.
- Rounds, D.E., 1980. The effect of methyl bromide on the frequency of SCEs in CHO cells. Pasadena Foundation for Medical Research. DPR Vol.123-044 #35750.
- Russo, J.M., W.K. Anger, J.V. Setzer, and W.S. Brightwell, 1984. Neurobehavioral assessment of chronic low-level methyl bromide exposure in the rabbit. *Journal of Toxicology and Environmental Health* 14:247-255.
- San, R.H.C. and C.I. Sigler, 1990a. L5178Y TK +/- mouse lymphoma mutagenesis assay with confirmation. Study # T9152.701020. Microbiological Associates, Inc. DPR Vol.199-041 #086982.
- San, R.H.C. and C.I. Sigler, 1990b. Salmonella/mammalian-microsome plate incorporation mutagenicity assay (Ames test) with a confirmatory assay. Study # T9152.501014. Microbiological Associates, Inc. DPR Vol.199-046 #088717.

- Sanders, J.S., 1997a. Monitoring methyl bromide field fumigations during winter months. Memorandum, July 18, 1997 from Sanders to Paul Gosselin. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Sanders, J.S., 1997b. Methyl bromide sensitive site monitoring, Monterey County, July - August, 1997. Memorandum, August 21, 1997 from Sanders to Paul Gosselin. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Sayers, R.R., W.P. Yant, B.G.H. Thomas, and L.B. Berger, 1929. Physiological response attending exposure to vapors of methyl bromide, methyl chloride, ethyl bromide and ethyl chloride. Public Health Bulletin No. 185., U.S. Public Health Service.
- Scheffrahn, R.H., L. Bodalbhai, and N-Y. Su, 1992. Residues of methyl bromide and sulfuryl fluoride in manufacturer-packaged household foods following fumigation. Bulletin of Environmental Contamination and Toxicology 48:821-827.
- Schardein, J.L., 1993. Inhalation developmental toxicity study in rats. Project #: 656-007. IRDC. DPR Vol.199-051 #122503.
- Schardein, J.L., 1994. Two generation inhalation reproduction/fertility study in rats. Study ID#: 656-011. IRDC. DPR Vol.199-056 and -064 #132463 and #138326.
- Schreider, J., 1986. Comparative anatomy and function of the nasal passages. In Toxicology of the Nasal Passages (C. Burrow, Ed.), pp.1-25. Hemisphere Publishing Company, New York.
- Schumacher, D.K., 1985. The determination of methyl bromide in cocoa beans and cocoa products. Chocolate Manufacturers Association. DPR Vol.123-090 #59574.
- Schwob, J.E., S.L. Youngentob, and R.C. Mezza, 1995. Reconstitution of the rat olfactory epithelium after methyl bromide-induced lesion. Journal of Comparative Neurology 359:15-37.
- Segawa, R., T. Barry and C. Gana. 1992. Results of Off-Site Air Monitoring Following Methyl Bromide Chamber Fumigations and Evaluation of the ISCST Model. Memorandum to John Sanders dated September 3, 1992. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Segawa, R., T. Barry, H. Biermann, and P. Wofford. 1994a. Methyl Bromide Monitoring: Building Fumigations, Sutter County. Memorandum to John Sanders dated June 16, 1994. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.

- Segawa, R., T. Barry, and P. Wofford. 1994b. Methyl Bromide Monitoring: Building Fumigation, Stanislaus County. Memorandum to John Sanders dated August 26, 1994. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Segawa, R., T. Barry, P. Wofford, and C. Ando. 1994c. Methyl Bromide Monitoring: Building Fumigation, Madera County. Memorandum to John Sanders dated December 2, 1994. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Segawa, R., T. Barry, and P. Wofford. 1995. Methyl Bromide Monitoring: Building Fumigation, Yolo County. Memorandum to John Sanders dated April 6, 1995. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Segawa, R. and J.S. Sanders, 1997. Methyl bromide monitoring in Santa Cruz county. Memorandum, October 2, 1997 from Segawa and Sanders to James W. Wells. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Seiber, J.N., M.M. McChesney, J.E. Woodrow, and T.S. Shibamoto, 1987. Final Report to the Air Resources Board Pilot Analysis of Methyl Bromide in Air. Contract #A5-169-43. Department of Environmental Toxicology, University of California, Davis.
- Sell, C.R., N.G. Klag, and A.K. Burditt, 1988. Methyl bromide residues in fresh cherries: Effects of parameters in fumigation. *Pesticide Science* 23:41-49.
- Siemer, S.R. 1992. Methyl Bromide Industry Panel Interim Report. Siemer and Associates, Inc. Study Number TC199.
- Sikov, M.R., W.C. Cannon, D.B. Carr, R.A. Miller, L.F. Montgomery, and D.W. Phelps, 1981. Teratologic assessment of butylene oxide, styrene oxide and methyl bromide. Contract no. 210-78-0025. Battelle, Pacific Northwest Lab. Submitted to the Division of Biomedical and Behavioral Science, National Institute for Occupational Safety and Health, U.S. Department of Health and Human Services. DPR Vol.123-092 #59690 (same study also in DPR Vol. 123-039 #26865 and 26866).
- Simmon, V.F., K. Kauhanen, and R.G. Tardiff, 1977. Mutagenic activity of chemicals identified in drinking water. In Progress in Genetic Toxicology (Scott, D., B.A. Bridges, and F.H. Sobels, Ed.), pp. 249-258. Elsevier/North Holland Biomedical Press, Amsterdam. DPR Vol.123-109 #87801.
- Singh, H.B., L.J. Salas, A.J. Smith, and H. Shigeishi, 1981. Measurements of some potentially hazardous organic chemicals in urban environments. *Atmospheric Environment* 15:601-612.

- Singh, G., L.E. Rippon, and W.S. Gilbert, 1982. Methyl bromide and inorganic bromide residues in avocados after fumigation and storage. *Australian Journal of Experimental Agriculture and Animal Husbandry* 22:343-347.
- Slauter, R.W., 1995. Two year oral (gavage) chronic toxicity study of chloropicrin in rats. Study ID #: 656-003. DPR Vol.199-066 #138871.
- Smith, P., and W. Hambourger, 1935. Antipyretic and toxic effects of combinations of acetanilide with sodium bromide and with caffeine. *Journal of Pharmacology and Experimental Therapeutics* 55:200-205.
- Soil Chemicals Corp., 1990. Worker exposure data, 12" tarpless. DPR Vol.123-120 #91483.
- Squier, M.V., J. Thompson, and B. Rajgopalan, 1992. Case report: neuropathology of methyl bromide intoxication. *Neuropathology and Applied Neurobiology* 18:579-584.
- Stein, E.R. and D.A. Wolfenbarger, 1989. Methyl bromide residues in fumigated mango. *Journal of Agricultural and Food Chemistry* 37:1507-1509.
- Swenson, E.J., B.Z. Horowitz, T.E. Albertson, and M. O'Malley, 1997. An unusual exposure to methyl bromide leading to fatality. *Clinical Toxicology* 35(5):513.
- Tanaka, S., H. Arito, S. Abuku, and S. Imamiya, 1988. Acute effects of methyl bromide on electroencephalographic activity and sleep-wakefulness in rats. *Industrial Health* 26:101-114.
- Tanaka, S., S. Abuku, Y. Seki, and S. Imamiya, 1991. Evaluation of methyl bromide exposure on the plant quarantine fumigators by environmental and biological monitoring. *Industrial Health* 29:11-21.
- Tebbets, J.S., P.L. Hartsell, H.D. Nelson, and J.C. Tebbets, 1983. Methyl bromide fumigation of tree fruits for control of the Mediterranean fruit fly: Concentrations, sorption, and residues. *Journal of Agricultural and Food Chemistry* 31:247-249. (in DPR Vol.123-089 #59572 and 123-144 #116226).
- The Merck Index, 1989. Eleventh edition. (S. Budavari, M.J. O'Neil, A. Smith, and P.E. Heckelman, Ed.), Merck & Co., Inc., Rahway, N.J. 1606 pp.
- Tompkins, E.C. 1995. A four week dietary range-finding toxicity study of methyl bromide in rats. Project number WIL-49013. WIL Research Laboratories, Inc. Methyl Bromide Industry Panel. DPR Vol.123-172 #143942.
- Trical, Inc., 1985. Methyl bromide and chloropicrin pre-plant soil fumigation residue trial. Pomona, CA. DPR Vol.123-096 #59688.
- Trical, Inc., 1900. Carrot section 18 application. DPR Vol.123-117 #88782.

- Trical, Inc., 1991. Summary of methyl bromide pre-plant soil fumigation data, CA. DPR Vol. 123-131 #92703.
- Uncini, A., M. Basciani, A. Di Muzio, D. Antonini, and M. Onofri, 1990. Methyl bromide myoclonus: an electrophysiological study. *Acta Neurologica Scandinavica* 81:159-164.
- UNEP (United Nations Development Programme), 1997. Report of the Ninth Meeting of the Parties to the Montreal Protocol on Substances that Deplete the Ozone Layer. September 15-17, 1997 in Montreal, Canada. (<http://www.unep.ch/ozone/9mop.htm>)
- USDA, 1993. Methyl bromide substitutes and alternatives. A research agenda for the 1990s. U.S. Department of Agriculture, Agriculture Research Service, Beltsville, Maryland.
- U.S. EPA, 1986a. Pesticide tolerance petitions; Methyl Bromide Industry Panel. Federal Register 51(53):9512.
- U.S. EPA, 1986b. Guidance for the Registration of Pesticide Products Containing Methyl Bromide as the Active Ingredient. Office of Pesticides and Toxic Substances, US Environmental Protection Agency, Washington, D.C.
- U.S. EPA, 1988. Recommendations for and documentation of biological values for use in risk assessment. Table 1-3. National Technical Information Service PB 88-179874, U.S. Environmental Protection Agency, Cincinnati, OH.
- U.S. EPA, 1990. Review draft interim methods for development of inhalation reference concentrations. EPA/600/8-90/066A. August 1990. Office of Research and Development, U.S. Environmental Protection Agency, Washington, D.C.
- U.S. EPA, 1991. Guidelines for developmental toxicity risk assessment. Federal Register 56(234):63798-63826.
- U.S. EPA, 1992a. Bromomethane, 4/6/92, Integrated Risk Information System (IRIS), U.S. Environmental Protection Agency, Washington, D.C.
- U.S. EPA, 1992b. EPA, State of California and registrants agree to interim label changes for home fumigation uses of pesticide methyl bromide. Environmental News, Communications, Education, and Public Affairs, U.S. Environmental Protection Agency, Washington, D.C.
- U.S. EPA, 1992c. Guidelines for exposure assessment; Notice. Federal Register 57(104):22888-26021.
- U.S. EPA, 1993. Protection of the stratospheric ozone. Federal Register 58(236): 65018-65082.
- U.S. EPA, 1997a. EPA Methyl Bromide Phase Out Web Site. U.S. Environmental Protection Agency, Washington, D.C. (<http://www.epa.gov/docs/ozone/mbr/mbrqa.html#q2>)

- U.S. EPA, 1997b. The Federal Insecticide Fungicide, and Rodenticide Act (FIFRA) and Federal Food, Drug, and Cosmetic Act (FFDCA) as Amended by the Food Quality Protection Act (FQPA) of August 3, 1996. Document no. 730L97001, March 1997. Office of Pesticide Programs, U.S. Environmental Protection Agency, Washington, D.C.
- U.S. EPA, 1997c. 1996 Food Quality Protection Act Implementation Plan. March, 1977. Office of Prevention, Pesticides and Toxic Substances (7506C), U.S. Environmental Protection Agency, Washington, D.C. (<http://www.epa.gov/fedrgstr/>)
- U.S. EPA, 1998a. Guidelines for Neurotoxicity Risk Assessment. Federal Register 63(93):26926-26954.
- U.S. EPA, 1998b. Protection of stratospheric ozone control of methyl bromide emission through use of tarps. Federal Register 63(24):6008-6010.
- U.S. EPA, 1998c. Toxicology data requirements for assessing risks of pesticide exposure to children's health, draft. Presented to the Scientific Advisory Panel on December 8-9, 1998, Washington, D.C.
- U.S. EPA, 1999a. Toxicology data requirements for assessing risks of pesticide exposure to children's health, April 28, 1999 Draft. Report of the Toxicology Working Group of the 10X Task Force. U.S. Environmental Protection Agency, Washington, D.C.
- U.S. EPA, 1999b. The Office of Pesticide Programs' policy on determination of the appropriate FQPA safety factor(s) for use in the tolerance-setting process. Office of Pesticide Programs, U.S. Environmental Protection Agency, Washington, D.C.
- U.S. EPA, 1999c. Standard operating procedures for the Health Effects Division FQPA Safety Factor Committee, April 26, 1999. Presented at the May 1999 FIFRA Scientific Advisory Panel meeting. U.S. Environmental Protection Agency, Washington, D.C.
- U.S. Testing Co., Inc., 1976. Toxicological and skin corrosion testing of selected hazardous materials. PB-264 975. National Technical Information Service, U.S. Department of Commerce. DPR Vol.199-003 #913911 (also in Vol.199-017 #23381).
- Van Assche, C., 1971. Behaviour and perspectives of chemical soil fumigation. Proceedings of the 6th British Insecticide Fungicide Conference, November 15-18. Brighton, England.
- van Gelderen, C.E.M., T.J.F. Savelkoul, J.L. Blom, W. van Dokkum, and R. Kroes, 1993. The no-effect level of sodium bromide in healthy volunteers. Human and Experimental Toxicology 12(1): 9-14.
- von Oettingen, W.F., 1946. The toxicity and potential dangers of methyl bromide with special reference to its use in the chemical industry, in fire extinguishers, and in fumigation. Federal Security Agency, U.S. Public Health Service, National Institute of Health Bulletin, No. 185.

- Watrous, R.M., 1942. Methyl bromide. *Industrial Medicine* 11:575-579.
- Watson, R.T., D.L. Albritton, S.O. Andersen, and S. Lee-Bapty, 1992. Synthesis report of the methyl bromide interim scientific assessment and methyl bromide interim technology and economic assessment. Montreal Protocol Assessment Supplement. Requested by United Nations Environment Programme on behalf of the Contracting Parties to the Montreal Protocol. June 25, 1992.
- WHO (World Health Organization), 1995. Methyl bromide. *Environmental Health Criteria* 166. International Programme on Chemical Safety. Geneva, Switzerland.
- WIL, 1984. A residue study with commodities grown on soil treated with methyl bromide and chloropicrin. Projects WIL-48001 and WIL-49001. DPR Vol.123-095 (same as 123-102 #65281).
- WIL, 1985a. A hydrolysis study with methyl bromide. WIL Project 49003, WIL Research Lab., Inc. DPR Vol.123-086 #59495, same as 123-079 #52426.
- WIL, 1985b. A laboratory volatility study with methyl bromide and chloropicrin. WIL Project 78001, WIL Research Lab., Inc. DPR Vol.123-079 #52428.
- WIL, 1986. An adsorption study with soil and methyl bromide. WIL Project 49002, WIL Research Lab., Inc. DPR Vol.123-085 #59494.
- Winteringham, F.P.W., A. Harrison, R.G. Bridges, and P. Bridges, 1955. The fate of labeled insecticide residues in food products. II. The nature of methyl bromide residues in fumigated wheat. *Journal of the Science of Food and Agricultural* 6:251-261.
- Wisler, J.A., 1994. Evaluation of chloropicrin in a one year oral (capsule) toxicity study in dogs. Study #: 656-005. IRDC. DPR Vol. 199-055 #129614.
- Wofford, P. and K. Bennett. 1993. Results of Off-Site Air Monitoring During a Methyl Bromide Warehouse Fumigation. Memorandum to Kean Goh dated May 21, 1993. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Wofford, P. and R. Segawa. 1997. Methyl Bromide Monitoring: Chamber Fumigation, Winters, California. Memorandum to John Sanders dated February 6, 1997. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Wofford, P., and R. Segawa, 1998a. Monitoring results from a deep untarped application in Madera county - Method 2. Memorandum, March 20, 1998 from Wofford and Segawa to Douglas Y. Okumura. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.

- Wofford, P., and R. Segawa, 1998b. Monitoring results from a shallow tarped broadcast application in Monterey county -Method 4/5. Memorandum, March 20, 1998 from Wofford and Segawa to Douglas Y. Okumura. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Wofford, P., and R. Segawa, 1998c. Methyl bromide monitoring results from a bedded tarped application in Orange county -Method 9.1. Memorandum, January 27, 1998 from Wofford and Segawa to John S. Sanders. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Wofford, P., and R. Segawa, 1998d. Monitoring results from a bedded tarped applications in Riverside county - Method 10. Memorandum, January 27, 1998 from Wofford and Segawa to John S. Sanders. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Wofford, P. and R. Segawa. 1998e. Monitoring Results From a Nontarped Shallow Broadcast Application In Merced County - Method 1.1 Memorandum, May 18, 1998 from Wofford and Kim to Douglas Y. Okumura. Environmental Monitoring and Pest Management Branch, Department of Pesticide Regulation, California Environmental Protection Agency, Sacramento, CA.
- Wofsy, S., M. McElroy, and Y. Yung, 1975. The chemistry of atmospheric bromine. *Geophysics Research Letters* 3:215-218.
- Wyers, H., 1945. Methyl bromide intoxication. *British Journal of Industrial Medicine* 2:24-29.
- Yagi, K., J. Williams, N.-Y. Wang, and R.J. Cicerone, 1995. Atmospheric methyl bromide (CH₃Br) from agricultural soil fumigations. *Science* 267:1979-1981.
- Yates, S.R., J.Gan, F.F. Ernst, A. Mutziger, and M.V. Yates, 1996a. Methyl bromide emissions from a covered field: I. Experimental conditions and degradation in soil. *Journal of Environmental Quality* 25:184-192.
- Yates, S.R., F.F. Ernst, J.Gan, F. Gao, and M.V. Yates, 1996b. Methyl bromide emissions from a covered field: II. Volatilization. *Journal of Environmental Quality* 25:192-202.
- York, R.G., 1993. Inhalation developmental toxicity study in New Zealand white rabbits. Project #: 656-009. IRDC. DPR Vol.199-051 #122504.
- Yoshida, M., T. Ikeda, M. Iwasaki, S. Tsuda, and Y. Shirasu, 1987a. Acute inhalation toxicity of chloropicrin vapor in rats. *Journal of Pesticide Science* 12:237-244. (also in DPR Vol. 199-038 #090204).

Yoshida, M., T. Ikeda, M. Iwasaki, M. Ikeda, T. Harada, K. Ebino, S. Tsuda, and Y. Shirasu, 1987b. Subchronic inhalation toxicity of chloropicrin vapor in rats. *Journal of Pesticide Science* 12:673-681. (also in DPR Vol.199-038 #090205).

Zielhuis, R.L., and F.W. van der Kreek, 1979. The use of a safety factor in setting health based permissible levels for occupational exposure. *International Archives of Occupational and Environmental Health* 42:191-201.

Zwaveling, J.H., W.L.A.M. de Kort, J. Meulenbelt, M. Hezemans-Boer, W.A. van Vloten, and B. Sangster, 1987. Exposure of the skin to methyl bromide: A study of six cases occupationally exposed to high concentrations during fumigation. *Human Toxicology* 6:491-495.

VIII. APPENDICES