

SUPPLEMENTAL LABEL

GROUP	INSECTICIDE
--------------	--------------------

FujiMite® SC Miticide/Insecticide EPA Reg. No. 71711-4

For Use on Citrus Fruits in the States of AZ, CA, HI, ID, OR, WA

This supplemental label expires August 16, 2020 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for FujiMite SC Miticide/Insecticide before applying. Use of FujiMite SC Miticide/Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for FujiMite SC Miticide/Insecticide.

This supplemental labeling provides for a reduced Preharvest Interval (PHI) for citrus from 14 to 3 days.

APPLICATION RATE CHART

Citrus Fruits (Crop Group 10-10)		
Australian desert lime; Australian finger lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin, clementine); tangor; trifoliolate orange; uni fruit; cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Asian citrus psyllid ¹ Citrus leafminer* Citrus thrips* Leafhoppers Mealybugs Other Mites (see Target Species)	2.0 to 4.0 pints (0.105 to 0.210 lb ai)	USE RESTRICTIONS <ul style="list-style-type: none"> Apply by ground using a minimum of 100 gallons of water per acre. For full size trees, use a minimum of 200 gallons of water per acre. Do not apply by air. Do not apply through any type of irrigation system. Do not apply to citrus nurseries or citrus in greenhouses. Do not apply more than 8.0 pints (0.420 lb ai) per acre per growing season. Do not make more than 2 applications per growing season. Allow 14 days between applications. Preharvest Interval (PHI): 3 days
*suppression		
¹ For best results, use for control of adults and nymphs present at time of application when newly expanding foliage flush is present.		

©2017 Nichino America, Inc. FujiMite and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA[®]
Nichino America, Inc.
4550 Linden Hill Road
Wilmington, DE 19808
888-740-7700